

DIRECTIONS FOR USE

GENERAL—Apply with a low-pressure sprayer equipped with a coarse spray nozzle. Spray surfaces only until wet. Avoid excessive run-off. Apply to all insect breeding sites and harborages. Also treat adjacent exposed surfaces where insects may crawl when they come out of hiding. Repeat treatment as necessary.

Cockroaches—Thoroughly treat the infested area. Spray baseboards, cracks and crevices; around window and door frames; the under surfaces of shelves, drawers and work tables; behind and beneath cabinets, refrigerators, sinks and stoves; in and around waste containers, switch panels, junction boxes, floor drains and other places where cockroaches hide or rest.

Ants—Apply to trails and places where ants enter premises such as window and door sills. Spray hills and runways outdoors.

Silverfish, Firebrats and Paperlice—Apply to localized areas infested by these insects.

Brown Dog Ticks—Spray sleeping quarters of pets, around baseboards, window and door frames, wall cracks, and local areas of floors. Repeat as needed. Fresh bedding should be placed in animal quarters after spraying. Do not spray animals.

Clover Mites, Crickets, Earwigs, Millipedes, Centipedes, Sowbugs, Scorpions and Spiders—Thoroughly spray the interior and exterior perimeter of the infested premises. For Clover Mites give special attention to window sills.

Fleas, Flies, Gnats, Mosquitoes, Punkies, and Sandflies (outdoor only)—Spray infested areas thoroughly including outside surfaces of screens, doors, window frames, foundations, patios and other areas where insects may enter.

F 69

CENTOKILL

Residual Spray-Contains Baygon

For Industrial and Institutional Use Such as in Food Processing Plants, Breweries, Canneries, and Warehouses.

CENTOKILL is a highly effective ready to use formulation for control of insects listed below, including those resistant to certain chlorinated hydrocarbon and phosphate insecticides. CENTOKILL drives insects from their hiding places; gives rapid kill and long residual activity.

Active Ingredients . . . 85%: Petroleum Distillates, o-Isopropoxyphenyl methylcarbamate*, N-octyl bicycloheptene dicarboximide, N-octyl sulfoxide of isosafrole and related compounds, Pyrethrins.

Inert Ingredients . . . 15%: Isopropanol, Methylene Chloride.

*U.S. Patent No. 3,111,539. This material used under license from Ciemagro Corporation. Baygon is the registered trademark of Farbenfabriken Bayer A. G., Chemagro Corporation licensee.

KILLS: Ants, Brown Dog Ticks, Centipedes, Cockroaches, Clover Mites, Crickets, Earwigs, Firebrats, Fleas, Flies, Gnats, Millipedes, Mosquitoes, Paperlice, Punkies, Sandflies, Scorpions, Silverfish, Sowbugs, and Spiders.

CAUTION

Keep Out of Reach of Children

May be harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing spray mist. Do not get in eyes, on skin, or on clothing. In case of contact, wash immediately with soap and warm water. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Should illness occur, get prompt medical aid. To physician—Atropine sulfate is antidotal.

See side panel for additional cautions.

USDA Reg. No. 1537-105

AGRICULTURAL CHEMICALS DIVISION



CENTER

BRANCHES IN PRINCIPAL CITIES

CHEMICAL COMPANY

ATLANTA, GEORGIA

PRINTED IN U.S.A.

DIRECTIONS FOR USE

For Use in Meat Packing and Food Processing Plants: Use only in the inedible product areas of meat packing plants, including those that are federally inspected. Any exposed food products, equipment, or packaging materials must be removed or covered before spraying. Rinse all equipment with water before resuming operations. Treat food processing plants early in the season before canning or other processing begins. Repeat application, if necessary, when plant is not in operation. Spraying unloading docks and outside areas where insects congregate will aid in preventing re-infestation. Good sanitation and proper disposal of wastes is essential in any effective insect control program.

Insect Control in Research Facilities: The use of CENTOKILL in and around research animal quarters for insect control will not induce changes in liver microsomal enzyme activity in these animals.

CAUTION

Do not use as a space spray. Provide adequate ventilation of area being treated. Do not allow children to contact treated areas until surfaces are thoroughly dry. Do not store near feed or food products. Avoid contaminating feed, food, food processing equipment, and food handling surfaces. Do not apply to animals. Remove pets and cover fish bowls before spraying. Do not spray plants.

ISC-50

Net Contents _____ Gallons

