

ACCEPTED
with COMMENTS
by EPA Label Desk

APR 4 1985

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
and 40 CFR Part 152, the pesticide
has been registered by EPA and the
EPA Label Desk.

1543-10



Ready to use Insecticide
(Aqueous)

ACTIVE INGREDIENTS:

Pyrethrins.....0.10
Piperonyl Butoxide, Technical..1.00
INERT INGREDIENTS..... 90.90
100%

*Equivalent to min. 0.80 (butylcarbityl)
(6-propylpiperonyl) ether and 0.20
related compounds

CAUTION

Keep Out of Reach of Children
See Other Precautions on the Back Panel

MANUFACTURED BY W.F. YOUNG, INC.,
111 LYMAN STREET, SPRINGFIELD, MA 01103

U.S.A. W.F. YOUNG, INC.
Net 128 oz. (1 gal.)
3.785 Liters

(Bottom of container)
Lot No.

Effective, fast-acting, ready to use Insecticide.

Directions for use:

BARN & STALL SPRAY

For rapid control of: stable flies, house flies, gnats, mosquitoes, wasps,
horns. Apply with conventional sprayers or compressed air equipment.

Indoor Application: Remove animals from the barn before spraying and ventilate before
reoccupying. For maximum effectiveness, a combination of surface treatment and fogging
is recommended.

Space Spraying: Close doors and windows and shut off ventilating equipment.
Apply with sprayer adjusted to deliver a fine mist or fog. Direct spray at an
upward angle distributing it uniformly throughout the entire area at a rate of
3-4 ounces per 1000 cubic feet of space.

Fogging: To kill exposed or accessible stages of the above named insects, use
a rate of 1 gallon to 20,000 to 40,000 cubic feet of space. Contact insects
directly whenever possible.

Outdoor Application: For adult mosquito control, apply as a fine mist to thoroughly
treat the entire area. For best results, treat when the wind velocity does not
exceed 3 miles per hour. Treat shrubbery and vegetation where mosquitoes may
rest. Apply at a rate of 0.1 ounce per 1000 square feet.

HORSE AND LIVESTOCK SPRAY

Effective, fast acting, ready to use Insecticide. Apply with conventional
sprayers or a cloth.

To protect cattle and horses from horn flies, house flies, mosquitoes
and gnats, apply a light mist sufficient to wet the surface of the hair.

To control stable flies, horse flies, and deer flies, apply at a rate of
2-4 ounces per adult animal sufficient to wet the hair thoroughly. Repeat
treatment daily or at intervals necessary to give overhead protection.

To control face flies, apply with a cloth dampened with the spray.

To control blood-sucking lice, apply to the infested areas of the animal.
Use a stiff brush to get the spray to the base of the hair. Repeat every
2-3 weeks if required.

PET INSECTICIDE

For control of fleas and brown dog ticks on premises and on pets. Effective,
fast-acting insecticide.

Surface spraying: Thoroughly spray infested areas, pet beds, resting quarters,
nearby cracks and crevices, along and behind baseboards, mouldings, window and
door frames and entire areas of floor covering. Fresh bedding should be placed
in animal quarters following treatment. Repeat treatment as needed.

On Pets: Start spraying at the tail, moving the dispenser rapidly and making sure
the animal's entire body is covered including the legs and under body. While
spraying, fluff the hair so spray will penetrate to the skin. Do not spray into
eyes or face. Avoid contact with genitalia. Repeat as needed.

HOUSEHOLD SPRAY

Effective fast acting ready to use spray for use
in homes. Leaves no oily or objectionable residues.

Indoor Application: Use of this product in food
processing areas should be limited to periods
when the kitchen is not in operation. Food should
be removed or covered during treatment. All food
processing surfaces should be covered during treat-
ment or thoroughly cleaned before use. Vacate areas
and ventilate before reoccupying.

For maximum results, a combination of spot treatment
and surface spray is recommended.

Surface spraying: To control cockroaches, palmetto
bugs, ants, silverfish, spiders, crickets, clover
mites, cheese mites, granary weevils, confused flour
beetles, spider beetles, cigarette beetles, drugstore
beetles, meal worms, grain mites and cadellies. Adjust
sprayer to a coarse wet spray. Direct the spray into
hiding places, cracks and crevices, around containers
of stored foods, behind shelves and drawers. For
silverfish, spray bookcases. For ants, spray trails,
nests, and points of entry. If surface application
only is to be used, spray floors and other surfaces
at the rate of 1 gallon to 750 square feet of area.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS &
DOMESTIC ANIMALS**

CAUTION: Harmful if swallowed. Avoid contamination of
feed and foodstuffs. Avoid contact with eyes. In case of
contact immediately flush eyes with plenty of water. Get
medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or
ponds. Do not apply where runoff is likely to occur. Do not
contaminate water by cleaning of equipment or disposal of
wastes. Apply this product only as specified on this label.

STORAGE AND DISPOSAL

STORAGE: DO NOT TRANSPORT OR STORE UNDER 32° F. Store in
original container only in a locked storage area. Leaking
packages of this product may be disposed of on site or at
an approved waste disposal facility.

DISPOSAL: Do not contaminate water, food, or feed by storage
or disposal. Wastes resulting from the use of this product
may be disposed of on site or at an approved waste disposal
facility. Triple rinse (or equivalent). Then offer for
recycling or reconditioning, or puncture and dispose of in
a sanitary landfill, or incineration, or, if allowed by
state and local authorities, by burning. If burned, stay
out of smoke.

Buyer assumes all risks of use, storage or handling of this
material not in strict accordance with directions given
herewith.

It is a violation of Federal Law to use this product in a
manner inconsistent with its labeling.

Establishment No. 1543-99-1
E.P.A. Reg. No. 1543-

Printed in U.S.A.