

ABSORBINE® BARN & STALL SPRAY



Ready To Use Insecticide
(Aqueous)

ACTIVE INGREDIENTS:

Pyrethrins	0.1%
*Piperonyl Butoxide, Technical	1.0%
INERT INGREDIENTS	98.9%
	100%

*Equivalent to min 0.8% (butylcarbityl) (6-propylpiperonyl) ether and 0.2% related compounds

CAUTION

Keep Out of Reach of Children
See Other Precautions on the Back Panel

Net 128 Fl oz. (1 Gal.) 3.785 Liters

EFFECTIVE, FAST-ACTING, READY TO USE INSECTICIDE.

DIRECTIONS FOR USE:

BARN & STALL SPRAY

For rapid control of: stable flies, house flies, gnats, mosquitoes, wasps, hornets. Apply with conventional sprayers or compressed air equipment.

Indoor Application: Remove animals from the barn before spraying and ventilate before reoccupying. For maximum effectiveness, a combination of surface treatment and fogging is recommended.

Space Spraying: Close doors and windows and shut off ventilating equipment. Apply with sprayer adjusted to deliver a fine mist or fog. Direct spray at an upward angle distributing it uniformly throughout the entire area at a rate of 3.6 ounces per 1000 cubic feet of space.

Fogging: To kill exposed or accessible stages of the above named insects, use a rate of 1 gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly whenever possible.

Outdoor Application: For adult mosquito control, apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery and vegetation where mosquitoes may rest. Apply at a rate of 0.1 ounce per 1000 square feet.

HORSE AND LIVESTOCK SPRAY

Effective, fast acting, ready to use insecticide. Apply with conventional sprayers or a cloth.

To protect cattle and horses from horn flies, house flies, mosquitoes and gnats, apply a light mist sufficient to wet the surface of the hair.

To control stable flies, horse flies, and deer flies, apply at a rate of 2.4 ounces per adult animal sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control face flies, apply with a cloth dampened with the spray.

To control blood-sucking lice, apply to the infested areas of the animal. Use a stiff brush to get the spray to the base of the hair. Repeat every 2-3 weeks if required.

PET INSECTICIDE

For control of fleas and brown dog ticks on premises and on pets. Effective, fast acting insecticide.

Surface spraying: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames and entire areas of floor covering. Fresh bedding should be placed in animal quarters following treatment. Repeat treatment as needed.

On Pets: Start spraying at the tail, moving the dispenser rapidly and making sure the animal's entire body is covered including the legs and under body. While spraying, fluff the hair so spray will penetrate to the skin. Do not spray into eyes or face. Avoid contact with genitalia. Repeat as needed.

HOUSEHOLD SPRAY

Effective fast acting ready to use spray for use in homes. Leaves no oily or objectionable residues.

Indoor Application: Use of this product in food processing areas should be limited to periods when the kitchen is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before use. Vacate areas and ventilate before reoccupying.

For maximum results, a combination of spot treatment and surface spray is recommended.

Surface spraying: To control cockroaches, palmetto bugs, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, confused flour beetles, spider beetles, cigarette beetles, drugstore beetles, meal worms, grain mites and cadellids. Adjust sprayer to a coarse wet spray. Direct the spray into hiding places, cracks and crevices, around containers of stored foods, behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests, and points of entry. If surface application only is to be used, spray floors and other surfaces at the rate of 1 gallon to 750 square feet of area.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Avoid contamination of feed and foodstuffs. Avoid contact with eyes. In case of contact immediately flush eyes with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

STORAGE AND DISPOSAL

STORAGE: DO NOT TRANSPORT OR STORE UNDER 32° F. Store in original container only in a locked storage area. Leaking packages of this product may be disposed of on site or at an approved waste disposal facility.

DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith. It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Establishment No. 1543 MA-1
E.P.A. Reg. No. 1543 ID

MANUFACTURED BY W. F. YOUNG, INC., 111 LYMAN STREET, SPRINGFIELD, MA 01103, U.S.A.

© W. F. Young, Inc.

