

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DILUTION INSTRUCTIONS

For an emulsion containing 0.35% IMPERIAL I dilute

Concentrate	To Volume Below With Water
1 1/2	8 fluid ounces
3	16 fluid ounces
6	32 fluid ounces
11 1/2	1/2 gallon
23	1 gallon

MIX WELL BEFORE USING.

APPLICATIONS:

INSTRUCTIONS FOR CONTACT SPACE AND RESIDUAL SPRAYS

FOR INDOOR USE: In non-food areas of food handling establishments, commercial, industrial, office, residential, public, institutional and storage buildings, homes, hotels, ship cabins and hospitals. In hospitals patients should be removed prior to treatment. Also for use in buses, automobiles, and non-food areas of trucks, boxcars and ship cargo holds. Do not use on electrical equipment or on surfaces that can be harmed or stained by water. Also for use indoors in dog kennels and horse stables. In horse stables, not for application if horses will be used for food.

CONTACT SPRAY—KILLS AND REDUCES INFESTATIONS: COCKROACHES, CRICKETS, SILVERFISH, ANTS, CARPET BEETLES, CENTIPEDES, EARWIGS, FLOUR BEETLES, GRAM MITES, HELLGRIPPERS, SCORPIONS, SPIDERS: Apply as a contact spray. Fill spray tank with IMPERIAL I 0.35% Transparent Emulsion Spray formulation. Keep spray nozzle about 8 to 12 inches from surface to be treated. Adjust pressure and nozzle to deliver a coarse wet spray. When possible, spray directly on pests to reduce infestations. Spray thoroughly into hiding places such as cracks, crevices, most areas, openings around pipes and sinks, under refrigerators and other appliances, baseboards and storage areas. Apply spray until area is moist, but not to point of runoff. Spray directly on insects where possible. Repeat as necessary.

ANTS: To kill ants. Spray trails, nests and points of entry within buildings and homes. Spray on ants where possible. Repeat as necessary.

FLEAS: Aids in the control of fleas on dogs and cats, on pet bedding and in horse stalls. Apply directly to animals along back, under forelegs and on other areas. Part hair to contact skin and insure good coverage. Hold nozzle about one foot away while spraying. Apply until hair is thoroughly moist. Do not directly spray in eyes, or mouth of animal. To reduce reinestation, thoroughly spray the animal's bedding and other resting areas. Treat those areas at the rate of two seconds per linear foot while spraying at a distance of 12 to 18 inches. Also spray entire inside surface of dog houses. Repeat as necessary.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

WARNING

Harmful if swallowed. Avoid contact with skin or clothing. Causes eye irritation. Do not get in eyes. Wash after handling. Avoid breathing vapor. Do not spray on humans. Do not spray where food or drink is exposed or being consumed.

ENVIRONMENTAL HAZARDS

Run off of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. When used on ships, do not wash or flush freshly sprayed surfaces into adjacent waters.

FIRST AID

If swallowed: Induce vomiting by sticking finger in back of throat or by administration of syrup of ipecac. Get medical attention. If in eyes, flush with water for 15 minutes and get medical attention. For skin, wash with plenty of soap and water. If irritation develops, get medical attention.

PHYSICAL OR CHEMICAL HAZARDS

DO NOT USE. POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. FLASH POINT MINIMUM 140° F.

IMPERIAL I 2% TRANSPARENT EMULSION** CONCENTRATE

A SYNTHETIC PYRETHROID

DUAL ACTION INSECT KILLER

KILLS AS CONTACT AND RESIDUAL SPRAY

KILLS ROACHES UP TO 4 WEEKS

ACTIVE INGREDIENTS:

resmethrin (15-(phenylmethyl)-3-(4-phenyl)-2,2-dimethyl-3-(2-methyl-1-propenyl) cyclopropanecarboxylate)	2.000% w/w
Related compounds	0.274%
INERT INGREDIENTS:	97.726%
	100.00% w/w

cis/trans isomers ratio: max. 30% (cis) and min. 70% (trans)
 *U.S. Patent Nos. 3,465,007 and 3,542,928
 Penick's SBP-1382* brand of resmethrin insecticide
 Penick Corp. 1050 Wall Street West, Lyndhurst, N.J. 07071
 **U.S. Patent No. 3,683,078

Contains 0.17 pounds resmethrin per gallon
 EPA REG. NO. 1853-112
 EPA EST. NO. 1553-6A-1

KEEP OUT OF REACH OF CHILDREN

WARNING

HAVE CLEAN WATER SUPPLY AT HAND IN CASE OF AN EYE EMERGENCY WHEN DILUTING THIS CONCENTRATE. SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS. CAN BE USED IN NON-FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS.

MANUFACTURED BY

MOMAR
 INCORPORATED
 ATLANTA, GEORGIA, U.S.A.

Providing Products of Momar Research in the U.S., Canada, Western Europe, and Africa
NET CONTENTS:

TO KILL BEDBUGS: A bedboards and walls that may harbor bedbug infestations. Consult pest control professional.
RESIDUAL SPRAY: A CENTIPEDES, EARWIGS, baseboards, cracks and closets, in back and under spray hiding places in 4 to 6 inch band until.
FRUIT FLIES, HOUSEFLIES: and nozzle to deliver a mist to all parts of the room. The spray nozzle at all mist at the dosage rate room closed for at least 24 hours.
OUTDOORS—To control: spray. Spray directly on ants and fleas: See wasps, yellow jackets at rest. Spray liberally as necessary.
CENTIPEDES, EARWIGS: indoor use.

— STOP
 — DO NOT
 BY S
 — OFF
 — DO NOT

Pesticide Disposal: chemically or by incineration. Container Disposal: rinse (or equivalent) and approved for disposal authority.

IMPERIAL I 2% TRANSPARENT EMULSION** CONCENTRATE A SYNTHETIC PYRETHROID DUAL ACTION INSECT KILLER KILLS AS CONTACT AND RESIDUAL SPRAY KILLS ROACHES UP TO 4 WEEKS

ACTIVE INGREDIENTS:	
resmethrin ((5-(phenylmethyl)-3-furyl) methyl 2,2-dimethyl-3-(2-methyl-1-propenyl) cyclopropanecarboxylate)	2.000% w/w
related compounds	0.274%
INERT INGREDIENTS:	97.726%
	100.000% w/w

cis/trans isomers ratio: max. 30% (cis) and min. 70% (trans)
 *U.S. Patent Nos. 3,465,007 and 3,542,928
 Penick's SBP-1382® brand of resmethrin insecticide
 Penick Corp. 1050 Wall Street West, Lyndhurst, N.J. 07071
 **U.S. Patent No. 3,683,078

Contains 0.17 pounds resmethrin per gallon
 EPA REG. NO. 1853-112
 EPA EST. NO. 1553-GA 1

KEEP OUT OF REACH OF CHILDREN WARNING

HAVE CLEAN WATER SUPPLY AT HAND IN CASE OF
 AN EYE EMERGENCY WHEN DILUTING THIS CONCENTRATE.
 SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.
 CAN BE USED IN NON-FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS

MANUFACTURED BY

MOMAR

INCORPORATED
 ATLANTA, GEORGIA, U.S.A.

Providing Products of Momar Research in the U.S., Canada, Western Europe, and Africa
NET CONTENTS:

TO KILL BEDBUGS: Apply as a spot treatment to cracks and crevices around baseboards, floorboards, bedboards and walls. First, thoroughly clean and air mattresses and springs. Then spot treat only areas that may harbor bedbugs, concentrating on tufts, folds and seams. Repeat as necessary. For persistent infestations, consult professional applicator.

RESIDUAL SPRAY: Aids in the control of COCKROACHES, CRICKETS, SILVERFISH, ANTS, CENTIPEDES, EARWIGS, GRAIN MITES, SOWBUGS, SPIDERS. Kills for up to 4 weeks. Spray baseboards, cracks and crevices, around door and window frames, and other hiding places such as closets, in back and under appliances, sinks and cupboards in kitchens and bathrooms. In basements, spray hiding places in cracks and crevices, lower walls and dark places. Apply as a coarse wet spray in a spray hiding places in cracks and crevices, lower walls and dark places. Apply as a coarse wet spray in a 4 to 6 inch band until area is thoroughly wet but not to the point of run-off. Repeat as necessary.

FRUIT FLIES, HOUSEFLIES, MOSQUITOES, MOTHS, GNATS: Apply as a space spray. Adjust pressure and nozzle to deliver a fine mist spray. Close all doors and windows. Point nozzle upward, direct spray mist to all parts of the room, especially windows and other light sources which attract these insects. Keep the spray nozzle at all times at least four feet from fabrics to avoid wetting or staining. Fill the room with mist at the dosage rate of one fluid ounce per 1,000 cu. ft. of space. Then leave the treated area. Keep the room closed for at least 15 minutes. Ventilate the room when treatment is completed and before re-entry.

OUTDOORS—To control nuisance pests around buildings, patios and paved areas. Apply as a contact spray. Spray directly on insects. Avoid spraying turf or plants.

ANTS AND FLEAS: Same as directed for indoor use.

WASPS, YELLOW JACKETS, HORNETS: Application should be made in late evening when insects are at rest. Spray liberally into hiding and breeding places contacting as many insects as possible. Repeat as necessary.

CENTIPEDES, EARWIGS, MILLIPEDES, SOWBUGS, SPIDERS: Apply as a contact spray, as for indoor use.

STORAGE AND DISPOSAL

- STORE AT TEMPERATURES ABOVE 32° F.
- DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL
- OPEN DUMPING IS PROHIBITED
- DO NOT REUSE EMPTY CONTAINER

Pesticide Disposal—Pesticide spray mixture or rinsate that cannot be used or chemically processed should be disposed of in a landfill approved for pesticide or buried in a safe place away from water supplies.

Container Disposal—Reseal container and offer for reconditioning, or triple rinse (or equivalent) and offer for recycling, reconditioning, or disposal in approved landfill, or bury in a safe place. Consult federal, state or local disposal authorities for approved alternate procedures.