

Reg # 1553-131

PM-10

191

BEST AVAILABLE COPY

Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.

1/25/90

CONTRACT



15 FOOT STREAM SPRAY REACHES OVERHEAD NESTS, NIVES, PLUS UNDERGROUND LAIRS

DIELECTRIC STRENGTH NON-CONDUCTING UP TO 26,600 VOLTS

A SYNTHETIC PYRETHROID FOR CONTROLLING THE FOLLOWING FLYING AND CRAWLING INSECTS WASPS HORNETS YELLOW JACKETS BEES ANTS COCKROACHES SPIDERS EARWIGS GRASS BITES SOUBUGS CRICKETS BEETLES SILVERFISH CENTIPEDES MILLIPEDES AND TENT CATERPILLARS

ACTIVE INGREDIENTS:

Permethrin (8-phenoxy-methyl-3-hydroxy-2-methyl-2-dimethyl-3-(2-methyl-1-propoxy)-cyclopropanecarboxylate)	0.250% w/w
Related Compounds	0.034%
INERT INGREDIENTS	99.716%
TOTAL	100.000% w/w

*Can treat roach infestations 20% (1-2) and can 10% (1-2) from U.S. Patent Nos. 3,486,987 and 3,542,888
Contract # 1283 brand of contractum insecticide
Peston Care 1000 West Street West, Lyndhurst, NJ 07031

KEEP OUT OF REACH OF CHILDREN.

CAUTION: SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.
NET WEIGHT 14 OZ

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS CAUTION

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Cover or remove exposed foods. Avoid contamination of cooking utensils and food preparation surfaces. Remove, seal and cover fish aquariums before spraying.

FIRST AID

In case of contact, immediately flush skin and eyes with plenty of water for at least 15 minutes. Get medical attention.

PHYSICAL AND CHEMICAL HAZARDS

Contents under pressure. Do not puncture. DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME. Exposure to temperatures above 130°F may cause bursting. Never throw container into fire or incinerator.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. See side panel for additional instructions.

For use in and around homes, hotels, greenhouses and laboratories, food handling establishments, packaging and processing plants, seed houses, commercial, industrial office, institutional and storage buildings, dairies, milk storage rooms, egg processing plants, supermarkets, restaurants, bottling plants, canneries, bars and food mills, bakeries, boats, gulf cabins and hospitals. In hospitals patients should be removed prior to treatment. Also for use in buses, automobiles, trucks, boats and camp sites. Also for use in dog kennels and horse stables. Not for application if horses will be used for food.

In food handling areas, food should be removed or covered previous to treatment. All food processing surfaces should be cleaned or thoroughly cleaned before using. Apply only when insects are not in operation.

DISPOSAL: DO NOT REUSE EMPTY CONTAINER, WRAP CONTAINER AND PUT IN TRASH COLLECTION

EPA REG NO 1553-131

EPA EST NO. 10007-8A-1

DIRECTIONS FOR USE (Continued)

Remove protective cap. Aim spray opening away from person. Push button to spray.

For industrial use, indoors and around premises.

INDOORS TO KILL WASPS, HORNETS, YELLOW JACKETS, BEES—Application should be made late in the evening when insects are at rest. Spray liberally into hiding and breeding places, particularly under rafters, contacting as many insects as possible. Saturate cracks in and around nest openings. Repeat as necessary.

TO KILL ANTS, COCKROACHES, SPIDERS, EARWIGS, GRASS BITES, SOUBUGS, CRICKETS, BEETLES, SILVERFISH, CENTIPEDES AND MILLIPEDES—Direct spray to contact these insects in cracks and crevices in floors, baseboards, ground tops, behind cabinets and other harborage. Spray these infested areas until thoroughly wet. Repeat as necessary.

TO KILL ANTS—Spray directly on insects, and around window frames, doors, and trails and hills.

TO KILL SPIDERS—Spray directly on these pests and use repeated applications to the webbing and all cracks and crevices which harbor spiders.

OUTDOORS—Apply as a contact spray. Spray directly on insects. Aim spray opening away from person. Avoid spraying plants or turf except to spray infested areas.

ANTS—Same as directed above for indoor use. **WASPS, HORNETS, YELLOW JACKETS, BEES**—Same as directed for indoor use. Aim spray at nest openings in trees, bushes, under eaves, chimneys, cracks and crevices to saturate in and around the nest openings. Contact as many insects as possible. Repeat as necessary.

EARWIGS, SOUBUGS, CRICKETS, BEETLES, SILVERFISH, CENTIPEDES AND MILLIPEDES—Apply as a contact spray for indoor use.

TO KILL TENT CATERPILLARS—Application should be made when caterpillars are young and tents are first noticed. For best results, apply in the late afternoon or evening when caterpillars have started to show signs. Aim spray directly at tents and tent openings on all sides. Repeat as necessary.

Manufactured by MOMAR Incorporated
P.O. Box 19667 Atlanta, GA 30330
Pounding Products of Momar Research in the USA Canada Western Europe and Africa