

**ACCURATE - CONVENIENT
EFFECTIVE**

This product fulfills the criteria of Appendix F of the Grade 'A' Pasteurized Milk Ordinance 1965 Recommendations of the U. S. Public Health Service in waters up to 500 ppm of hardness calculated as CaCO₃ when tested by the AOAC Germicidal and Detergent Sanitizers-Official Method.

DANGER: Keep out of reach of children. Tablets may be harmful or fatal if swallowed. May cause skin irritation or eye damage. Avoid prolonged skin contact. Do not get in eyes. In case of contact flush with plenty of water. If irritation persists, get medical attention. Avoid contamination of food.

FIRST AID: If tablets are swallowed, drink promptly a large quantity of milk, egg whites or gelatin solution. Avoid alcohol.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

**ALWAYS FOLLOW YOUR
HEALTH DEPARTMENT
REGULATIONS**

S-9-172



**For the SANITIZATION of
DISHES-GLASSES-UTENSILS**

EPA REG. NO. 1561-9

ACTIVE INGREDIENTS 50%
Alkyl (C14 95%, C12 3%, C16 2%) dimethyl benzyl
ammonium chloride dihydrate
INERT INGREDIENTS 50%

DANGER
Keep Out of Reach of Children
**See Left Panel For Additional
Precautionary Statements**

200 TABLETS

NET WEIGHT 8 OUNCES

Manufacturing Chemists

EDWARDS-COUNCILOR CO.
NORFOLK, VIRGINIA 23510

DIRECTIONS

When preparing sanitizing solutions, allow several minutes for tablets to dissolve in water, stir well before using.

For sanitizing Dishes, Glasses, and, Utensils in Restaurants, Taverns, etc.:

1. Scrape and prewash utensils and glasses whenever possible.
2. Wash with a good detergent or compatible cleaner in first sink compartment.
3. Rinse with clean water in second sink compartment.
4. Sanitize in a solution of 1 tablet per 3 quarts of water (200 ppm) in third sink compartment. Immerse all utensils for at least two minutes or for contact time specified by governing sanitary code.
5. Place sanitized utensils on a rack or drainboard to air dry.

Steramine sanitizing solutions of 200 ppm concentration may be safely used on food processing equipment and utensils and on other food contact articles as specified under Subpart F of the Food Additives Amendment by the U. S. Food and Drug Administration.

2-14-72
1561-9