

Proposed additional label claims are shown in typewritten, yellow highlighted text.

This sanitizing product is registered with the U.S. Environmental Protection Agency as required by Federal law to insure that when used as directed, it has proven to be effective as claimed without causing unreasonable adverse effects to humans or the environment.

Steramine sanitizing solutions of 200 ppm concentration may be safely used on food processing equipment and utensils and on other food contact articles as specified under Subpart F of the Food Additives Amendment by the U.S. Food and Drug Administration.

This product meets the sanitizing criteria of the U.S. PUBLIC HEALTH SERVICE in waters up to 500 ppm of hardness calculated as CaCO₃ when tested by the AOAC Germicidal and Detergent Sanitizers - Official Methods PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING Keep out of Reach of Children Tablets dust causes eye damage and skin irritation. Do not get dust in eyes, on skin, or on clothing. Avoid prolonged skin contact. In case of contact, flush with plenty of water. If irritation persists, get medical attention. Avoid contamination of food.

STATEMENT OF PRACTICAL TREATMENT If tablets are swallowed, drink promptly large quantities of water. Avoid alcohol. Get medical attention immediately.

NOTE TO PHYSICIAN Probable no real damage may corroborate the use of epinephrine. STORAGE DISPOSAL Do not reuse empty container. Wrap container and put in trash.

S-10-1282

STERAMINE 20 TABLETS THE MOST EFFECTIVE SANITIZER

USE 1 TABLET PER TWO GALLONS OF WATER
For the **SANITIZATION** of
DISHES-GLASSES-UTENSILS
and other food contact surfaces

KILLS HIV-1 (AIDS Virus)
when used as directed for sanitizing

EPA REG. NO. 1981-14
ACTIVE INGREDIENTS 80%
Allyl (C14 95%, C12 3%, C16 2%) dimethyl benzyl ammonium chloride hydrate
INERT INGREDIENTS 20%

WARNING
Keep Out of Reach of Children
See Left Panel For Additional
Precautionary Statements
For Institutional and Commercial Use Only
75 TABLETS NET WEIGHT 8 OUNCE

Manufacturing Chemists
EDWARDS-COUNCILOR CO.
1427 BAKER ROAD, VA. BEACH, VA 23455

ACCEPTED
JAN 31 1994
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 1561-10

DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Prepare sanitizing solutions with warm water. Allow several minutes for tablets to dissolve before using.

For sanitizing Dishes, Glasses, and Utensils in Restaurants, Taverns, etc.

1. Scrub and pre-rinse utensils and glasses whenever possible.
2. Wash with a good detergent or commercial cleaner in first sink compartment.
3. Rinse with clean water in second sink compartment.
4. Soak in a solution of 1 tablet per 2 gallons of water (200 ppm) in third sink compartment. Immerse all utensils for at least one minute or for contact time specified by governing sanitary code.
5. Place sanitized utensils on a rack or drain-board to air dry.
6. A fresh sanitizing solution must be prepared at least daily or more often if the solution becomes diluted or soiled.

For Sanitizing Food Processing Equipment, Dairy Equipment, Sinks, Countertops, Refrigerated storage and Display Equipment and other hard non-porous food contact articles and surfaces.

1. Wash and Rinse all articles and surfaces thoroughly.
2. Apply a solution (Spray On or Wipe On) of 200 ppm concentration and allow surface to remain wet for at least one minute followed by adequate draining and air drying.

KILLS HIV-1 (AIDS Virus) when used as directed for sanitizing.

EPA REG. NO. 1981-14

BEST AVAILABLE COPY