

This product fulfills the sanitizing criteria of the US PUBLIC HEALTH SERVICE in waters up to 500 ppm of hardness calculated as CaCO3 when tested by the AOAC-Germicidal and Detergent Sanitizers - Official Method.

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS and DOMESTIC ANIMALS**  
**WARNING:** Tablet dust causes substantial but temporary eye injury. Do not get dust in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Tablets may be harmful if swallowed.

**FIRST AID**

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advise.  
**IF ON SKIN:** Rinse immediately with water for 15-20 minutes. If irritation persists, call poison control center or doctor for treatment advise.  
**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advise. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by poison control center doctor. Do not give anything by mouth to an unconscious person. Have the product label or container with you when calling a poison control center or doctor, or going for treatment.

**STORAGE and DISPOSAL:** Store in original container in areas inaccessible to persons unfamiliar with it's use. Do not reuse empty container. Wrap container and discard in trash.



**FOR SANITIZING FOOD CONTACT SURFACES**

USE ONE TABLET PER TWO GALLONS OF WATER

Kills HIV-1 (AIDS Virus) when used as directed for sanitizing.

EPA REG. NO. 1561-10

Active Ingredient (Quaternary).....50%  
 Alkyl (C14 95%, C12 3%, C16 2%) dimethyl benzyl ammonium chloride dihydrate  
 INERT INGREDIENTS.....50%

**WARNING**

**KEEP OUT OF REACH OF CHILDREN**

See left panel for additional precautionary statements. NET WEIGHT 8 OUNCES  
 For Institutional and Commercial use only.

75  
Tablets

Manufacturing Chemists  
 EDWARDS-COUNCILOR CO. INC., 1427 Baker Road, Virginia Beach, VA 23455

**DIRECTIONS**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Prepare sanitizing solutions with warm water. Allow several minutes for tablets to dissolve before using. **FOR SANITIZING DISHES, GLASSES, AND UTENSILS IN RESTAURANTS, TAVERNS, ETC.**

1. Scrape and prewash utensils and glasses whenever possible.
2. Wash with a good detergent or compatible cleaner in first sink compartment.
3. Rinse with clean water in second sink compartment.
4. Sanitize in a solution of 1 TABLET PER 2 GALLONS OF WATER (200 ppm) in third sink compartment. Immerse all utensils for at least one minute or for contact time specified by governing sanitary code.
5. Place sanitized utensils on a rack or drainboard to air dry.
6. A fresh sanitizing solution must be prepared at least daily or more often if the solution becomes diluted or soiled.

**DIRECTIONS FOR SPRAYING**

**FOR SANITIZING FOOD PROCESSING EQUIPMENT, DAIRY EQUIPMENT, SINKS, COUNTERTOPS, REFRIGERATED STORAGE AND DISPLAY EQUIPMENT** and other hard non-porous food contact articles and surfaces.




1. Wash and rinse all articles and surfaces thoroughly.
2. Apply a solution (SPRAY ON WITH HAND TRIGGER SPRAYER OR WIPE ON) of 200 ppm concentration and allow surface to remain wet for at least one minute followed by adequate draining and air drying.

KILLS HIV-1 (AIDS Virus) when used as directed for sanitizing.

EPA EST. 1561-VA-1

**DEODORIZATION**

For controlling odor causing bacteria in ostomy, colostomy, and ileostomy appliance pouches. Use 1 or 2 tablets in a clean ostomy, colostomy or ileostomy appliance pouch. Replace tablets after each pouch emptying.



1561-10

3/4/2005

Page 181