

The sanitizing product is registered with the U.S. ENVIRONMENTAL PROTECTION AGENCY as required by Federal law to insure that when used as directed, it has proven to be effective as claimed without causing unreasonable adverse effects to humans or the environment. This sanitizing product is authorized by USDA for use in federally inspected meat and poultry plants.

This product fulfills the sanitizing criteria of the U.S. PUBLIC HEALTH SERVICE in waters up to 500 ppm of hardness calculated as CaCO<sub>3</sub>, when tested by the AOAC Germicidal and Detergent Sanitizers - Official Method.

Steramine sanitizing solutions of 200 ppm concentration may be safely used on food processing equipment and utensils and on other food contact articles as specified under Subpart F of the Food Additives Amendment by the U.S. Food and Drug Administration.

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**WARNING** Keep out of Reach of Children. Tablet dust causes eye damage and skin irritation. Do not get dust in eyes, on skin, or on clothing. Avoid prolonged skin contact. In case of contact, flush with plenty of water. If irritation persists, get medical attention. Avoid contamination of food.

**STATEMENT OF PRACTICAL TREATMENT.** If tablets are swallowed, drink promptly large quantities of water. Avoid alcohol. Get medical attention immediately.

**NOTE TO PHYSICIAN.** Probable mucosal damage may contraindicate the use of gastric lavage.

**DISPOSAL:** Do not reuse empty container. Wrap container and put in trash.

EPA Est. 1561-VA-1

# STERAMINE 1-G TABLETS THE THIRD SINK SANITIZER

USE 1 TABLET PER GALLON OF WATER  
For the **SANITIZATION** of  
**DISHES-GLASSES-UTENSILS**

and other food contact surfaces

**KILLS HIV-1 (AIDS Virus)**

when used as directed for sanitizing

EPA REG. NO. 1561-11

**ACTIVE INGREDIENTS** 50%  
Alkyl (C14 85%, C12 3%, C16 2%) dimethyl benzyl ammonium chloride dihydrate  
**INERT INGREDIENTS** 50%

**WARNING**

**Keep Out of Reach of Children**

See Left Panel For Additional

Precautionary Statements

For Institutional and Commercial Use Only

150 TABLETS NET WEIGHT 8 OUNCES

Manufacturing Chemists

**EDWARDS-COUNCILOR CO.**

1427 BAKER ROAD, VA. BEACH, VA 23455

**DIRECTIONS**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Prepare sanitizing solutions with warm water. Allow several minutes for tablets to dissolve before using.

For sanitizing Dishes, Glasses, and Utensils in Restaurants, Taverns, etc.

1. Scrape and prewash utensils and glasses whenever possible.
2. Wash with a good detergent or compatible cleaner in first sink compartment.
3. Rinse with clean water in second sink compartment.
4. Sanitize in a solution of 1 tablet per 1 gallon of water (200 ppm) in third sink compartment. Immerse all utensils for at least one minute or for contact time specified by governing sanitary code.
5. Place sanitized utensils on a rack or drain-board to air dry.
6. A fresh sanitizing solution must be prepared at least daily or more often if the solution becomes diluted or soiled.

For Sanitizing Food Processing Equipment, Dairy Equipment, Sinks, Countertops, Refrigerated storage and Display Equipment and other hard non-porous food contact articles and surfaces.

1. Wash and Rinse all articles and surfaces thoroughly.
2. Apply a solution (Spray On or Wipe On) of 200 ppm concentration and allow surface to remain wet for at least one minute followed by adequate draining and air drying.

**KILLS HIV-1 (AIDS Virus)** when used as directed for sanitizing.

**ACCEPTED**  
**MAR 25 1994**  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 1561-11

**BEST AVAILABLE COPY**