

1561-11

3/4/2005

Page 182

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

MAR 04 2005

Mr. William Edwards
Edwards-Councilor Co., Inc.
1427 Baker Road
Airport Industrial Park
Virginia Beach, VA 23455

Subject: Steramine 1-G Tablets
EPA Registration No. 1561-11
Amendment Date: February 17, 2005
EPA Receipt Date: February 23, 2005


Dear Mr. Edwards,

The following amendment submitted in connection with registration under FIFRA, as amended, is acceptable.

A stamped copy of the accepted labeling is enclosed. Submit a copy of your final printed label before distributing or selling the product bearing the revised labeling.

If you have any questions regarding this letter, please contact Jacqueline McFarlane at (703) 308-6416.

Sincerely,



Velma Noble
Product Manager (31)
Regulatory Management Branch I
Antimicrobials Division (7510C)

Enclosure: Stamped Label

CONCURRENCES

SYMBOL							
SURNAME							
DATE							

This product fulfills the sanitizing criteria of the US PUBLIC HEALTH SERVICE in waters up to 500 ppm of hardness calculated as CaCO₃ when tested by the AOAC Germicidal and Detergent Sanitizers – Official Method.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS and DOMESTIC ANIMALS

WARNING: Tablet dust causes substantial but temporary eye injury. Do not get dust in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Tablets may be harmful if swallowed.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advise.

IF ON SKIN: Rinse immediately with water for 15-20 minutes. If irritation persists, call poison control center or doctor for treatment advise.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advise. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by poison control center doctor. Do not give anything by mouth to an unconscious person.

Have the product label or container with you when calling a poison control center or doctor, or going for treatment.

STORAGE and DISPOSAL: Store in original container in areas inaccessible to persons unfamiliar with it's use. Do not reuse empty container. Wrap container and discard in trash.



FOR SANITIZING FOOD CONTACT SURFACES

USE ONE TABLET PER ONE GALLON OF WATER

Kills HIV-1 (AIDS Virus) when used as directed for sanitizing.

EPA REG. NO. 1561-11

Active Ingredient (Quaternary).....50%
Alkyl (C14 95%, C12 3%, C16 2%) dimethyl benzyl ammonium chloride dihydrate
INERT INGREDIENTS.....50%

WARNING

KEEP OUT OF REACH OF CHILDREN

See left panel for additional precautionary statements. NET WEIGHT 8 OUNCES
For Institutional and Commercial use only.

150
Tablets

Manufacturing Chemists
EDWARDS-COUNCILOR CO. INC., 1427 Baker Road, Virginia Beach, VA 23455

DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Prepare sanitizing solutions with warm water. Allow several minutes for tablets to dissolve before using. **FOR SANITIZING DISHES, GLASSES, AND UTENSILS IN RESTAURANTS, TAVERNS, ETC.**

1. Scrape and prewash utensils and glasses whenever possible.
2. Wash with a good detergent or compatible cleaner in first sink compartment.
3. Rinse with clean water in second sink compartment.
4. Sanitize in a solution of 1 TABLET per 1 GALLON OF WATER (200 ppm) in third sink compartment. Immerse all utensils for at least one minute or for contact time specified by governing sanitary code.
5. Place sanitized utensils on a rack or drainboard to air dry.
6. A fresh sanitizing solution must be prepared at least daily or more often if the solution becomes diluted or soiled.


DIRECTIONS FOR SPRAYING

FOR SANITIZING FOOD PROCESSING EQUIPMENT, DAIRY EQUIPMENT, SINKS, COUNTERTOPS, REFRIGERATED STORAGE AND DISPLAY EQUIPMENT and other hard non-porous food contact articles and surfaces.

1. Wash and rinse all articles and surfaces thoroughly.
2. Apply a solution (SPRAY ON WITH HAND TRIGGER SPRAYER OR WIPE ON) of 200 ppm concentration and allow surface to remain wet for at least one minute followed by adequate draining and air drying.

KILLS HIV-1 (AIDS Virus) when used as directed for sanitizing.

EPA EST. 1561-VA-1




FOR CLEANING AND SANITIZING SOFT SERVE – FROZEN DESSERT MACHINES AND EQUIPMENT

CLEANING INSTRUCTIONS

1. Drain all product from machine.
2. Rinse machine thoroughly by adding 2 gallons of water and operating for 2 minutes, then drain all rinse water from machine.
3. Wash machine thoroughly by adding 2 gallons of a suitable detergent solution and operating for 5 minutes, then drain detergent solution.
4. Rinse machine again by adding 2 gallons of water and operating for 2 minutes, then drain all rinse water from machine.

SANITIZING INSTRUCTIONS

5. Dissolve 2 tablets in 2 gallons of warm water to prepare 200 ppm sanitizing solution – allow several minutes for tablets to dissolve completely before using. Add this sanitizing solution to machine and operate for 5 minutes, then drain sanitizing solution from machine. Allow 5 minutes for machine to dry.
6. Machine is now ready for refill.



2822