

Proposed addition 1 label claims are shown in typewritten, yellow highlighted text.

The containing product is registered with the U.S. ENVIRONMENTAL PROTECTION AGENCY as required by Federal law to insure that when used as directed, it has proven to be effective as claimed without causing unreasonable adverse effects to humans or the environment.

Steramine sanitizing solution of 200 ppm concentration may be safely used on food processing equipment and utensils and on other food contact surfaces as specified under Subpart F of the Food Additives Amendment by the U.S. Food and Drug Administration.

STERAMINE 60 TABLETS THE THIRD SINK SANITIZER

USE 1 TABLET PER 1½ GALLONS OF WATER
For the **SANITIZATION** of
DISHES-GLASSES-UTENSILS
and other food contact surfaces

KILLS HIV-1 (AIDS Virus)
when used as directed for sanitizing

EPA REG. NO. 1561-12
ACTIVE INGREDIENTS 50%
Allyl (C14 95%, C12 3%, C10 2%), dimethyl benzyl ammonium chloride dihydrate
INERT INGREDIENTS 50%

WARNING
Keep Out of Reach of Children
See Left Panel For Additional
Precautionary Statements

For Institutional and Commercial Use Only
100 TABLETS NET WEIGHT 6 OUNCES

Manufacturing Chemists
EDWARDS-COUNCILLOR CO.
1427 BAKER ROAD, VA. BEACH, VA 23455

This product fulfills the sanitizing criteria of the U.S. PUBLIC HEALTH SERVICE in waters up to 500 ppm of hardness calculated as CaCO₃, when tested by the AOAC Germicidal and Detergency Standards - Office Method

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING Keep out of Reach of Children
Tablet dust causes eye irritation and is a skin irritant. Do not get dust in eyes, on skin, or on clothing. Avoid prolonged skin contact. In case of contact, flush with plenty of water. If irritation persists, get medical attention. Avoid contamination of food.

STATEMENT OF PRACTICAL TREATMENT

If tablets are swallowed, drink promptly large quantities of water. Avoid alcohol. Get medical attention immediately.

NOTE TO PHYSICIAN Possible medical damage may contraindicate the use of gastric lavage.

DISPOSAL: Do not reuse empty container. Wrap container and put in trash.

S-12-1282-1

DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Prepare sanitizing solutions with warm water. Allow several minutes for tablets to dissolve before using.

For sanitizing Dishes, Glasses, and Utensils in Restaurants, Taverns, etc.

1. Spray and prewash utensils and glasses whenever possible.
2. Wash or prewash detergent or compatible cleaner in first sink compartment.
3. Rinse with clean water in second sink compartment.
4. Sanitize in a solution of 1 tablet per 1½ gallons of water (200 ppm) in third sink compartment. Immerse all utensils for at least one minute or for contact time specified by governing sanitary code.
5. Place sanitized utensils on a rack or drainboard to air dry.
6. A fresh sanitizing solution must be prepared at least daily or more often if the solution becomes diluted or soiled.

For Sanitizing Food Processing Equipment, Dairy Equipment, Sinks, Countertops, Refrigerated storage and Display Equipment and other hard non-porous food contact articles and surfaces.

1. Wash and Rinse all articles and surfaces thoroughly.
2. Apply a solution (Spray On or Wipe On) of 200 ppm concentration and allow surface to remain wet for at least one minute followed by adequate draining and air drying.

KILLS HIV-1 (AIDS Virus) when used as directed for sanitizing.

EPA Est. 1561-VA-1

ACCEPTED
JAN 31 1994
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 1561-12

BEST AVAILABLE COPY