PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS and DOMESTIC ANIMALS WARNING: Tablet dust causes substantial but temporary eye injury. Do not get dust in eyes, on skin or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Tablets may be harmful if swallowed.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advise. IF ON SKIN: Rinse immediately with water for 15-20 minutes. If irritation persists, call poison control center or doctor for treatment advise.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advise. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by poison control center doctor. Do not give anything by mouth to an unconscious person,

Have the product label or container with you when calling a poison control center or doctor, or going for treatment.

STORAGE and DISPOSAL: Store in original container in areas inaccessible to persons unfamiliar with it's use. Do not reuse empty container. Wrap container and discard in trash.



FOR SANITIZING FOOD CONTACT SURFACES

USE ONE TABLET PER 11/2 GALLONS OF WATER

Kills HIV-1 (AIDS Virus) when used as directed for sanitizing. EPA REG. NO. 1561-12

Active ingredient (Quaternary)...

WARNING



KEEP OUT OF REACH OF CHILDREN

See left panel for additional precautionary statements. 8 OUNCES

Manufacturing Chemists
EDWARDS-COUNCILOR CO. INC., 1427 Baker Road, Virginia Beach, VA 23455

DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Prepare sanitizing solutions with warm water. Allow several minutes for tablets to dissolve before using.

FOR SANITIZING DISHES, GLASSES, AND UTENSILS IN RESTAURANTS, TAVERNS, ETC.

1. Scrape and prewash utensits and glasses whenever possible. 2. Wash with a good detergent or compatible cleaner in first sink

3. Rinse with clean water in second sink compartment.

4. Sanitize in a solution of 1 TABLET per 11/2 GALLONS OF WATER (200 ppm) in third sink compartment. Immerse all utensils for at least one minute or for contact time specified by governing sanitary code.

5.Place sanitized utensils on a rack or drainboard to air dry. 6. A fresh sanitizing solution must be prepared at least daily or more often if the solution becomes diluted or soiled.

DIRECTIONS FOR SPRAYING FOR SANITIZING FOOD PROCESSING EQUIPMENT, DAIRY EQUIPMENT, SINKS, COUNTERTOPS, REFRIGERATED STORAGE AND DISPLAY EQUIPMENT and other hard non-

porous food contact articles and surfaces. 1. Wash and rinse all articles and surfaces thoroughly

2. Apply a solution (SPRAY ON WITH HAND TRIGGER SPRAYER OR WIPE ON) of 200 ppm concentration and allow surface to remain wet for at least one minute followed by adequate draining and

KILLS HIV-1 (AIDS Virus) when used as directed for sanitizing.

EPA EST. 1561-VA-1

To control ODOR CAUSING and slime FORMING BACTERIAL / ALGAE GROWTH in all manually filled EVAPORATIVE CONSOLE and PAPER FILTER TYPE HUMIDIFIERS. Not for use in Heat Vaporizing, Atomizing or Ultrasonic type Humidifiers.

Start with a clean humidifier tank and filter

Initially add I tablet per every 4 gallons of water in humidifier tank. Thereafter add 1 tablet per every 8 gallons of refill water added to tank. Tablets are effervescent and will dissolve completely.

For best results, avoid using more tablets than required. Rinse residual deposits from tank periodically. Clean your humidifier thoroughly every 4 weeks.

Do not use the product in heat vaporizing atomizing or ultrasonic humidifiers.

To control the growth of odor causing and slime forming Bacteria / Algae in waterbed mattress.

DOSAGE: To control the growth of odor causing and slime forming Bacteria / Algae in water bed mattress.

Add one (1) tablet for every 20 gallons of waterbed mattress capacity.

In a twin size waterbed mattress of 90 to 100 gallons capacity, use 5 tablets.

In a waterbed mattress of 180 to 200 gallons capacity, use 10 tablets.

Repeat this application of tablets at 4 to 6 month intervals to maintain the control of Bacterial and Slime Growth in waterbed mattresses.

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide, registered under FPA Reg. No. 156/-



