

NET CONTENTS 30 GALLONS

LIME-SULPHUR SOLUTION

GUARANTEED ANALYSIS

Active Ingredients:

Calcium Polysulphide, not less than	28%
Calcium Thiosulphate, not more than	2%
Inert Ingredients, not more than	70%
Total	100%

Baume Test at 60° F., not less than 32°.

WARNING

WARNING! Avoid contact with eyes and skin. In case of contact, wash eyes or skin with water immediately. Get medical attention for eyes.

KEEP OUT OF REACH OF CHILDREN

CAUTION

Do not use Lime Sulfur on apricots.

Do not use Oil with Lime Sulfur in summer applications.

Do not apply Oil after an application of Lime Sulfur, nor Lime Sulfur after application of Oil, until sufficient time has elapsed for safety; at least 30 days.

NOTICE

The use of this material being subject to conditions beyond their control, neither Yakima Farmers' Supply Co. nor the seller makes any representation or warranty, express or implied, with respect to results from use, whether or not used in accordance with directions. The buyer accepts and uses this material subject to these terms and shall not hold either Yakima Farmers' Supply Co. or seller liable for the results of any use of this material.

MANUFACTURED BY

YAKIMA FARMERS' SUPPLY CO.

10 East Washington Avenue

Yakima, Washington 98903

Phone GLencourt 7-5121

SUGGESTIONS FOR USE

Dormant or Delayed Dormant Sprays

On Apples and Pears for control of scale insects, blister mites, rust mites, use at rate of 11 gallons per 100 gallons water as a conventional or full gallonage spray; or at the rate of 3 gallons with 1½ gallons of superior type oil for additional control of red and brown mites.

On Stone Fruits (except Apricots) for the control of scale insects, silver mites, peach twig borer, aphids, coryneum blight, peach leaf curl — use at the rate of 11 gallons per hundred gallons of spray; or at the rate of 3 gallons with 1½ gallons of superior type oil for additional control of brown and red mites.

Pre - Pink, Pink Sprays

On all sulfur tolerant* varieties of apples and pears for control of scab and powdery mildew, use at the rate of 2 to 3 gallons per 100 gallons of spray.

Post - Harvest Spray

On pears for control of pear leaf blister mites, use at the rate of 4 gallons per 100 gallons of spray.

*Consult your fieldman or Extension Service for specific information.

