

[1588-13]

Extermo Vaporant Insecticide is specifically prepared for use around food. Extermo Vaporant is especially designed for use in cafes, restaurants, food storage warehouses, flour mills, meat packing plants with electric mist sprayer equipment.

Use of the product in food processing plants should be limited to those times when the plant is not in operation.

Food and foodstuffs must be covered as well as food handling equipment. In meat packing plants washing down of walls, meat handling equipment and surfaces with which the meat will come in contact is necessary prior to reusing the treated room.

DO NOT REUSE EMPTY DRUM. RETURN TO DRUM RECONDITIONER OR DESTROY BY PERFORATING OR CRUSHING AND BURYING IN A SAFE PLACE AWAY FROM WATER SUPPLIES.

CAUTION


- Do not spray near fire or open flame.
- Do not spill, pour, or store near fire or flame.
- Harmful if swallowed.
- Store in a cool place and keep container tightly closed.
- Keep out of reach of children.
- Remove pets, birds and cover aquariums before spraying.

EPA Est. 1588-IA-1

EPA Est. 1588-NY-1



**VAPORANT
CONCENTRATED**



EPA Reg. No. 1588-13

PERMETHRIN BASED INSECTICIDE SPECIFIC, LLY
COMPOUNDED FOR USE IN ELECTRIC
MIST SPRAYERS

- ACTIVE INGREDIENTS 100% CONSISTING OF:
- PETROLEUM DISTILLATES - 98.600%
 - N-OCTYL BICYCLOHEPTENE DICARBOXIMIDE - 0.625%
 - PYRETHRINS - 0.400%
 - PIPERONYL BUTOXIDE, TECHNICAL* - 0.375%

*Equivalent to 0.300% (butylcarbityl) (6-propylpiperonyl) ether and 0.075% related compounds

CAUTION: Keep Out Of Reach Of Children.

Harmful If Swallowed.

See left panel for additional cautions



ROCHESTER GERMICIDE CO.
ROCHESTER, NEW YORK 14603

DIRECTIONS

For Control of Flies, Mosquitoes, Gnats, Wasps, and Adult Moths:

Close all windows and stop all ventilation. Set machine to deliver mist in as fine a particle size as possible. Use not less than 5 oz. per 10,000 cu. ft. Keep area closed for a minimum of 20 minutes after mist has been applied. Sweep up and destroy fallen insects. Repeat treatment as necessary.

DO NOT REMAIN IN TREATED AREAS. VENTILATE THE AREA AFTER TREATMENT IS COMPLETED.

For Control of Crawling Insects such as Roaches, Silverfish, Ants, Spiders:

Set equipment to deliver a coarse spray. Direct spray at baseboards and into all cracks and crevices, behind switch boxes, around motor mounts, under skids, behind sinks, at door and window casings, in double wall partitions, around scales and conduits and behind conduit terminal boxes. Spray should be applied to penetrate into these areas. Spray should be applied to point of run off for control of crawling insects. If ant control is also desired, thoroughly spray door and window casings and all trail ways and nests near the building being treated. Check the building's foundations for cracks, spraying these cracks and patching them if possible.

If aerosol space treatment for exposed crawling insects is desired, set machine for fine mist, using 6-13 oz. per 10,000 cu. ft. It is generally preferable, however, to use a coarse spray as described above. Repeat treatment for crawling insects as necessary.

DO NOT REMAIN IN TREATED AREAS. VENTILATE THE AREA AFTER TREATMENT COMPLETED.

For Control of Bedbugs:

Spray mattress, bedstead, corners and all cracks and crevices in the room. Use coarse spray. Repeat as needed. Allow treated