

UNDER THE FEDERAL INSECTICIDE ACT
AND THE FEDERAL RESIDUE ACT
THIS PRODUCT IS SUBJECT
TO ATTACHED COMMENTS.



FCX SEVIN 10 "D"



An insecticidal dust containing 10 % Sevin* Carbaryl insecticide

ACTIVE INGREDIENTS:	
Carbaryl (1-naphthyl N-methylcarbamate)	10.0 %
INERT INGREDIENTS:	
	90.0 %
TOTAL	100.0 %

* Sevin is Union Carbide Corporation's trademark for Carbaryl, U.S. Pat. Nos. 2,903,478 and 3,009,855.

USDA Registration Number 1598-82

CAUTION KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or swallowed. Avoid breathing of dust. Avoid contact with skin. Wear clean long-sleeved clothing and change daily. Wash hands and face thoroughly after use and before eating, drinking, or smoking.

Note to Physician: Carbaryl is a moderate reversible, cholinesterase inhibitor. Atropine is antidotal.

DIRECTIONS FOR USE

FCX Sevin 10 "D" is a ready to use product and should not be mixed with water. Read entire label and follow directions and precautions.

Tobacco: To control Budworm, use a rotary hand duster to apply insecticide to each bud, applying 4 pounds per acre. Begin applications when budworms first appear and repeat as necessary when worms are less than 1/2 inch long.

To control Hornworm, Flea Beetle, Japanese Beetle, and Grasshoppers, use 10 to 15 pounds per acre applied with power driven ground equipment or aircraft. Thorough coverage is essential for effective control. Usually, use lesser amounts on smaller plants and slight infestations and greater amounts on larger plants and heavy infestations.

To avoid injury to **tender foliage**, do not apply when foliage is wet or when rain or excessive humidity is expected for the next two days.

Soybeans: To control Corn Earworm, Mexican Bean Beetle, Bean Leaf Beetle, Japanese Beetle, Armyworm, Green Cloverworm, Blister Beetle, Thrip, Cucumber Beetle, and Velvetbean Caterpillar, use 10 to 15 pounds per acre applied with power driven ground equipment or aircraft. Usually, use

lesser amounts on smaller plants and slight infestations and greater amounts on larger plants and heavy infestations.

To avoid injury to **tender foliage**, do not apply when foliage is wet or when rain or excessive humidity is expected for the next two days.

Cotton: For early season Thrip, Flea Beetle, and Cotton Fleahopper control, use 7 to 10 pounds per acre applied with power driven ground equipment or aircraft.

For mid and late season control of Boll Weevil, Bollworm, Fall Armyworm, Cotton Leaf Perforator, Leaf Roller, Leaf hopper, and Tarnish Plant Bug, use 10 to 20 pounds applied with power driven ground equipment or aircraft. Usually, use lesser amounts on smaller plants and slight infestations and greater amounts on larger plants and heavy infestations.

To avoid injury to **tender foliage**, do not use when foliage is wet or when rain or excessive humidity is expected in the next two days.

CAUTION

May kill honeybees in substantial numbers. This product is highly toxic to bees exposed to direct treatment or residues on crops. Do not use when the value of bees as pollinators is more important than insect control. Before applying, warn beekeepers to locate hives beyond bee flight range until one week after application or to take other effective precautions. Additional protective information may be obtained from the Cooperative Agriculture Extension Service.

Do not allow to contaminate feed and foodstuffs.

Do not allow to contaminate any body of water.

Do not use in any manner not specified on this label.

Clean equipment and dispose of waste in a safe place away from water supplies. Dispose of empty bag in same manner or by burning. If burned, keep out of smoke.

NOTICE

Follow directions carefully. Timing and method of application, weather and crop conditions, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given on this label.

Net Contents

MANUFACTURED BY

U.S. Pounds



1598-82
ACCEPTED
 Apr 22 1971
 UNDER THE FEDERAL INSECTICIDE ACT
 THE UNITED STATES DEPARTMENT OF AGRICULTURE
 HAS REVIEWED THIS PRODUCT SUBJECT
 TO ATTACHED COMMENTS



FCX SEVIN 10 "D"



An insecticidal dust containing 10 % Sevin[®] Carbaryl insecticide

ACTIVE INGREDIENTS:
 Carbaryl (1-naphthyl N-methylcarbamate) 10.0 %

INERT INGREDIENTS: 90.0 %

TOTAL 100.0 %

[®] Sevin is Union Carbide Corporation's trademark for Carbaryl, U.S. Pat. Nos. 2,903,478 and 3,009,855.

USDA Registration Number 1598-82

CAUTION
KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or swallowed. Avoid breathing of dust. Avoid contact with skin. Wear clean long-sleeved clothing and change daily. Wash hands and face thoroughly after use and before eating, drinking, or smoking.

Note to Physician: Carbaryl is a moderate reversible, cholinesterase inhibitor. Atropine is antidotal.

DIRECTIONS FOR USE

FCX Sevin 10 "D" is a ready to use product and should not be mixed with water. Read entire label and follow directions and precautions.

Tobacco: To control Budworm, use a rotary hand duster to apply insecticide to each bud, applying 4 pounds per acre. Begin applications when budworms first appear and repeat as necessary when worms are less than 1/2 inch long.

To control Hornworm, Flea Beetle, Japanese Beetle, and Grasshoppers, use 10 to 15 pounds per acre applied with power driven ground equipment or aircraft. Thorough coverage is essential for effective control. Usually, use lesser amounts on smaller plants and slight infestations and greater amounts on larger plants and heavy infestations.

To avoid injury to tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected for the next two days.

Soybeans: To control Corn Earworm, Mexican Bean Beetle, Bean Leaf Beetle, Japanese Beetle, Armyworm, Green Cloverworm, Blister Beetle, Thrip, Cucumber Beetle, and Velvetbean Caterpillar, use 10 to 15 pounds per acre applied with power driven ground equipment or aircraft. Usually, use

lesser amounts on smaller plants and slight infestations and greater amounts on larger plants and heavy infestations.

To avoid injury to tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected for the next two days.

Cotton: For early season Thrip, Flea Beetle, and Cotton Fleahopper control, use 7 to 10 pounds per acre applied with power driven ground equipment or aircraft.

For mid and late season control of Boll Weevil, Bollworm, Fall Armyworm, Cotton Leaf Perforator, Leaf Roller, Leaf hopper, and Tarnish Plant Bug, use 10 to 20 pounds applied with power driven ground equipment or aircraft. Usually, use lesser amounts on smaller plants and slight infestations and greater amounts on larger plants and heavy infestations.

To avoid injury to tender foliage, do not use when foliage is wet or when rain or excessive humidity is expected in the next two days.

CAUTION

May kill honeybees in substantial numbers. This product is highly toxic to bees exposed to direct treatment or residues on crops. Do not use when the value of bees as pollinators is more important than insect control. Before applying, warn beekeepers to locate hives beyond bee flight range until one week after application or to take other effective precautions. Additional protective information may be obtained from the Cooperative Agriculture Extension Service.

Do not allow to contaminate feed and foodstuffs.

Do not allow to contaminate any body of water.

Do not use in any manner not specified on this label.

Clean equipment and dispose of waste in a safe place away from water supplies. Dispose of empty bag in same manner or by burning. If burned, keep out of smoke.

NOTICE

Follow directions carefully. Timing and method of application, weather and crop conditions, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given on this label.

Net Contents MANUFACTURED BY U.S. Pounds

BEST DOCUMENT AVAILABLE **FCX, Inc. RALEIGH, N.C.**