

FCX FERBAM NO. 10 DUST
FUNGICIDE
For Control of Blue Mold and Anthracnose on Tobacco Plant Beds



INGREDIENTS

ACTIVE INGREDIENT:	
Ferbam (ferric dimethyl dithiocarbamate)	11.4 %
INERT INGREDIENTS:	88.6 %
Total	100.0 %

EPA Reg. No. 1598-90

CAUTION
KEEP OUT OF REACH OF CHILDREN

May cause irritation of nose, throat, eyes, and skin. Harmful if swallowed. Do not breathe dust. Wear a respirator. Avoid contact with eyes, skin, or clothing. In case of contact with skin, wash with soap and water; for eyes, flush with plenty of water and get medical attention. Avoid contamination of feed and foodstuffs. Store in a cool dry place. Keep container closed when not in use.

This product is toxic to fish. Keep out of any body of water. Apply only in the manner and for the purpose stated in this label. Do not apply when weather conditions favor drift of dust from area treated. Clean application equipment and dispose of wastes and unused chemical by burial in a pit on non-crop land well away from water supplies.

DESTROY EMPTY CONTAINERS. Do not re-use this container for any purpose. Bury empty containers (paper bags) deeply in a pit in an isolated area on non-crop land well away from water supplies.

DIRECTIONS FOR USE

To control blue mold and anthracnose in tobacco plant beds, begin applications when tobacco plants are about the size of a dime, or when blue mold is first reported in your area, and repeat at intervals of every 4 to 5 days, depending upon the severity of the diseases and upon weather conditions which favor development of the diseases, until the tobacco plants are transplanted into the field. Under warm humid conditions, frequent rainfall, and heavy dews, certain applications at intervals of 2 to 3 days may be required, particularly if blue mold is epidemic in your area, until the disease is controlled.

The dust should be applied when the air is calm -- in the early morning or in the evening -- preferably when the tobacco plants are moist with dew. Use a good rotary-type hand duster for large beds; a plunger-type duster may be used on small beds. Do not over-fill the duster at any time beyond its capacity for uniformly thorough application. Make certain that all leaves on the tobacco plants are uniformly covered with a light film of dust. An application that is washed off the leaves by rainfall should be repeated as soon as possible. Usually, 8 to 12 applications are required per season, and 35 pounds of dust should be sufficient to treat 100 square yards of plant bed for the season.

The quantities to apply per 100 square yards of plant bed are:

1st to 4th application	1 1/2 pounds
5th and 6th application	2 1/2 to 3 pounds
7th and later, if required	3 1/2 pounds

NOTICE

Follow directions carefully. Timing and method of application, weather and crop conditions and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given on this label.

1598-90
ACCEPTED
Jul 3 1972
UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR FOOD AND DRUGS REGISTERED
 UNDER NO. 1598-90
 NO ADVERSE COMMENTS

Net Weight

Manufactured By

25 Pounds

FCX, Inc. RALEIGH, N.C.