




FCX

SEVIN 5 "D"



An Insecticidal Dust Containing 5% Sevin* Carbaryl Insecticide.

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) 5.0%

INERT INGREDIENTS: 95.0%

TOTAL 100.0%

*Sevin is Union Carbide Corporation's trademark for Carbaryl, U.S. Pat. Nos. 2,903,478 and 3,009,855.

EPA Registration No. 1598-92

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or swallowed. Avoid breathing of dust. Avoid contact with skin. Wear clean long-sleeved clothing and change daily. Wash hands and face thoroughly after use and before eating, drinking or smoking.

Note to physician: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal.

FCX Sevin 5 "D" is a ready to use dust and should not be mixed with water. Read entire label and follow directions and precautions carefully.

SEE BACK PANEL FOR ADDITIONAL DIRECTIONS

Net Contents 5 US Pounds

MANUFACTURED BY



Home Gardens

To control the insects on the crops listed below, apply the dust uniformly and thoroughly to upper and lower surfaces of the foliage, but do not apply excessive amounts. Begin applications when insects or their damage first appears. Repeat as necessary, generally at weekly intervals. May be applied on day of harvest unless otherwise stated.

Beans: Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetle, Japanese Beetle Adults, Leafhoppers, Flea Beetles, and Corn Earworm.

Corn (Sweet): Corn Earworms, Fall Armyworms, Flea Beetles, Japanese Beetle Adults, Sap Beetles, and Leafhoppers. For larvae in the whorls (Fall Armyworms) and foliage feeders, apply to entire plant before tasseling. For insects attacking silks and ears apply when silks first appear and repeat at 2 to 3 day intervals until silks become dry. Three or more applications may be required. Proper timing and uniformly thorough coverage are essential. Application of this product to the tassel region of corn during the pollen-shed period will seriously reduce bee populations.

Cucumbers, Melons, and Squash: Pickle Worm, Melonworm, Cucumber Beetles, Squash Bug, and Leafhopper. Avoid excessive applications. Do not use on watermelons.

Potatoes, Tomatoes, Eggplant, and Peppers: Colorado Potato Beetle, Leafhoppers, Flea Beetle, Fall Armyworm, Tomato Fruit Worm, Tomato Hornworm, Blister Beetle, and Tarnished Plant Bug.

Cabbage, Broccoli, and Cauliflower: Harlequin Bug, Leafhoppers, Imported Cabbage Worm, and Corn Earworm. Do not apply within 3 days of harvest.

Garden Beets, Lettuce, Radishes, Spinach, Turnips, and Carrots: Flea Beetles, Leafhoppers, Harlequin Bug, Fall Armyworm, Corn Earworm, Spittlebugs, and Tarnished Plant Bug.

Do not apply to garden beets, head lettuce, radishes, and turnips within 3 days of harvest. Do not apply to leaf lettuce and spinach or to garden beets and turnips, if tops are to be used for food or feed, within 14 days of harvest.

Black-eyed Peas: Cowpeas Curculio and Lygus Bug.

Okra: Japanese Beetle Adults and Blister Beetles.

Strawberries: Meadow Spittlebug, Strawberry Leaf Roller, and Strawberry Weevil. Do not apply within 1 day of harvest.

Some injury to the crops named above may occur if treatments are made when foliage is wet or during periods of high humidity for two or three days following an application.

CAUTION

May kill honeybees in substantial numbers. This product is highly toxic to bees exposed to direct treatment or residues on crops. Do not use when the value of bees as pollinators is more important than insect control. Before applying warn beekeepers to locate hives beyond bee flight range until one week after application or to take other effective precautions. Additional protective information may be obtained from the Cooperative Agriculture Extension Service.

Do not allow to contaminate feed and foodstuffs.

Do not allow to contaminate any body of water.

Do not use in any manner not specified on this label.

Clean equipment and dispose of waste in a safe place away from water supplies. Dispose of empty bag in same manner or by burning. If burned, keep out of smoke.

NOTICE

Follow directions carefully. Timing and method of application, weather and crop conditions and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given on this label.

ODF 3-71

FCX SEVIN 5 "D"