



FCX 1% LINDANE DUST



ACTIVE INGREDIENTS

Lindane (gamma isomer of benzene
hexachloride) 1%
INERT INGREDIENTS 99%
TOTAL 100%

EPA Reg. No. 1598-115

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing dust. Avoid contact with skin and eyes. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs.

This product is poisonous to fish and wildlife. Birds feeding in treated areas may be killed. Keep out of any body of water. Apply only for the purposes and in the manner directed in this label. Do not apply when weather conditions favor drift of dust from treated. Do not use where runoff may occur as a result of heavy rain. Clean application equipment and dispose of wastes and unused chemical in a pit on non-crop land in an isolated area well away from water supplies.

This product is toxic to bees exposed to direct treatment or to residues, on treated crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. Bury empty containers in a pit on non-crop land in an isolated area well away from water supplies.

DIRECTIONS FOR USE

CUCUMBERS AND MELONS--To control cucumber beetles and pickleworm, apply the dust uniformly to upper and lower surfaces of foliage and into blossoms (30 to 40 pounds of dust per acre.) Begin applications when insects first appear, which is usually when the plants are in the 4 to 6 leaf stage, and repeat if necessary at 5 to 10 day intervals, depending upon severity of insect infestation and weather conditions favorable to insect development. **DO NOT APPLY TO CUCUMBERS WITHIN 1 DAY OF HARVEST.**

SQUASH AND PUMPKIN--To control aphids on foliage, pickleworm, and squash vine borer, apply the dust uniformly to upper and lower surfaces of foliage, to stems and vines at base of plants, and into blossoms (30 to 40 pounds of dust per acre). Begin applications when the plants are small (4 to 6 leaf stage) and repeat if necessary at 7 day intervals depending upon insect infestation and weather conditions favorable to insect development.

NOTICE

Follow directions carefully. Timing and method of application, weather and crop conditions, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given on this label.

NET WEIGHT

MANUFACTURED BY

25 POUNDS

FCX, Inc. RALEIGH, N.C. 27602

ACCEPTED WITH COMMENTS

