



"73" POTATO AND TOMATO DUST



Combination Insecticide and Fungicide for Potatoes, Tomatoes, and Certain Other Vegetable Crops.

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate)*	3.00 %
Cuprous oxide	7.10 %
Cupric oxide	.88 %
INERT INGREDIENTS	89.02 %
Total	100.00 %

CONTAINS 7% COPPER AS ACTIVE

EPA REG. NO. 7-136
EPA EST 1598-G-1

*SEVIN

Trademark of Union Carbide Corp. for 1-naphthyl N-methylcarbamate

CAUTION

KEEP OUT OF REACH OF CHILDREN

(See remainder of CAUTION Statement on side panel of this label.)

Harmful if inhaled or swallowed. Do not breathe dust. Do not take internally. Skin contact may be harmful. Avoid contact with skin. Wash hands before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing daily. Do not contaminate feed and foodstuffs.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is ant dotal.

This product is toxic to fish. Keep out of streams, lakes or ponds; do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or to residues on crops or foliage. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DISPOSAL OF EMPTY CONTAINERS: Do not reuse empty container. Destroy it by perforating or tearing. Bury or discard in a safe place away from water supplies.

DIRECTIONS FOR USE

To control the insects and the plant diseases on the field-grown crops listed below, apply uniformly and thoroughly to upper and lower surfaces of the foliage, but do not apply excessive amounts. Begin applications when the insects or their damage first appear and before diseases appear (when disease is first reported in the area). Repeat as necessary, generally at 7-day intervals. Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration.

TOMATOES: Tomato fruitworm, tomato hornworm, leafhoppers, European corn borer, fall armyworm, tarnished plant bug, Colorado potato beetle, early blight, late blight, septoria leaf spot, and anthracnose on field grown tomatoes. Apply 3 to 3½ lb. of dust per 1000 linear feet of row.

POTATOES: Colorado potato beetle, leafhoppers, flea beetles, fall armyworm, European corn borer, early blight, and late blight. Apply 3 pounds of dust per 1000 linear feet of row to foliage only.

GARDEN BEETS: Flea beetles, tarnished plant bug, stink bug, and cercospora leaf spot. Apply 2½ to 3 lbs. of dust per 1000 linear feet of row. **DO NOT APPLY TO BEETS WITHIN 14 DAYS OF HARVEST IF TOPS ARE TO BE USED AS FOOD OR FEED; IF TOPS WILL NOT BE USED AS FOOD OR FEED, DO NOT APPLY WITHIN 3 DAYS OF HARVEST.**

EGGPLANT AND PEPPERS: Leafhoppers, fall armyworm, European corn borer, tarnished plant bug, alternaria and phomopsis leaf spots. Apply 2 to 2½ lbs. of dust per 1000 linear feet of row.

CUCUMBERS, MELONS, AND SQUASH: Cucumber beetles, leafhoppers, squash bug, pickleworm, melonworm, downy mildew, and anthracnose. Apply 2½ lbs. of dust per 1000 linear feet of row. Some leaf injury may occur if applications are made when foliage is wet or during periods of prolonged high humidity. Do not use on watermelons.

NOTICE

Follow directions carefully. Timing and method of application, weather and crop conditions and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given on this label.

ODF 1171

NET WEIGHT

MANUFACTURED BY



INCORPORATED

25 POUNDS

RALEIGH, N.C. 27602