



# FCX 20-0 TOXAPHENE DUST



### INGREDIENTS

#### ACTIVE INGREDIENT:

Toxaphene (Technical chlorinated camphene containing 67-69 per cent chlorine) 20 %

#### INERT INGREDIENTS

Total 80 %  
100 %

### INSECTICIDE

FOR CONTROL OF FALL ARMYWORM, THRIPS, COTTON FLEAHOPPER, AND LYGUS (STINK) BUGS ON GROWING COTTON

## CAUTION

**KEEP OUT OF REACH OF CHILDREN**

See Remainder of CAUTION Statement on Side Panel

EPA Reg. No. 1598-154

### DIRECTIONS FOR USE

This product is a ready-to-use insecticide dust to be applied without mixing or diluting with any substance, intended for use on growing cotton plants to control the insects listed previously in this label. Apply with rotary hand dusting equipment, power-driven ground dusting machines, or by aircraft. Uniformly thorough dust coverage of the foliage of the cotton plants is essential.

During the early season (seedling to 8-leaf stage), apply at the rate of 15 to 20 pounds of dust per acre to control thrips, Lygus bugs, and cotton fleahopper. Begin applications when one or more of the insects is determined to be present in sufficient numbers to cause substantial damage. In some instances the early-season applications have been omitted without extensive damage from the insects, but this is dependent upon seasonal conditions which cannot be predicted in advance. If early-season treatment is required, apply only when the insects are determined to be threatening in each particular field. Repeated applications at this stage may or may not be necessary, depending upon the occurrence of the insects and upon seasonal conditions favorable to their development. Continued occurrence and development of the insects will make two or more repeated applications necessary at subsequent 7 to 10-day intervals.

During the mid-season (squaring to full bloom), apply at the rate of 15 to 20 pounds of dust per acre to control cotton fleahopper and Lygus bugs. Apply when the insects are first determined to be present in sufficient numbers to cause substantial damage in each particular field. One or more repeated applications, based on determinations that the insects are actually threatening in each particular field, made at successive intervals of 7 to 10 days, are usually necessary.

During the late season (bolls set), apply at the rate of 15 to 20 pounds of dust per acre to control fall armyworm. Begin applications when the insects are determined to be present in sufficient numbers to cause substantial damage in each particular field. Two or more repeated applications, based on determinations that the insects are actually threatening, made at successive intervals of 7 to 10 days, are usually necessary.

Apply the dust only when the air is calm. Speed of air movement must not exceed 5 miles per hour. If higher air speed occurs, suspend dusting until air movement becomes calm again.

**DO NOT GRAZE LIVESTOCK IN COTTON FIELDS TREATED WITH THIS PRODUCT. DO NOT FEED GIN TRASH OR WASTE FROM COTTON TREATED WITH THIS PRODUCT TO LIVESTOCK.**

Do not apply the dust on or near areas where dairy animals or animals being finished for slaughter will graze. Do not apply the dust on or near any body of water.

**AIRCRAFT APPLICATIONS**--This product may be applied only by operators, applicators, and-or pilots licensed by competent regulatory authority which has inspected and certified the aircraft and dusting equipment, and which as a prerequisite for obtaining and maintaining the license requires compliance with its established regulations for application of insecticide dusts to cotton fields.

## CAUTION

Harmful if swallowed. Avoid inhalation of dust. Avoid contact with skin, eyes, and clothing. In case of skin contact, wash thoroughly with soap and water; if in eyes, flush with plenty of water and get medical attention. Wash thoroughly after using and before eating or smoking. Wash contaminated clothing with soap and water before reuse. Do not contaminate feed and food-stuffs.

This product is poisonous to fish, bird, and other wildlife. Keep out of any body of water. Apply only for the purposes and in the manner directed in this label. Do not apply when weather conditions favor drift of dust from area treated. Do not use where runoff may occur as a result of heavy rain. Clean application equipment and dispose of waste and unused chemical in a pit on non-crop land well away from water supplies.

This product is toxic to bees exposed to direct treatment and to residue, on treated crop. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**DESTROY EMPTY CONTAINERS.** Do not reuse this container for any purpose. Bury empty containers in a pit on non-crop land in an isolated area well away from water supplies.

## NOTICE

Follow directions carefully. Timing and method of application, weather and crop condition and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage, or handling of this material in strict accordance with directions given on this label.

ACCEPTED  
JAN 29 1973  
FUNGICIDE AND PESTICIDE DIV.  
U.S. DEPARTMENT OF AGRICULTURE  
ED UNDER NO. 100-100000-1  
TO ATTACHED COMMENTS.

Net Weight

Manufactured By

Pounds