FCX POLYRAM 3.5 "D"

ACTIVE INGREDIENTS:

A mixture of 5.2 parts by weight (83.9%) of ammoniates of [ethylenebis (dithiocarbamate)] zinc with 1 part by weight (16.1%) ethylenebis [dithiocarbamic acid] bi- molecular and tri-molecular cyclic anhydrosulfides	
and disulfides	3.5 %
TOTAL	

USDA Registration No. 1598-160

CAUTION

KEEP OUT OF REACH OF CHILDREN

May be harmful if swallowed or inhaled. May cause irritation of eyes, nose, throat, and skin. In case of contact with skin, wash with soap and water; eyes, flush with clear water and get medical attention. Wash thoroughly after use and before eating, drinking, or smoking.

ACCEPTED 3-10-71 1598-160 Manufactured by Net Weight 25 Pounds FCX, MC. RALEIGH, N.C.

Follow directions carefully. Timing and method of application, weather and crop conditions, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given on this label.



DIRECTIONS FOR USE

FCX Polyram 3.5"D" is a ready to use fungicidal dust to be used as a preventive in the control of certain plant diseases on those crops as listed below. Use equipment, such as a rotary or power duster, that will give thorough, complete coverage of plants.

Tobacco Plant Beds: To control Blue Mold, use two (2) to four (4) pounds per 100 square yards of plant bed. Make first application when plants are about the size of a dime. Repeat applications twice weekly until plants are transplanted.

Do not use this product in any manner not specified on this label.

Do not re-use empty container. Clean equipment and dispose of waste by burying in non-crop land away from water supplies. Dispose of empty container in same manner or by burning. If burned, keep out of smoke.

NOTICE

