

## FCX POLYRAM' 5 DUST W/BORAX



(FUNGICIDE-NUTRITIONAL)

## **ACTIVE INGREDIENT:**

A mixture of 5.2 parts by weight (83.9%) of ammoniates of [ethylenebis (dithiocarbamato)] zinc with 1 part by weight (16.1%) ethylenebis (dithiocarbamic acid) bimolecular and trimolecular cylic anhydrosulfides and disulfides\*..... 5.00 %

INERT INGREDIENTS: \*\* 95.00 % TOTAL ..... 100.00 %

\*Zinc expressed as metallic 0.85 \*\*Boron expressed as metallic

USDA Registration No. 1598—173

## CAUTION

KEEP OUT OF REACH OF CHILDREN

May be harmful if swallowed or inhaled. Do not breathe dust. May cause irritation of eyes, nose, throat, and skin. In case of contact, flush with plenty of water; for eyes get medical attention.

Polyram is a registered Trademark of Badische Anilin & Soda Fabrik, A.G.

ACCEPTED

73 STENE CTIONS FOR USE

PEANUTS — For the control of Cercospora Leaf Spot and as an aid in correcting boron dificiency, apply 20 pounds of dust per acre by power driven ground applicator or aircraft. Apply when disease spots first appear on old leaves at the base of the plants. Make only one application per season. Apply only when foliage is dry. If applied when foliage is wet, severe burning may occur.

Do not feed treated forage to dairy animals or animals being finished for slaughter.

CAUTION: Use of this material in a manner or at a time other than in accordance with directions may produce plant injury, excessive residues or other undesirable results.

Dispose of waste by burying in a pit in non-crop lands located away from water supplies. Dispose of empty bags in the same manner or by burning. If burned, keep out of smoke.

## NOTICE

Follow directions carefully. Timing and method of application, weather and cropconditions and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given on this label.

Net Weight 50 Pounds

Manufactured By

