

FCX COPPER - SULPHUR - SEVIN* DUST

A PEANUT INSECTICIDE - FUNGICIDE COMBINATION

ACTIVE INGREDIENTS:

Copper, Expressed as metallic (From cuprous and cupric oxides)	4%
Sulphur	80%
Carbaryl (1-naphthyl N-methylcarbamate)*	5%

INERT INGREDIENTS

	11%
TOTAL	100.0%

*Sevin is the Registered Trade Mark of Union Carbide Corporation for Carbaryl (1-naphthyl N-methylcarbamate)
USDA Registration Number 1598-1-6

HARMFUL if swallowed, inhaled or absorbed thru skin. Do not breathe dust. May cause irritation of eyes, nose, throat or skin. In case of contact flush with plenty of water; for eyes, get medical attention. Wash thoroughly with soap and water after handling and before smoking, eating or drinking. Wash all contaminated clothing with soap and water before re-using.

ANTIDOTE: Carbaryl is a moderate, reversable, cholinesterase inhibitor. Atropine is antidotal. (See additional precautionary statements following Directions For Use)

PEANUTS: FCX Copper-Sulphur-Sevin* Dust will effectively control these listed disease and insects on peanuts when used as directed: Cercospora leafspot; Spidermites; Corn earworm; Fall armyworm; Southern armyworm; Potato Leafhopper; Green cloverworm; Velvetbean caterpillar and Thrips.

MANUFACTURED BY

Net Contents

Raleigh, N. C.

50 US Pounds

ACCEPTED

Apply 20 to 25 pounds per acre with ground or aircraft equipment beginning about the last week in June or first week in July. To control Cercospora leafspot begin applications before disease appears. Continue making applications at 10 to 14 day intervals as needed. Usually about 5 applications will be sufficient. If material should be washed off plants within a few hours after application, repeat application as soon as weather conditions permit.

Dust respirators and protective clothing should be used during frequent and prolonged exposure.

This material is toxic to honey bees. When it is necessary to use during periods of bee activity on the crop, warn beekeepers in advance to remove hives to a safe distance from area to be treated until one week after application.

Avoid contamination of feed and foodstuffs.

Do not apply when weather conditions favor drift from treated areas to unprotected areas.

Do not allow to contaminate any body of water.

Do not use in any manner not specified on this label.

Dispose of waste by burying in non-crop land away from water supply. Destroy container in same manner or by burning. If container is burned,

Read and follow directions carefully. Timing and method of application, weather and crop conditions and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given on this label.