

ACTIVE INGREDIENTS:

Copper, Expressed as Metallic _____ 4.00% (From Cuprous & Cupric Oxides) Sulphur _____ 70.00% Boron _____ 3.34% INERT ING 22.66%

> 100.00% TOTAL

1598-188

CHILDREN

& absorbed through skin. irritation of eyes, nose, at with skin, wash thoroughin case of contact with eyes, __u get medical attention. Wash and water after handling and before or drinking. Wash contaminated clothing smok... with soap and water before re-using.

(See additional precautionary statements following Directions for Use.)

DIRECTIONS FOR USE

PEANUTS: FCX Copper-Sulphur Peanut Dust with Boron will effectively control Cerecospora leafspot while adding plant nutrition. Apply 15 pounds per acre with ground or aircraft equipment. Check with the local Agriculture Extension Service as to best date for application. This should replace one treatment of the regular disease control program.

Do not apply over 15 pounds our acre or make more than one application per season. The recommended rate wil give 1/2 pound of Boron per acre per season. More than this could damage the crop. Do not use this product on any crop other than peanuts.

CAUTION

Dust respirators and protective clothing should be used during frequent and prolonged exposure.

Do not contaminate feed and foodstuffs.

Do not apply when weather conditions favor drift from treated rreas to unprotected areas.

Do not allow to contaminate any body of water.

Do not use in any manner not specified on this label.

Dispose of waste by burying in non-crop land away from water supplies. Destroy empty bag in same manner or by burning. If container is burned, keep out of smoke.

NOTICE

Follow directions carefully. Timing and method of application, weather and crop conditions and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given on this label.

50 U.S. Pounds

