

DIRECTIONS FOR USE

HOME GARDENS

Mix FCX Cythion 5 at the rate of 1 1/4 teaspoons per gallon of water. Agitation is desired. Make thorough and complete coverage of the entire plant to control the pests on the specified crops listed below.

VEGETABLES

APHIDS: (PLANT LICE): Beans, Beets, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Eggplant, Lettuce, Melons, Mustard, Okra, Peas (cowpeas and garden), Peppers, Rutabaga, Squash, Spinach, Tomato, Turnips.

CUCUMBER BEETLE: Beans, Cucumbers, Melons.

DIAMOND BACK MOTH LARVAE AND IMPORTED CABBAGE WORM: Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Collards, Lettuce, Mustard, Rutabaga, Turnips.

JAPANESE BEETLES: Beans.

LEAFHOPPERS: Beans, Lettuce, Melons, Okra.

MEXICAN BEAN BEETLE: Beans.

TWO-SPOTTED SPIDER MITES: Cucumber, Eggplant, Tomatoes.

DO NOT USE within 1 day of harvest of Beans, Cucumbers, Melons, Squash, or Tomatoes. DO NOT USE within 3 days of harvest of Broccoli, Eggplant, Peas, Peppers, Rutabagas, or Turnips (roots only; wait 10 days for greens).

DO NOT USE within 14 days of harvest of Leaf Lettuce. DO NOT USE on Head Lettuce within 7 days of harvest.

DO NOT USE on Okra after the pods form.

If forage is to be fed to livestock, do not apply to peas within 7 days of harvest.

DO NOT USE on Cucumbers when foliage is wet.

BERRIES

APHIDS (Plant Lice): Blackberry, Boysenberry, Dewberry, Loganberry, Strawberry.

CHERRY FRUIT WORM: Blueberry.

JAPANESE BEETLE: Blackberry, Blueberry, Boysenberry, Dewberry, Loganberry.

LEAFHOPPERS: Blackberry, Boysenberry, Dewberry, Loganberry, Strawberry.

ALSO ON STRAWBERRIES for Lygus Bug, Spider Mites, Spittlebug, Strawberry Leaf Roller, Thrips and White Flies.

DO NOT USE on any of the above named berry crops within 3 days of harvest.

ORNAMENTAL PLANTS

FCX Cythion 5 applied in a thorough, complete coverage at these listed rates will give

Accepted
Container
Sizes
PT
3
FCX



FCX CYTHION 5

AN EMULSIFIABLE CONCENTRATE CONTAINING 5 LB. LOW ODOR PREMIUM GRADE MALATHION PER GALLON.

ACTIVE INGREDIENTS:

Malathion (0, 0-dimethyl dithiophosphate of diethyl mercaptosuccinate)	56.6 %
Xylene	35.4 %
INERT INGREDIENTS:	8.0 %
Total	100.0 %

EPA Reg. No. 1598-190

CAUTION KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist or vapor. Wash thoroughly after using. SEE REMAINDER OF CAUTION STATEMENT ON SIDE PANEL.

NET VOLUME 1 U. S. Gallon
CYTHION IS A REGISTERED TRADE MARK OF AMERICAN CYANAMID COMPANY

Manufactured by



effective control of certain pests that are harmful to Ornamental Plants, with the exceptions noted.

Mix 1 1/4 teaspoons per gallon of water and uniformly spray the entire plant to control APHIDS, JAPANESE BEETLES, LEAFHOPPERS, MEALYBUG, TARNISH PLANT BUG, THRIP and WHITE FLY.

Mix 3 1/2 teaspoons per gallon of water and uniformly spray the entire plant to control AZALEA SCALE, EUONYMUS SCALE, JUNIPER SCALE, MONTEREY SCALE, OYSTER SHELL SCALE, SCURFY SCALE, SOFT BROWN SCALE, BOXWOOD LEAFMINER AND TWO-SPOTTED SPIDER MITES.

SCALE INSECTS are easier to control if insecticide is applied when "crawlers" are present. Weekly applications during the season may be necessary.

DO NOT USE on Boston Maidenhair, or Pteris Ferns, or on Petunias. May cause injury to Crassula.

CAUTION

Do not contaminate feed or foodstuffs. This product is toxic to fish. Keep out of any body of water. Apply only for the purposes and in the manner directed in the label. Do not apply when weather conditions favor drift of spray from area treated. Clean application equipment and dispose of rinse solutions, wastes, and unused chemical in a pit on non-crop land in an isolated area well away from water supplies.

This product is highly toxic to bees exposed to direct treatment or to residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. If container is glass, break into a previously prepared pit on non-crop land well away from water supplies, and bury fragments deeply. If container is metal, punch holes in top and bottom, crush under wheels of heavy vehicle or with an axe, and bury crushed container in a pit on non-crop land in an isolated area well away from water supplies.

DO NOT USE, STORE, POUR, OR SPILL NEAR HEAT OR OPEN FLAME. KEEP CONTAINER CLOSED WHEN NOT IN USE.

NOTICE

Follow directions carefully. Timing and method of application, weather and crop conditions and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions on this label.

1598-190

REC'D

NOV 9 1972

U.S. DEPARTMENT OF AGRICULTURE
FEDERAL INSECTICIDE
AND RODENTICIDE ACT
TOXIC SUBSTANCE CONTROL
BOARD
REGISTERED UNDER NO. 1598-190 SUBJECT
TO ATTACHED COMMENTS.