



FCX SEVIN® ZINEB 3-5.2 "D"



ACTIVE INGREDIENTS

Carbaryl (1-naphthyl N-methylcarbamate)	3.00
Zineb (zinc ethylene bis dithiocarbamate)	5.20

INERT INGREDIENTS

	91.80
TOTAL	100.00

Sevin is the Trademark of Union Carbide Corp. for the Active ingredient 1-naphthyl N-methylcarbamate.

USDA Reg. No. 1598-196

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if inhaled or swallowed. May cause irritation of nose and throat. Do not breathe dust. Do not take internally. Skin contact may be harmful. Avoid contact with skin. Wash hands before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing daily. Do not contaminate feed and foodstuffs.

NOTE FOR PHYSICIANS: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

For protection of honey bees avoid application during flowering or pollinating period when bees may be visiting crop. If necessary to apply during such periods, warn bee keepers well in advance to locate hives at a safe distance until one week after the application.

DESTROY EMPTY CONTAINERS: Do not reuse this container for any purpose. Burn empty containers (paper) in a hot fire, outdoors, in an isolated area. The smoke may be hazardous. Stay well away from the smoke.

DIRECTIONS FOR USE

To control the insects and the plant diseases on the crops listed below, apply at the rate specified for each crop to upper and lower surfaces of the foliage. Dust coverage of foliage should be uniform and thorough, but do not apply excessive amounts. Begin

applications when the insects or their damage first appear and before diseases appear (when disease is first reported to the area). Repeat as necessary, generally at 7 day intervals unless a shorter interval is stated. Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration.

TOMATOES Tomato fruitworm, tomato hornworm, leathoppers, European corn borer, fall armyworm, tarnished plant bug, Colorado potato beetle, early blight, late blight, leaf spot, and anthracnose on field grown tomatoes. Apply 40 to 50 pounds of dust per acre.

BEETS Flea beetles, tarnished plant bug, stink bug, cabbage looper (when larvae are small, apply at 4 to 5 day intervals), and leaf spot. Apply 30 to 40 pounds of dust per acre. **DO NOT APPLY TO BEETS WITHIN 14 DAYS OF HARVEST IF TOPS ARE TO BE USED AS FOOD OR FEED. IF TOPS WILL NOT BE USED, DO NOT APPLY WITHIN 3 DAYS OF HARVEST.**

EGGPLANT AND PEPPERS Leathoppers, fall armyworm, European corn borer, tarnished plant bug, leaf spot, and phomopsis. Apply 30 to 40 pounds of dust per acre.


BEANS Mexican bean beetle, bean leaf beetle, leathoppers, rust, anthracnose, and downy mildew (on lima beans). Apply 30 to 40 pounds of dust per acre. **DO NOT APPLY TO BEANS WITHIN 7 DAYS OF HARVEST.**

CABBAGE, CAULIFLOWER, AND BROCCOLI Harlequin bug, imported cabbage worm, larvae of diamond back moth, cabbage looper (when larvae are small, apply at 4 to 5 day intervals) and downy mildew. Apply 30 to 40 pounds of dust per acre. **DO NOT APPLY WITHIN 7 DAYS OF HARVEST.**

CUCUMBERS, PUMPKINS, AND SQUASH Pickle worm, melon worm, cucumber beetles, flea beetles, leathoppers, squash bug, downy mildew, Alternaria (Macrosporium) blight (on cucumbers) and anthracnose (on squash). Apply 30 to 33 pounds of dust per acre.

NOTICE

Follow directions carefully. Timing and method of application, weather and crop conditions and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material in strict accordance with directions given on this label.



Net Contents 25 U.S. Pounds

MANUFACTURED BY

FCX, Inc. RALEIGH, N.C.