




DANGER - POISON
PRECAUTIONS

POISONOUS IF SWALLOWED

This product can kill you if swallowed even in small amounts. Spray mist may be fatal if swallowed.



POISONOUS BY SKIN CONTACT

Poisonous if touched by hands or spilled or splashed on skin in eyes or on clothing. Liquid goes through clothes.



POISONOUS IF BREATHED

Breathing vapors or spray mist may be fatal.

POISONOUS

TO FISH AND WILDLIFE
The product is toxic to fish, birds, and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams, or ponds.

USE PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

USE ONLY WHEN WEARING THE FOLLOWING PROTECTIVE EQUIPMENT AND PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN

NET VOLUME

CONTENTS



STOP

READ LABEL BEFORE USING!

POISON
See side panels
for antidote
and precautions



DANGER
Keep out of reach
of children

CAN KILL YOU

METHYL PARATOX 63

DO NOT STORE OR USE
IN OR AROUND THE HOME

EPA EST. 1598-21
F. A. Reg. No. 1598-21

INGREDIENTS

Methyl Parathion

Inert

Water

Surfactant

Stabilizer

Emulsifier

Colorant

Odorant

Antifoam

Preservative

Buffer

Chelator

Sequestering Agent

Surfactant

Emulsifier

Colorant

Odorant

Antifoam

Preservative

Buffer

Chelator

Sequestering Agent

Surfactant

Emulsifier

Colorant

Odorant

Antifoam

Preservative

Manufactured by

FCX, Inc.

POISON
 DANGER
 READ LABEL BEFORE USING!
 CAN KILL YOU
 DO NOT STORE OR USE IN OR AROUND THE HOME
 EPA EST. 1598 SC-1
 EPA Reg. No. 1598-212
 INGREDIENTS
 ACTIVE INGREDIENTS
 Toxaphene 52.6%
 O,O Dimethyl O-p-nitrophenyl phosphorothioate 26.3%
 Xylene 16.1%
 INERT INGREDIENTS 5.0%
 Total 100.0%
 * Technical chlorinated camphene (chlorine content 67% to 69%)
 ** Methyl parathion
 Contains 6 Pounds TOXAPHENE and 3 Pounds METHYL PARATHION Per Gallon
 NET VOLUME GALLONS
 Manufactured By
FCX, Inc.



METHYL PARATOX 63

DO NOT STORE OR USE IN OR AROUND THE HOME

EPA EST. 1598 SC-1
 EPA Reg. No. 1598-212

INGREDIENTS

ACTIVE INGREDIENTS	
Toxaphene *	52.6%
O,O Dimethyl O-p-nitrophenyl phosphorothioate **	26.3%
Xylene	16.1%
INERT INGREDIENTS	5.0%
Total	100.0%

* Technical chlorinated camphene (chlorine content 67% to 69%)
 ** Methyl parathion
 Contains 6 Pounds TOXAPHENE and 3 Pounds METHYL PARATHION Per Gallon

Do not attempt to use this product unless you will comply with all the information provided in the label attached to the container.

NET VOLUME GALLONS

Manufactured By



NOTE TO PHYSICIAN

ANTIDOTE-Administer atropine sulfate in large doses. TWO to FOUR mg. intravenously or intramuscularly as soon as cyanosis is over come. Repeat at 5 to 10 minute intervals until signs of atropinization appear. 2 PAM chloride is also antidotal and may be administered in conjunction with atropine. **DO NOT GIVE MORPHINE OR TRANQUILIZERS.** Methyl parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement. **VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.**

DIRECTIONS FOR USE

This product must be diluted with water for application as a spray. To prepare the spray mixture, proceed as follows: 1. Introduce into the spray tank about one half of the total amount of water to be used. 2. Start agitation or pump bypass, and then add required amount of this product. 3. With agitation continuing, add remainder of the water, and then continue agitation during spray application.
 This product may be applied with ground equipment or with aircraft. To apply with ground equipment, prepare the spray mixture in a volume of water sufficient for uniformly thorough spray coverage of the foliage, usually 10 to 12 gallons of water per acre for low volume boom sprayers. To apply with aircraft, prepare the spray mixture in a volume of water which will assure application of at least 3 to 4 gallons per acre.
COTTON: During the early season, beginning at the two leaf stage, apply at the rate of 1/4 to 1 pint per acre to control thrips, apply at the rate of 1 to 1 1/2 pints per acre to control plant bugs and fleahoppers, apply at the rate of 1/2 to 1 pints per acre to control cutworms. Two or more applications made at intervals of 7 days apart may be required at this stage, depending upon the occurrence of insect populations and weather conditions favorable to their development, but do not apply unless damaging populations of insects are actually determined to be present in a particular field.
 During the mid season, beginning at the time squares form, apply at the rate of 2 to 2 1/2 pints per acre to control late wintering boll weevils, adults, plant bugs, aphids, garden webworms, and cotton leafworms. Two or more applications made at intervals of 7 days apart are likely to be required at the mid season stage, particularly if late wintering boll weevils are to be controlled, depending upon the occurrence of insect populations and weather conditions favorable to their development, so that the necessity for and time of repeated applications cannot be accurately predicted in advance. Do not apply unless potentially damaging populations of insects are actually determined to be present in each field.
 During the late season, beginning at the time a substantial number of bolls have formed, apply at the rate of 3 to 4 pints per acre to control boll weevils, cotton worms, garden webworms, and ichids. Apply at the higher rate for heavy bollworm outbreaks. Usually the applications should be repeated at 5 to 7 day intervals. 5 day intervals if bollworms are present in outbreak proportions. Two or more repeated applications are likely to be required at this stage, depending primarily upon weather conditions favorable to the development of the insects, but do not apply unless potentially damaging populations of insects are actually determined to be present in each particular field.

WHEAT, OATS, AND BARLEY: Apply within 21 days of handpicking or harvest. DO NOT GRAZE OR FEED WHEAT, OATS, OR BARLEY TO DAIRY ANIMALS OR ANIMALS BEING FINISHED FOR SLAUGHTER. DO NOT FEED WHEAT, OATS, OR BARLEY TO LIVESTOCK OR POULTRY.

SOYBEANS: Apply at the rate of 2 1/2 pints per acre to control aphids, armyworms, bean leaf beetle, corn ear worm, grasshopper, green cloverworm, stink bugs, and velvet bean caterpillars. Apply when a potentially damaging population of one or more of the insects is determined to be present in each particular field. Repeated applications may be necessary if the initial application is properly timed in relation to the actual presence of insects and the stage of growth of the soybeans in each particular field. However, if weather conditions are dry, the application may be necessary and continued development of the insects, a second application may be needed in 7 to 10 days after the first treatment. Do not apply more than twice per growing season.

DO NOT APPLY TO SOYBEANS WITHIN 21 DAYS OF HARVEST. DO NOT GRAZE OR FEED SOYBEAN PLANTS TO OTHER FORAGE FROM TREATED FIELDS OR ENSILAGE MADE FROM THESE SOURCES TO DAIRY, BEEF, OR ANIMALS BEING FINISHED FOR SLAUGHTER. DO NOT FEED WHEAT, OATS, OR BARLEY TO LIVESTOCK OR POULTRY.

WHEAT, OATS, AND BARLEY: Apply at the rate of 2 pints per acre to control cutworms. Apply when caterpillars are beginning to appear in small quantities in the field. Do not apply more than once per growing season. Repeated applications may be necessary if the initial treatment is made at the proper time in relation to the presence of cutworms and the stage of growth of these small grains.

DO NOT GRAZE OR FEED WHEAT, OATS, OR BARLEY TO DAIRY ANIMALS OR ANIMALS BEING FINISHED FOR SLAUGHTER. DO NOT FEED WHEAT, OATS, OR BARLEY TO LIVESTOCK OR POULTRY.

Keep away from children and animals. Do not get on face and away from the eyes, when spraying or pouring. If it gets in eyes, wash immediately with water. If it gets on skin, wash immediately with soap and water. Do not get on face and away from the eyes, when spraying or pouring. If it gets in eyes, wash immediately with water. If it gets on skin, wash immediately with soap and water.

Do not store in a place where children or animals can get at it. Do not apply to areas near water. Do not apply to areas where children or animals are present. Do not apply to areas where children or animals are present.

Do not store in a place where children or animals can get at it. Do not apply to areas near water. Do not apply to areas where children or animals are present. Do not apply to areas where children or animals are present.

Do not store in a place where children or animals can get at it. Do not apply to areas near water. Do not apply to areas where children or animals are present. Do not apply to areas where children or animals are present.

DESTROY EMPTY CONTAINERS. Do not reuse for any purpose. Do not burn, do not put in fire. Do not reuse for any purpose. Do not burn, do not put in fire. Do not reuse for any purpose. Do not burn, do not put in fire.

NOTICE

Do not use on crops or plants which are sensitive to this product. Do not use on crops or plants which are sensitive to this product. Do not use on crops or plants which are sensitive to this product.