

ACCEPTED

1598-215

1-23-75

UNITED STATES FEDERAL INSECTICIDE  
 AND FUNGICIDE ACT  
 FOR ESTABLISHED FLOWERS



# GREEN SHRUB WEEDER

Prevents weed growth in flower beds, shrub areas and vegetables



Contains Treflan\*

**ACTIVE INGREDIENTS:** Trifluralin (a,a,a trifluoro-2,6-dinitro-N,N-dipropyl-p-touidine)..... 93%  
**INERT INGREDIENTS**..... 99.07%  
 Total..... 100.00%

\***TREFLAN:** The Trade Mark for Trifluralin - Elanco Products Co., A Division of Eli Lilly & Co.

EPA Reg. No. 1598-215

EPA EST 1598-SC-1

## CAUTION

**KEEP OUT OF REACH OF CHILDREN**  
 (See remainder of Warning statements following directions)

### GREEN SHRUB WEEDER

For weed control in flower beds, shrub areas and vegetables. Stops broadleaf weeds\* and grass seeds\* as they sprout. Will not control established weeds, grasses or perennial roots.

For Flower Bed and Shrub Areas

This product may be applied to trees, shrubs and most established flowers\*\* Do not apply to gladiolus, aiuga, myrtle, pachysandra, and ornamental ground covers as injury may result.

### DIRECTIONS:

Before applying remove all existing broadleaf and grassy weeds, since this product kills sprouting seed but not established plants. Where mixing into the soil is impractical, apply uniformly around established flowers and shrubs with a cyclone or similar type spreader, making sure that the contents of this bag covers 5000 square feet. An additional similar application should be made about four weeks after the first application for season long weed control. The product will provide greater coverage where mixing with the soil is possible. This type of application can be made with good results by following directions for mixing into soil.

**\*ESTABLISHED FLOWERS:** Recommended for use on: Achillea, Ageratum, Arctotis, Aster, Balsam, Calendula, California Poppy, Calliopsis, Carnation, Centaurea, Chrysanthemum, Cosmos, Dahlia, Dianthus, Dimorphotheca, Forget-me-not, Four O'Clock, Gaillardia, Hollyhock, Ixora (Coccinea), Lobelia, Lupines, Marigold, Morning Glory, Nasturtium, Nicotiana, Periwinkle, Petunia, Phlox, Portulaca, Rudbeckia, Salvia, Scabiosa, Shasta Daisy, Snapdragon, Stocks, Snow on the Mountain, Sunflower, Sweet Alyssum, Sweet Pea, Sweet William, Zinnia.

**DIRECTIONS FOR MIXING INTO SOIL:** Apply to established trees, ornamentals, roses, and flowers that are 2 to 3 inches or more in height. Apply uniformly at the recommended rate. The rate of application varies with soil type. Use the table below to determine the proper rate of your type soil. Mix into the soil surface with hand rake being careful not to disturb the roots of established plants.

Shallow cultivation during the growing season will not reduce effectiveness when the product has been adequate mixed into the soil.

**FOR VEGETABLES:** Apply this product thoroughly and uniformly at the recommended rates over the entire area. For large areas, use a seeder or broadcast applicator. Immediately after application, mix into the soil to a depth of 1 to 3 inches using a rotary tiller or hand rake. Mixing should be done in two directions, at right angles, to insure thorough incorporation with the soil. Use only on vegetables listed below.

### SPECIAL INSTRUCTIONS FOR VEGETABLES

#### DIRECT SEEDED

Carrots

Okra

Southern Peas

(Cowpeas)

Field Peas

Black eyed Peas

Apply and incorporate into the soil before planting, anytime from 10 weeks before planting up to planting these vegetables from seed. See application rate table for proper rate for your type of soil.

#### DIRECT SEEDED

Lima Beans

Snap Beans

Apply and incorporate into the soil before planting, anytime from 10 weeks before planting up to planting. In light and medium soils apply at the rate of 25 lb. per 20,000 square feet. In heavy soils apply at the rate of 25 lb. per 15,000 square feet.

#### TRANSPLANTS

Broccoli

Brussel Sprouts

Cabbage

Cauliflower

Peppers

Tomatoes

Apply and mix into soil before transplanting. Do not apply after transplanting. See application rate table for proper rate for your type of soil.

### GENERAL APPLICATION RATES WHEN MIXING THIS PRODUCT INTO SOIL

SOIL TEXTURE	Light	Medium	Heavy
	Sandy & Sandy Loam	Loam	Silt & Clay
APPLICATION RATES	25 lb. per 20,000 sq. ft.	25 lb. per 15,000 sq. ft.	25 lb. per 10,000 sq. ft.

\***ANNUAL GRASSES:** Annual Bluegrass, Barnyardgrass (watergrass), Brachiaria, Bromegrass, Cheat, Crabgrasses, Foxtails, Goosegrass, Johnsongrass (from seed), Junglerice, Sandbur, Stinkgrass, Sprangletop, Texas Panicum, Wildcane.

Carelessweed, Carpetweed, Chickweed, Florida Purglane (Pusley), Goosefoot, Knotweed, Kochia, Lambsquarters, Pigweeds (spiny, redroot), Purslane, Russian Thistle, Stinging nettle.

**NOTE:** Avoid over application since this product may cause injury to crops sensitive to TREFLAN planted the following season.

### CAUTION

Harmful if swallowed. Avoid contact with skin.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not reuse container. Destroy when empty.

### NOTICE

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given on this label.

NET WEIGHT \_\_\_\_\_ POUNDS

MANUFACTURED BY

**FCX, Inc.** RALEIGH, N.C. 27602