

CAUTION

Harmful if taken internally. Avoid contamination of teed and foodstuffs, milking utensils or equipment, water and watering devices, and feed troughs.

Do not wet the skin of animals or poultry, since this may caus blistering. Do not apply to plants or vegetation

This pladuct is toxic to fish and birds. Do not contaminate water by cleaning of equipment or disposal of wastes

DESTROY EMPTY CONTAINERS. Do not reuse this container for any purpose. Promptyly dispose of empty container in refuse which will be buried in a landfill disposal dump, or bury empty container after being perforated and crushed in an isolated area well away from water supplies.

NOTICE

Follow directions carefully. Timing and method of application weather and crop conditions, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given on this label.

ACCEPTED:

AUG 3 1973

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER
ED UNDER NO
TO ATTACHED COMENT



MIST SPRAY

For Mist Spraying by Mechanical Atomization Without Heat or Steam

INGREDIENTS

CTIVE INGREDIENTS	
Technical piperonyl butowide	3 0
Pyrethrins	6
Petroleum basillori	96 .:
Total	190-6

Equivalent to 2 4 percent (but (carbity)) 6 prop (piperons) 6 ther and 0.6 percent related compounds

CAUTION KEEP OUT OF REACH OF CHILDREN

Sec Remainder of CAUTION Statement on Side Panel EPA Reg. No. 1898-225

Net Volume

Manufactured By

' Gallor



DIRECTIONS FOR USE

The oduct is a ready to use spray solution for "foggers, and mist erators which mechanically dispense a finely atomized mist hout using heat or steam, to control house files, stable files, it horn mes in dairy barns, poultry houses, and other livestic, quarters, when applied over the animals or poultry as a finely dimized mist.

BAR AND POULTRY HOUSES—To control house flies stable flies of thorn flies, apply as a fine mist over the livestock in stanch. Sor in holding pens, or over the poultry in cage or in pens to a fogger or mist generator for each 2,500 sq. ft. or less of floor as a placed above the animals or birds and preferably near the center of the area. Doors and windows should be closed, for to fogging openings may permit escape of the mist. Operate the generator unit until the mist is thoroughly distributed throughout the entire area, spray out 1 fl. oz. uniformly per 1,000 cu. ft. Do not ventilate the area for at least 10 minutes after the unit has been stupped it used in a dairy barn, areate the barn thoroughly before milking. Repeat daily morning, and afternooil, or as needed. Keep generator unit reservoir and nozzle clean to enjure proper delivery of spray mist or fog.

IMPORTANT INFORMATION—This product used as directed is intended to i. If adult flies immediately present in the treated area which are contacted by the mist. It does not have any effect on flies which subsequently migrate into the area or on fly ey, and larvae in brieding media on the premises and in surrounding areas. These sources of flies should be cleaned up, and sanitation procedures should be circled, as a primary requisite in a fix control program.

AYAYAYAYAYAYAYAYAYAYAYAYA