



FCX 2% Methomyl Insecticide Dust



For Control of Certain Leaf-Eating Insects on Cabbage and Tobacco Plants

ACTIVE INGREDIENT: Methomyl	
S-methyl-N-(methylcarbamoyl) oxy-	
thioacetimide	2.0 %
INERT INGREDIENTS	98.0 %
TOTAL	100.0 %

EPA Reg. No. 1598-230

EPA EST. 1598-SC-1

ACCEPTED
12-7-74

WARNING

KEEP OUT OF REACH OF CHILDREN

DO NOT USE OR STORE IN OR AROUND HOME

HARMFUL IF SWALLOWED OR INHALED

Do not breathe dust. Do not get in eyes, on skin, on clothing. In case of contact, wash skin with plenty of soap and water. For eyes, flush with water for 15 minutes and get medical attention. Remove and wash contaminated clothing before re use. Wear goggles and a mask or respirator of a type suitable for protection against insecticide dust when loading, applying or otherwise handling. Wear clean clothes daily. Wash thoroughly after handling and before eating or smoking. Do not contaminate or store near food or foodstuffs.

ANTIDOTE AND FIRST AID

ATROPINE IS AN ANTIDOTE. CONSULT PHYSICIAN FOR EMERGENCY SUPPLY OF 1/400 GRAIN ATROPINE TABLETS. CALL A PHYSICIAN AT ONCE IN ALL CASES OF REPORTED POISONING.

IF SWALLOWED, give a tablespoonful of salt in a glass of warm water and repeat until vomit is clear. **IF INHALED**, remove from exposure. Have patient lie down and keep quiet. If patient is not breathing, start artificial respiration immediately. **Never give anything by mouth to an unconscious person.**

IF WARNING SYMPTOMS APPEAR (See warning symptoms below) before physician arrives, immediately swallow two atropine tablets (each 1/100 gr.), thereafter, every 10 to 15 minutes, take one atropine tablet (1/100 gr.) until throat becomes dry and skin becomes dry and flushed. Take additional tablets as necessary to maintain a moderately dry throat and dry, flushed skin until physician arrives.

NOTE TO PHYSICIAN

WARNING SYMPTOMS - Methomyl poisoning produces effects associated with anti-cholinesterase activity which may include weakness, blurred vision, headache, nausea, abdominal cramps, discomfort in the chest, constriction of pupils, sweating, slow pulse, muscle tremors.

TREATMENT - Atropine sulphate should be used for treatment. Administer repeated doses, 1.2 to 2.0 mg intravenously every 10 to 30 minutes until full atropinization is achieved. Maintain atropinization until the patient recovers. Artificial respiration or oxygen may be necessary. Allow no further exposure to any cholinesterase inhibitor until recovery is assured. Do not use morphine or 2 PAM.

Do not use or store in or around the home. Keep unprotected persons and beneficial animals out of operational areas during treatment and while there is danger of drift.

This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and it should not be applied when bees are actively visiting the area.

DIRECTIONS FOR USE

FCX 2% Methomyl Insecticide Dust may be used to control certain insects on cabbage and tobacco. Apply only with rotary hand operated dust application equipment or with power driven ground dusting equipment. Use only in commercial plantings, do not use in home plantings.

APPLICATION Apply at the rates specified below when insects first appear. Repeat applications at 5 to 7 day intervals, or less frequently as needed. Apply the low rates on small plants, small worms and light infestations of insects. Use intermediate rates on larger worms and heavier infestations of insects. Use 1 to 3 applications of the highest recommended rate for controlling severe infestations. Thereafter, use the lowest rate which is adequate to maintain control.

CABBAGE - Cabbage looper, imported cabbage worm, diamondback moth larvae: 12 to 45 lbs/acre. **DO NOT APPLY TO CABBAGE WITHIN 1 DAY OF HARVEST.**

TOBACCO (Except Shade Tobacco) - Flea Beetle: 12 to 23 lbs/acre, begin when plants are set in the field. Hornworm: 12 to 23 lbs/acre, begin before largest worms exceed 1 1/2" in length. Cabbage Looper: 23 lbs/acre, begin when worms first appear. Budworm: 23 lbs/acre, include 2 to 3 applications before flower buds open. Direct dust into buds. **DO NOT APPLY TO BLUE CURED TOBACCO WITHIN 10 DAYS OF HARVEST. DO NOT APPLY TO WILD BLUE CURED TOBACCO WITHIN 14 DAYS OF HARVEST.**

DESTROY EMPTY CONTAINERS Do not reuse this container for any purpose. Bury empty container in a pit for non-crop land in an isolated area well away from water supplies.

MANUFACTURED BY

NET WEIGHT

FCX, Inc. RALEIGH N.C. 27602

25 POUNDS