

DIRECTIONS FOR USE

FOLIAGE APPLICATION Apply the dust when insects or their damage first appear on the plants and repeat applications as needed to maintain effective control. For best results apply the dust in the early morning or late evening when the temperature is lower and the air is still. Apply the dust uniformly and thoroughly so that a thin film of dust covers both the upper and the lower surfaces of all foliage. Do not use on food crops within the number of days of harvest shown in parentheses () after each crop.

- BEANS** Lima, Pole, Snap (7) Aphids, Mexican bean beetle, leafhoppers
- BROCCOLI** (5), **CABBAGE** (7), **CAULIFLOWER** (5) Aphids, diamond back moth, larva, imported cabbage worm
- CANTALOUPE** (3), **CUCUMBER** (7), **MUSKMELON** (3), **WATERMELON** (3), **SUMMER SQUASH** (3), **WINTER SQUASH** (3) Aphids, mites, thrips
- CORN** Sweet (2) Corn earworm. Apply to stalks as soon as they appear. Make applications at 4- to 6-day intervals. Reapply immediately after rain.
- LETTUCE** (10), **ENDIVE** (10) Aphids, leaf miners
- ONIONS** (10) Onion thrips
- PEAS** (9), **PEPPERS** (5), **PARSNIPS** (10), **RADISHES** (10), **SPINACH** (10), **TOMATOES** (3), **POTATOES** (14) Aphids, band-a-banana beetle, fall armyworm, southern armyworm, leaf miners, vinegar flies, Drosophila sp.
- BUFF BERBERIS** (7) Rubberry maggot
- GRAPE** (18) Grape berry moth, grape leaf folder, leafhoppers, Pacific spider mite
- SHRUBS** (4), (5) Aphids, leafhoppers, two spotted mite. For Cycadaria, mix dust with water and spray with a fine nozzle to ensure thorough coverage.
- FLOWERS, SHRUBS, ORNAMENTALS** Aphids, sawflies, leafhoppers, thrips, mites, scale insects, weevils, whiteflies, scale insects, thrips, mealybugs, etc. Do not use on plants which are susceptible to phytotoxicity. Do not use on plants which are susceptible to phytotoxicity.
- SOIL APPLICATION - PRE-PLANTING ONLY** To control cutworms and wireworms prevent soil-borne transmission of the following crops:

Asparagus	Celery	Onion	Spinach
Beans	Corn	Parsley	Swiss chard
Cantaloupe	Cucumbers	Peas	Tomatoes
Corn	Endive	Peppers	Watermelons
Corn cobs	Kale	Potatoes	
Cucumbers	Lettuce	Squash	

Apply 1/2 ounce of dust per broadcast treatment over 100 square feet of soil to be planted at the rate of 1/2 ounce per 100 square feet (10 x 20' area) immediately after application. For row crops, apply the dust to the soil to the depth and distance of 2 to 3 inches (2 to 3' row) on both sides of the row. For cutworms, wireworms, etc. apply 1/2 ounce of dust per 100 square feet of soil.

FOR ROOT MAGGOTS on broccoli, cabbage, and cauliflower, before transplanting, in formaldehyde, apply 1 pound of dust over 800 square feet (20 x 40' of area) to be planted, and mix the dust thoroughly into the top 3 to 4 inches of soil, or apply in the furrow at the rate of 1 ounce (2 tablespoons) per 100 feet of row and mix with the soil, or 1/2 ounce of dust. For radishes, apply 3 ounces (12 tablespoons) per 100 feet of row in the furrow at the time of planting.

ORNAMENTAL LAWNS Apply the amount of dust indicated on the label for 1000 square feet of lawn, ornamental only, to control:

Chinch bugs, Journeaux (12 tablespoons), cutworms, etc. (1/2 ounce per 1000 sq. ft. of grass), apply 7 ounces of dust.

Ants, armyworms, clover mites, springtails, etc. (1/2 ounce per 1000 sq. ft. of grass), apply 7 ounces of dust.

Beetle mites, chiggers, fleas, leafhoppers, etc. (1/2 ounce per 1000 sq. ft. of grass), apply 7 ounces of dust.

Dug-out spots and isolated ant hills, apply the dust liberally directly onto and around the spots and into the nest openings.

KEEP CHILDREN AND PETS OFF TREATED AREA UNTIL THE INSECTICIDE HAS BEEN WORKED INTO THE SOIL AND THE GRASS IS DRY.

NOTICE

Read the label carefully. Do not use this product if the label is missing or if the product is not in its original container. Do not use this product if the label is missing or if the product is not in its original container. Do not use this product if the label is missing or if the product is not in its original container.



GARDEN INSECTICIDE DUST
Contains 5% DIAZINON[®]

INGREDIENTS


ACTIVE INGREDIENT:	O,O diethyl O (2-isopropyl 6-methyl 4-pyrimidinyl) phosphorothioate	5%
INERT INGREDIENTS		95%
	Total	100%

* DIAZINON[®] - Trademark of CIBA-GEIGY Corporation

EPA Reg. No. 1598-236
EPA EST. 1598 S.C. 1

CAUTION

Keep Out of Reach of Children



Read the label carefully. Do not use this product if the label is missing or if the product is not in its original container. Do not use this product if the label is missing or if the product is not in its original container. Do not use this product if the label is missing or if the product is not in its original container.

MANUFACTURED BY **FCX** CORPORATION

RALEIGH, N.C. 27602