



**FCX
FLY BAN 7.76 RABON
ORAL LARVICIDE
TOP DRESS**

To prevent the development of house fly larvae and other fly larvae in manure.

ACTIVE INGREDIENT
2-chloro-1-(1,4-dichlorophenyl)-3-dimethyl phosphorane

INERT INGREDIENT

**KEEP OUT
OF REACH OF CHILDREN
CAUTION**

See below for additional precautionary statements.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS
CAUTION**

Harmful if swallowed. Irritant to the skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritated, consult a physician.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not contaminate streams or ponds. Do not contaminate water by cleaning of equipment or disposal of waste.

**SEE BACK OF LABEL
FOR DIRECTIONS FOR USE**

Manufactured By
FCX, Inc., Farmville, North Carolina

ACCEPTED
Net Weight 50 Lbs.
EPA Reg. No. 1598-242
EPA Est. No. 01598-NC-04
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 1598-242

DIRECTIONS FOR USE GENERAL CLASSIFICATION

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

When fed to cattle, PABON® Oral Larvicide passes through the digestive system into the animal's manure where it kills fly larvae on contact shortly after fly eggs hatch. By preventing larval development, PABON® Oral Larvicide helps to control horn flies and face flies on pasture and house flies and stable flies in and around cattle lots.

Fly Ban 7.76 Rabon Oral Larvicide Top Dress will mix uniformly in cattle feeds when standard mixing procedures are followed. Thus usual problems that are common to all feed preparation and which cause stratification, such as excessive free-fall or excessive handling, are to be avoided. It is recommended that appropriate preblending techniques be employed to assure adequate distribution of Fly Ban 7.76 Rabon Oral Larvicide Top Dress throughout the feed mix. The premix should be preblended with ground shays, protein supplements, mineral mixes, etc. before being added to roughages such as chopped hay or silage. Mixing time should be adequate to assure uniform dispersion. Optimum performance can be assured only if Fly Ban 7.76 Rabon Oral Larvicide Top Dress is dispersed uniformly with the feedstuff at the recommended level. Common feed mixing equipment (i.e., vertical mixer, horizontal blenders, mixer/feeder truck) may be used to prepare formulated feeds.

Feeds prepared with Fly Ban 7.76 Rabon Oral Larvicide Top Dress should not be pelleted, nor be mixed with feeds containing predominantly pellets. Further, Fly Ban 7.76 Rabon Oral Larvicide Top Dress should not be mixed in liquid feed supplements.

In A Concentrate Feed -- Roughage Fed Separately
Fly Ban 7.76 Rabon Oral Larvicide Top Dress can be used to prepare concentrate feeds that will provide 100 mg of PABON per animal per day.

To prepare a larvicidal concentrate feed, mix Fly Ban 7.76 Rabon Oral Larvicide Top Dress according to the amount of concentrate to be fed per animal per day. Use the following table as a guide for determining the proper mixing rate.

MIXING DIRECTIONS		
Pounds of Concentrate Consumed per Animal Per Day	PABON	
	Pounds of Concentrate	Pounds of Fly Ban 7.76 Rabon Oral Larvicide Top Dress per Ton of Concentrate
5	0.3500	90.0
10	0.1800	45.0
15	0.1200	30.0
20	0.0870	22.5
25	0.0350	9.0
30	0.0180	4.5
35	0.0120	3.0
40	0.0087	2.3
45	0.0063	1.8
50	0.0059	1.5

Feed the appropriate larvicidal concentrate to cattle weighing between 400 and 1200 pounds. For larger cattle weighing between 1200-1700 pounds, increase the amount of concentrate fed to 1 1/2 times that recommended.

Start feeding PABON larvicidal feeds early in the spring before flies begin to appear and continue feeding throughout the summer and into the fall until cold weather restricts fly activity.

Rations containing Fly Ban 7.76 Rabon Oral Larvicide Top Dress may be fed up to slaughter and to lactating dairy cows without withholding the milk from market during or after treatment. Fly Ban 7.76 Rabon Oral Larvicide Top Dress prevents the development of Horn Flies, Face Flies, House Flies and Stable Flies in the manure of treated cattle but is not effective against existing adult flies. In some cases, supplementary fly control measures may be needed in and around cattle lots and barns to control adult House Flies and Stable Flies which can breed not only in cattle manure but other decaying vegetable matter or silage on the premises. In order to achieve optimum fly control, Fly Ban 7.76 Rabon Oral Larvicide Top Dress should be used in conjunction with other good management and sanitation practices.

In A Complete Ration -- No Other Roughage Fed.
Fly Ban 7.76 Rabon Oral Larvicide Top Dress can be used to prepare rations that contain 25.4 mg of PABON per pound of complete ration.

To prepare a larvicidal ration, mix 1.5 pounds of Fly Ban 7.76 Rabon Oral Larvicide Top Dress per ton of complete mixed ration containing both grain and roughage. Full-feed this larvicidal complete ration to feeder cattle weighing from 400 to 1400 pounds and to dairy cattle at a rate to sustain milk production, but not less than 2.6 pounds of the ration per 100 pounds of body weight daily.

STORAGE AND DISPOSAL

Do not contaminate water or food by storage or disposal. Open dumping is prohibited. Store in an incinerator or landfill approved for pesticide containers, or bury in a safe place.