

[1621-2]  
2 pages


3/2

1621

**CONTAINS  
PYRETHRUM**

USDA Reg.No. 1621-2

CAUTION:



**CAUTION - COMBUSTIBLE MIXTURE**

N.Y.F.D. C. of A. No. 1404

**DIRECTIONS FOR USE**

**QUANTITY:** For exposed stages of common industrial and institutional insect pests, use 1 gallon per 50,000 cubic feet of space. Increase quantity for hardy insects more difficult to kill. Rapidity of initial kill may be accelerated by increasing dosage.

**FLYING INSECTS:** For control of flying insects such as flies, mosquitoes, gnats, and flying grain moths, direct spray or fog toward the upper portion of the enclosure, distributing throughout space.

**CRAWLING INSECTS:** To control crawling insects such as grain weevils, flour beetles, rice weevils, ants, and roaches, direct mist toward and into all places commonly inhabited by these insects, and spray liberally all protected hiding and breeding places.

**FOOD PRODUCTS INSECTS:** Spray the outside of packaged cereals or food; spraying shelves and other places where the pests are seen to alight, crawl, and congregate except on exposed food. Avoid sources of reinfestation by destroying wormy flour, grain, and cereals before applications.

For maximum effectiveness, keep doors and windows closed for 15 minutes after spraying. Repeat applications as necessary.

For best results, it is recommended that this insecticide be applied with mechanical aerosol generators or foggers, fog applicators, electric, steam, or compressed air atomizers in accordance with equipment manufacturers instructions. This is a non-corrosive solution that can be used in any

Space spraying in food processing plants should be done when plant is not in operation. Used as a contact spray, food handling surfaces should be covered during treatment or cleaned before re-use.

**CAUTION: HARMFUL IF SWALLOWED.**

**FIRE HAZARD: DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME.**

**TANGLEFOOT**