


FEB 14 1949

with letter

GROSS 101 LBS.
TARE 1 LB.
NET 100 LBS.

ACCEPTED
Mar. 1, 1949
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PESTS
ED UNDER NO. 1624-1
TO ATTACHED COMMENTS

**PACIFIC COAST
BORAX CO.**



BORASCU

FOR WEED CONTROL PURPOSES

ACTIVE INGREDIENT: BORAX, $\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O}$ 93%
INERT INGREDIENTS: 7%
GUARANTEED MINIMUM BORON TRIOXIDE, B_2O_3 34%

- CAUTION -

AVOID CONTAMINATION OF FOODSTUFFS
KEEP AWAY FROM CHILDREN AND DOMESTIC ANIMALS

GENERAL DIRECTIONS FOR USE

Borascu is a dry, white, crystalline solid. Subsequent rainfall or irrigation is necessary to dissolve the material and render it available for absorption by the roots of weeds. It becomes effective.

HOW TO APPLY: Borascu may be applied by a mechanical applicator, such as a fertilizer spreader, adjusted to deliver at a pre-determined rate. Where the use of a spreader is not feasible, weighed amounts may be scattered by hand over the weed areas. For a uniform distribution is essential to uniform control. If weed growth is prostrate, such as to interfere with even distribution on the soil surface, remove such material before applying Borascu. On steep slopes or in regions of excessive rainfall, raking into the soil or lightly covering with soil or gravel assists in retaining the Borascu in place.

WHEN TO APPLY: Any time of year when rains may be expected, but best results on annual and biennial weeds are obtained by applying when plants are young and tender. For deep-rooted perennials, fall or winter applications are recommended.

DOSE RATE: For general weed and grass growth, the following minimum rate of application, in pounds of Borascu per 100 square feet of ground area, are recommended: Eastern States, 8 to 10 pounds. All others, 11 to 12 pounds. Experience with field conditions has shown that the following weeds have been controlled by the average rates of application in pounds per 100 square feet as indicated:

Klamath Weeds (St. Johnswort)
Poison Ivy

2 1/2 - 3 1/2 lbs.
4 - 5 lbs.

Leafy Spurge
Canada Thistle
Wild Morning Glory, Wild Oats, Rye Grass
Johnson Grass, Poison Oak

6 - 8 lbs.
8 - 10 lbs.
11 - 12 lbs.
12 - 15 lbs.

It is advisable to consult local agricultural authorities on specific problems. Borascu is not recommended on Russian Thistle.

MAINTENANCE: If necessary, make spot applications where weeds re-appear, or general re-treatment using from one-half to full amount of initial rate of application. Such subsequent treatments tend to build up an area of unproductive soil, thereby minimizing re-infestations.

CAUTION

As this material is not selective in action and affects all types of vegetation, care should be taken to avoid contact with lawns, trees, shrubs and other growing plants. It is not intended to destroy. This includes precautions in treating areas which may be underlain by roots of adjacent valuable growths.

NOTICE

The use of this material being subject to conditions beyond their control, neither the manufacturer nor the seller makes any representation or warranty, express or implied, with respect to results from such use, whether or not in accordance with directions. The buyer accepts and uses this material subject to these terms and shall not hold the manufacturer or seller liable for the results of any use of this material.

YORK AS ANGELS

2/2