
MANUFACTURERS OF INDUSTRIAL MAINTENANCE MATERIALS



iune 5-1964

BODE AND ROBENTICIDE ACT FOI ECONOMIC POISON REGISTER-ED HITE NO LE SUBJECT TO ATTACHED COMMENTS.

U.S. Patent No. 3,032,405

Nonselective Weed and Grass Killer

FORMULA

Active Ingredients Sodium Metaborate Tetrahydrate (Na₂B₂O₄·4H₂O) . . (Boron Trioxide (B₂O₃) Equivalent . . 22.7%) Sodium Chlorate (NaClO₃) 30.00% Diuron [3-(3, 4-dichlorophenyl)-1, 1-dimethylurea] . 1.25%

CAUTION

Avoid contamination of feed and food-66.50% stuffs. Do not leave container where children or animals may take internally. Harmful if swallowed. Avoid contact with skin and clothing. Keep out of the reach of children.

Net Wt. - 125 lbs.



ICO INDUSTRIAL PRODUCT



13311 UNION AVENUE

CLEVELAND, OHIO 44120

SALES OFFICES IN PRINCIPAL CITIES

RGR

APPLICATION — For all uses, uniform application is essential. NEVRGREEN may be applied either as a dry granular material or as a water spray. Dissolve in water at rates up to 4 pounds per gallon. Both spray and dry applications become effective as soil treatments following rainfall or irrigation. For DRY application, conventional herbicide spreaders may be used. The hand operated NEVRGREEN Spreader is especially recommended for use where power equipment is not desired.

For SPRAY application use equipment with mechanical agitation. Add up to 4 pounds NEVRGREEN per gallon of water, depending on job requirements, to spray tank. Continue agitation until the herbicide is completely dissolved and during spraying operation. To minimize fire hazard due to dry yegetation, treat-

ment while weeds are small is suggested. GENERAL VEGETATION CONTROL - NEVRGREEN will give quick knock-down of vegetation and residual control of broadleaved weeds and grasses on parking lots, fence lines, highway and railroad sites, storage yards and other similar areas. For spray applications, add 1/2 to 2 pounds of NEVRGREEN to sufficient water to cover 100 square feet and apply as a contact spray thoroughly wetting all parts of the plants, particularly the growing tips 2 yaquare feet is usually adequate for young growth up to 1/2 to 1 gallon per square feet is usually adequate for young growth up to a jacket in higher rates may be necessary to insure thorough coverage of heavy get or to provide extended residual control. For DRY granular treater to 2 pound per 100 square feet. The lighter rates should be used of weed emergence on moist soil or when precipitation is antic weeds are still young. Persistent perennial weeds are most effectively regized when nearing maturity or during dormant stages in the fall. Spring may also be made under moderate rainfall conditions. Summer are recommended only where rainfall is dependable for leaching into the soil. Apply 2 to 4 pounds per 100 square feet in the dry grant or as a spray. On deep-rooted perennials such as bindweed, Can Bermuda grass and quack grass the heavier rates may be required. JOHNSON GRASS CONTROL — NEVRGREEN is particularly effective rooted perennial grasses such as Johnson grass and Dallis grass on sites, roadsides, fence lines and storage yards. Apply dry at 1 to 3 p 100 square feet or add 1 to 3 pounds to 1 or 2 gallons of water to a

treating after heading to late fall.

I M P O R T A N T

This material is not selective in action and may be taxic to all types of vegetally and may render the entire treated area totally or partially unproductive to or more years. Care should be taken to confine the use or application to particular area intended to be treated and to avoid its contact with lawns, shrubs, crops and other desirable plants which are not intended to be dest or injured. This includes precautions in treating areas which may be under to by roots of adjacent valuable growths. Careless application of this materia; washing by water run-off, to areas where desirable plants are growing which will be used for later planting may result in injury to desirable plants. Do not drain or flush equipment on or near such areas. Do not contaminate domestic or irrigation water. Wash clothing and equipment after application. Keep animals off treated areas until rainfall has leached the herbicide into the soil. Mechanical equipment should be thoroughly cleaned after

cover 100 square feet. Maximum efficiency on Johnson grass is obto

NOTICE WE

No responsibility is assumed by manufacturer or seller for unproductivity of the soil, injury to plant life or other results of the use of this material whether or not used or applied in accordance with directions. Because of the many varying conditions affecting use and application, such as location, seasonal weather, soil composition and type of vegetation, manufacturer makes no representation or warranty with respect to results or effects of such use whether or not used or applied in accordance with directions herein contained. Neither manufacturer nor seller makes, and their employees and agents are not permitted to make, any representations or warranties, express or implied by law, not specifically set forth on the label. Complete elimination of weeds is not guaranteed. The buyer accepts this material subject to these terms and assumes all risk of usage and handling. It is advisable to consult agricultural authorities or our field representatives on specific problems.