

5/11/77

20 MULE TEAM
162477

Apply to lawn and turf areas. The growth of the turf is retarded by an extremely fine, light green color of the grass, and by extensive control of weeds. Applications should be made throughout the growing season for effective broadleaf weed control.

RATES: On established turf and as listed broadleaf weed species, 2 to 3 gallons of MAINTAIN CF 125 (2 to 3 pounds active material, per acre, or a combination of 1 gallon of MAINTAIN CF 125 plus 1 gallon of MAINTAIN 3 (3 pounds maleic hydrazide) per acre in 50 gallons of spray solution are recommended. Even application to insure uniform wetting of grass and broadleaf weed foliage is essential for most effective results.

The combination gives the user the benefit of the separate retarding action of each material, plus an increase in effectiveness over either material used independently. MAINTAIN CF 125 should be well mixed with water before MAINTAIN 3 is added. Continue agitation during spraying operation.

Do not mow turf and weeds until all evidence of growth retardant has disappeared. Early mowing will greatly reduce the effectiveness of growth retardant treatment.

On vines such as A. ... of up to ... on deciduous broadleaves.

Gymnosperms (conifers, junipers, etc.) before buds expand in order to prevent the normal growth. Treatment when buds are tender causes developing branches to drop.

RATES: 1 pint/100 gal of water

	1 lb. / 100 gal	2 lb. / 100 gal
Deciduous Hardwood		
Alder Red	2.5	
Ash	1	
Bottlebrush Black	6.8	
Elm Siberian	1	
Maple Silverleaf	6	
Yew	6	
Silver	4.8	
Willow Golden	2.2	
Gymnosperms		
Fir	1.5	
Juniper	1.5	
Pine	1.5	
Spruce Sitka	1.5	
Redwood	1.5	
Vines and Ground Cover		
Ice Plant	4	
Ivy, English	6	
English (Mahoe)	1.2	
Kudzu	6.8	
Shrub		
Abelia	1.2	
Aspidistra	1.3	
Calluna	1.5	
Crataegus	1.2	
Elaeagnus	1.2	
Euonymus	1.5	
Hibiscus	1.5	
Japanese	1.5	
Meibomia	1	
Osageoak	1.3	
Plumbago	1	
Prunella	1.5	
Xylocopa	1	

* Australian or Lodgespole

IMPORTANT

This material is not selective in action and may affect all types of vegetation. Care should be taken to avoid its use or application to the particular area intended to be treated, and to avoid its contact with desirable plants. Keep livestock off treated areas. Wash hands, face and clothing after application. Machinery and equipment should be thoroughly cleaned after use.

NOTICE

Because of many varying conditions affecting use and application, such as soil conditions, position and type of vegetation, manufacturer warrants only that to the best of his knowledge and belief the results or effects of the use of this product in any event will be in accordance with the label. It is advisable to consult your State Extension Service or Agricultural Experiment Station with respect to any problems. Neither manufacturer nor seller shall be held liable to any injury or damage resulting from the use of this product for a purpose not intended by the manufacturer, or from any application not intended hereon nor with respect to breach of any warranty, not expressly set forth herein. Buyer accepts this material subject to these terms, and assumes all risk of use and handling, except when used in accordance with this label. Neither manufacturer nor seller, maker, and their agents and agents are not permitted to make any representations or warranties, express or implied by law, not specifically set forth herein.

Contains Chlorfuretol, a Product of E. Merck, Darmstadt, West Germany. **Patents Pending.
 ©Trademark of United States Borax & Chemical Corporation
 © 1976 U.S.B. & C.C.

A 20 MULE TEAM Product
 A MEMBER OF THE RTZ GROUP
 United States Borax & Chemical Corporation
 3075 Wilshire Boulevard / Los Angeles, California 90005

20 MULE TEAM
 162477

MAINTAIN CF 125 is a growth retardant which causes portions of the plant which are young and tender and which are growing after treatment, thus causing the plant to remain green but growing very slowly for an extended period.

The persistence of the retarding effect of MAINTAIN CF 125 in the plant may last a full season, depending on the species, rate and time of application. The effect on herbaceous broadleaf species is generally of longer duration than on turf grasses.

Do not leave containers where children or animals may reach them. Avoid contamination of food and feed-stuffs.

MAINTAIN CF 125 is a growth retardant which causes portions of the plant which are young and tender and which are growing after treatment, thus causing the plant to remain green but growing very slowly for an extended period.

The persistence of the retarding effect of MAINTAIN CF 125 in the plant may last a full season, depending on the species, rate and time of application. The effect on herbaceous broadleaf species is generally of longer duration than on turf grasses.

Do not leave containers where children or animals may reach them. Avoid contamination of food and feed-stuffs.

MAINTAIN CF 125 is toxic to fish. Keep out of water. Apply this product only as directed on this label.

DO NOT REUSE CONTAINERS. Destroy contents.