

SUPPLEMENTAL LABELING

POST EMERGENCE APPLICATION OF COBEX IN COTTON CROPS IN ARIZONA AND ALABAMA ONLY


Apply COBEX in cotton at least 6 inches high, prior to weed emergence, and not later than 80 days before harvest. Direct the spray to the soil surface between the rows and do not allow spray or drift to contact cotton foliage or stems. Incorporate COBEX into the top 1½ to 2 inches of soil within 24 hours of application. A rolling cultivator (Lilliston type) or other implement that can be adjusted to the contours of the furrows and beds is suitable. Use the rate recommended on the COBEX label for the soil type. Make only one application of COBEX per year.

IMPORTANT

BEFORE USING COBEX, CAREFULLY READ THE COBEX LABEL AND FOLLOW THE PRECAUTIONS AND DIRECTIONS.

THIS BULLETIN CONTAINS NEW OR SUPPLEMENTAL DIRECTIONS FOR THE USE OF COBEX. USE THIS BULLETIN IN CONJUNCTION WITH THE REGULAR PACKAGE LABEL (EPA Reg. No. 1624-104-AA).

UNITED STATES BORAX & CHEMICAL CORPORATION
3075 Wilshire Boulevard
Los Angeles, California 90010



UNITED STATES BORAX & CHEMICAL CORPORATION
A MEMBER OF THE RTZ GROUP
3075 Wilshire Blvd. • Los Angeles, California 90010