

# cobex

SUPPLEMENTAL LABELING  
EPA REG. NO. 1624-104-AA  
COBEX HERBICIDE FOR ENGLISH PEAS

COBEX (dinitramine) applied preplant and incorporated in the top 1½ to 2 inches of the final seedbed will control barnyard grass, crabgrass, kochia, lambsquarters, nightshade (black), pigweed, purslane and smartweed (Pennsylvania) and reduces population of morningglory (annual), ragweed (annual) and velvetleaf at the following rates.


#### RECOMMENDED BROADCAST RATES PER ACRE

Soil Texture	COBEX 2EC	
	Organic Matter (0-2.5%)	Organic Matter (2.5-6.0%)
Coarse (sand, sandy loam, loamy sand)	1 1/3 pints	1 1/3 pints
Medium (loam, silt loam, silt, clay loam, silty clay, silty clay loam)	1 1/2 pints	1 1/2 to 2 pints
Fine (clay)	2 pints	2 pints

#### IMPORTANT

BEFORE USING COBEX, CAREFULLY READ THE COBEX LABEL (EPA Reg. No. 1624-104) AND FOLLOW THE PRECAUTIONS AND DIRECTIONS.

THIS BULLETIN CONTAINS NEW OR SUPPLEMENTAL DIRECTIONS FOR THE USE OF COBEX. USE THIS BULLETIN IN CONJUNCTION WITH THE REGULAR PACKAGE LABEL. (EPA Reg. No. 1624-104-AA).



8/76

**BORAX®**  
A MEMBER OF THE RTZ GROUP  
3075 WILSHIRE BLVD. • LOS ANGELES, CALIF. 90010

# cobex

SUPPLEMENTAL LABELING  
EPA REG. NO. 1624-104-AA  
COBEX<sup>(a)</sup>, PLUS KARMEX AS AN OVERLAY APPLICATION IN  
THE UNITED STATES EXCEPT ARIZONA AND CALIFORNIA  
FOR COTTON ONLY

In addition to those weeds controlled by COBEX AND Karmex alone, COBEX applied preplant incorporated treatment and followed by Karmex after planting and before cotton emerges, either as a broadcast or band application will give control of tall morning-glory (annual morningglory) and prickly sida.

#### COBEX FOLLOWED BY KARMEX OVERLAY APPLICATION

Apply COBEX as a preplant incorporated treatment at the soil rates. This treatment should be incorporated in the top 1½ to 2 inches of the final seedbed within 24 hours of application.

#### RECOMMENDED BROADCAST RATES PER ACRE

SOIL TEXTURE	COBEX 2EC
Coarse	1 1/3 pints
Medium	1 1/2 to 2 pints
Fine	2 2/3 pints

APPLY KARMEX AS A PREEMERGENCE TREATMENT AFTER PLANTING AS A SEPARATE OPERATION AT THE FOLLOWING BROADCAST RATES: FOR BAND TREATMENT USE PROPORTIONATELY LESS; FOR EXAMPLE, FOR 14 INCH BAND ON 42 INCH ROW, USE 1/3 OF THE BROADCAST RATE.

SOIL TEXTURE	KARMEX POW
Coarse	2/3 lb.
Medium	1 lb.
Fine	1 1/4 to 2 lbs.

Do not use Karmex on sand or loamy sand soils or exposed subsoils containing less than 1 1/2% organic matter.

#### PRECAUTIONS

Do not exceed the following amounts of Karmex per acre during a single crop season as injury to subsequent crops may result: 1 lb. on loamy sand; 1 1/2 lbs. on sandy loam; 2 lbs. on clay loam; 2 3/4 lbs. on clay.

#### IMPORTANT

THIS BULLETIN CONTAINS NEW OR SUPPLEMENTAL DIRECTIONS FOR THE USE OF COBEX AND KARMEX. BEFORE USING COBEX WITH A KARMEX OVERLAY, CAREFULLY READ THE REGULAR LABELS OF EACH PRODUCT AND FOLLOW THE PRECAUTIONS AND DIRECTIONS.

Karmex<sup>®</sup> - Registered Trademark of E. I. duPont de Nemours & Co.

**BORAX®**  
A MEMBER OF THE RTZ GROUP  
3075 WILSHIRE BLVD. • LOS ANGELES, CALIF. 90010