

CONTENTS

POUNDS

DIRECTIONS

Punch out perforated holes in sifter top for easy application.

For control of house flies in and around dairy barns, horse barns, dog kennels, livestock sheds, hog pens, poultry houses, for each 500 sq. ft. of floor space, sprinkle 2 ounces of FLY KILLER in a space about 6 feet x 2 feet in size on floor. This application will adequately bait an area of 500 sq. ft. and thus one pound FLY KILLER properly used that will treat approximately 4,000 sq. ft. Give particular attention to areas around doorways, window ledges, or other flat surfaces where flies are seen to congregate.

Calf pens and bull pens, sprinkle FLY KILLER at the rate of 1/4 lb. per 500 sq. ft. around the outside edges of pens to insure best and most economical results. For inside pens, clean all pens well. Before applying new bedding, sprinkle FLY KILLER at the rate of 1/4 lb. per 100 sq. ft. of pen space, and cover with bedding. This method will help control maggots.

For reducing fly populations in and around Milk Houses, Creameries, Ice Cream Stands and Canneries, sprinkle FLY KILLER on the outside near the loading docks, doorways or refuse containers. Use approximately 1/4 lb. for every 500 sq. ft. Garbage areas, outside camps and outside latrines, apply bait around doorways, refuse containers or directly on refuse piles. There is no set application rate, but apply enough to insure good coverage. Maintain sanitation.


In all cases application should be made every two or three days until fly population is greatly reduced, and then repeated as necessary. For successful fly control, fly breeding sites such as manure piles and other waste materials should be eliminated.

CAUTION

Avoid breathing of dust. Harmful if swallowed or absorbed through the skin. Wash hands immediately if exposed to Malathion. Do not contaminate feeds or apply to animals. Keep out of reach of children, livestock and poultry. Do not use in areas where the affected insect may fall into food, food containers or in food processing equipment. If poisoning should occur call a physician immediately. Atropine is antidotal.

WARRANTY: The manufacturer warrants this product to kill flies if used as directed, or it will refund purchase price upon delivery of the container with unused contents to the dealer. No other warranty is made either expressed or implied.

16
ACC
July
UNDER THE FE
FUNGICIDE AND
FOR ECONOMIC
ED UNDER NO.
TO ATTACHED C



**KILLS FLIES
IN DAIRY BARN,
HOG PENS,
POULTRY HOUSES,
GARBAGE AREAS,
AND LIVESTOCK
SHEDS.**



CAUTION KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL CAUTIONS.

FLY BAIT

Contains 2% Malathion


ACTIVE INGREDIENTS: Malathion* 2%
INERT INGREDIENTS 98%

*O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate.

CHICAGO **KNOX CHEMICAL COMPANY** ILLINOIS

CONTENTS

POUNDS



Punch out perforated holes in sifter top for easy application.

For control of house flies in and around dairy barns, horse barns, dog kennels, livestock sheds, hog pens, poultry houses, for each 500 sq. ft. of floor space, sprinkle 2 ounces of FLY KILLER in a space about 6 feet x 2 feet in size on floor. This application will adequately bait an area of 500 sq. ft. and thus one pound FLY KILLER properly used that will treat approximately 4,000 sq. ft. Give particular attention to areas around doorways, window ledges, or other flat surfaces where flies are seen to congregate.

Calf pens and bull pens, sprinkle FLY KILLER at the rate of 1/4 lb. per 500 sq. ft. around the outside edges of pens to insure best and most economical results. For inside pens, clean all pens well. Before applying new bedding, sprinkle FLY KILLER at the rate of 1/4 lb. per 100 sq. ft. of pen space, and cover with bedding. This method will help control maggots.

For reducing fly populations in and around Milk Houses, Creameries, Ice Cream Stands and Canneries, sprinkle FLY KILLER on the outside near the loading docks, doorways or refuse containers. Use approximately 1/4 lb. for every 500 sq. ft. Garbage areas, outside camps and outside latrines, apply bait around doorways, refuse containers or directly on refuse piles. There is no set application rate, but apply enough to insure good coverage. Maintain sanitation.

In all cases application should be made every two or three days until fly population is greatly reduced, and then repeated as necessary. For successful fly control, fly breeding sites such as manure piles and other waste materials should be eliminated.

Avoid breathing of dust. Harmful if swallowed or absorbed through the skin. Wash hands immediately if exposed to Malathion. Do not contaminate feeds or apply to animals. Keep out of reach of children, livestock and poultry. Do not use in areas where the affected insect may fall into food, food containers or in food processing equipment. If poisoning should occur call a physician immediately. Atropine is antidotal.

WARRANTY: The manufacturer warrants this product to kill flies if used as directed, or it will refund purchase price upon delivery of the container with unused contents to the dealer. No other warranty is made either expressed or implied.

**KILLS FLIES
IN DAIRY BARNs,
HOG PENS,
POULTRY HOUSES,
GARBAGE AREAS,
AND LIVESTOCK
SHEDS.**



CAUTION KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL CAUTIONS.

Contains 2% Malathion

ACTIVE INGREDIENTS: Malathion*	2%
INERT INGREDIENTS	98%

*O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate.