

Hillyard

ITEM 179

EPA Reg. No. 1658-20

Hil-phene II

DISINFECTANT-DETERGENT

FOR INDUSTRIAL, INSTITUTIONAL AND COMMERCIAL USE. FOR USE IN CLEANING AND DISINFECTING CONDUCTIVE FLOORING. ALSO SUITABLE FOR TREATING MANY OTHER HARD SURFACES SUCH AS FLOORING, WALLS, CEILINGS, WOODWORK, DOORS AND FURNITURE.



* VIRUCIDAL	•	DEODORANT
TUBERCULOCIDAL	•	PSEUDOMONACIDAL
STAPHYLOCIDAL	•	FUNGICIDAL

ACTIVE INGREDIENTS	14.00%
Ortho-phenyl-phenol 5%; Ortho-benzyl-para-chlorophenol 3%; Para-tertiary-amyl-phenol 1%; Sodium dodecylbenzenesulfonate 5%.	
INERT INGREDIENTS	86.00%
TOTAL INGREDIENTS	100.00%

DANGER KEEP OUT OF REACH OF CHILDREN
(SEE BACK PANEL FOR ANTIDOTE STATEMENT AND FURTHER PRECAUTIONS)

DIRECTIONS: Diluted solutions can be applied in any convenient manner such as mopping, flooding or wiping. In cases where excessive amounts of soil and contamination are present, clean before disinfecting for maximum effectiveness. The surface should be thoroughly wetted with the disinfecting solution and allowed to dry without rinsing. For cleaning and disinfecting surfaces suspected of being contaminated, use the following dilutions (parts Hil-phene II/parts water):

- 1-128 for Mycobacterium tuberculosis
- 1-192 for Pseudomonas aeruginosa
- 1-256 for routine use, Staphylococcus aureus, Salmonella choleraesuis and Tricophyton interdigitale.

*Virucidal Action - for inactivation of Asian Flu (Flu A₂ Taiwan) and Hong Kong Flu A₂ viruses on environmental surfaces, use a 1-192 Hil-phene II solution.

DILUTION TABLE:	1 GALLON	10 QUARTS	5 GALLONS
1-128 (0.8%)	1 ounce	2½ ounces	5 ounces
1-192 (0.5%)	¾ ounce	2 ounces	3¾ ounces
1-256 (0.4%)	½ ounce	1½ ounce	2½ ounce

DANGER: Keep Out of Reach of Children. Corrosive. Causes eye damage and severe skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Store in closed and properly labeled container. Do not re-use empty container. Rinse empty container thoroughly with water and discard it. Wash thoroughly after using. Avoid contamination of food, feed stuffs, processing equipment and water supplies.

173

EPA Reg. No. 1658-20



MANUFACTURED BY
HILLYARD CHEMICAL COMPANY
HOSPITAL PRODUCTS RESEARCH DIVISION
St. Joseph, Missouri 64502 U.S.A.

ACCEPTED