



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

May 13, 2020

Jennifer Brandon
Agent
James Austin Company
c/o Delta Analytical Corp.
12510 Prosperity Drive, Suite 160
Silver Spring, MD 20904

Subject: Label Amendment: Emerging Viral Pathogens Claim
Product Name: Austin A-1 Ultra Disinfecting Bleach
EPA Registration Number: 1672-65
Application Date: 04/06/2020
Decision Number: 561529

Dear Ms. Brandon:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Because you have opted to add statements pertaining to emerging viral pathogens to your label as described in the August 19, 2016, Guidance to Registrants: Process For Making Claims Against Emerging Viral Pathogens Not On EPA-Registered Disinfectant Labels ("Guidance"), https://www.epa.gov/sites/production/files/2016-09/documents/emerging_viral_pathogen_program_guidance_final_8_19_16_001_0.pdf, you are subject to the following additional terms of registration:

1. You may make statements pertaining to emerging viral pathogens only through the following communications outlets: technical literature distributed exclusively to health care facilities, physicians, nurses and public health officials, "1-800" consumer information services, social media sites and company websites (non-label related). These statements shall not appear on marketed (final print) product labels.


2. Your statements pertaining to emerging viral pathogens must adhere to the format approved on the Agency-accepted master label.
3. You may make statements pertaining to emerging viral pathogens only upon a disease outbreak that meets all the following criteria:
 - a. The causative organism must be a virus that causes an infectious disease that has appeared in a human or animal population in the U.S. for the first time, or that may have existed previously but is rapidly increasing in incidence or geographic range.
 - i. For human disease, the outbreak is listed in one of the following Centers for Disease Control (CDC) publications:
 - A. CDC Current Outbreak List for “U.S. Based Outbreaks” (www.cdc.gov/outbreaks),
 - B. CDC Current Outbreak List for “Outbreaks Affecting International Travelers” with an “Alert” or “Advisory” classification (www.cdc.gov/outbreaks) (also released through the CDC’s Health Alert Network (HAN) notification process)
 - C. Healthcare-Associated Infections (HAIs) Outbreaks and Patient Notifications page (www.cdc.gov/hai/outbreaks)
 - ii. For animal disease, the outbreak is identified as an infectious disease outbreak in animals within the U.S. on the World Organization for Animal Health (OIE) Weekly Disease Information page (www.oie.int/wahis_2/public/wahid.php/Diseaseinformation/WI).
 - A. The CDC or OIE has identified the taxonomy, including the viral family and/or species, of the pathogen and provides notice to the public of the identity of the emerging virus that is responsible for an infectious disease outbreak. Based on the taxonomy of the outbreak pathogen identified by the CDC or OEI, the pathogen's viral subgroup is small non-enveloped, large non-enveloped, enveloped.
 - B. The virus can be transmitted via environmental surfaces (non-vector transmission), and environmental surface disinfection has been recommended by the CDC, OIE or EPA to control the spread of the pathogen.
4. You may begin communicating statements pertaining to emerging viral pathogens only upon CDC or OIE’s publication per term 3.a. of an outbreak of an emerging viral pathogen meeting all of the criteria of term 3. You must cease and remove all such non-label communications intended for consumers no later than 24 months after the original publication of the outbreak per term 3.a., unless the Agency issue written guidance to the contrary due to continued public health concerns. The emerging pathogen claim language may remain on the master label.

5. Terms from points 1 through 4 above shall become immediately void and ineffective if registration for use against Hepatitis A and Canine Parvovirus is suspended or cancelled or no longer meets the criteria for a disinfectant claim (see EPA Product Performance Test Guideline 810.2200). In addition, terms B.1 through B.4 above shall become immediately void and ineffective upon your receipt of evidence of ineffectiveness against any pathogen in a less-resistant Spaulding category.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, you may contact the disinfectants list at disinfectantslist@epa.gov.

Sincerely,



Steven Snyderman, Acting Product Manager 33
Regulatory Management Branch 1
Antimicrobials Division (7510P)
Office of Pesticide Programs

Enclosure: stamped label

AUSTIN

A-1 ULTRA DISINFECTING BLEACH

(ABN's: Austin's A-1 Bleach, Austin's A-1 Concentrated Bleach, Crystal Bleach, Sno-ee Bleach, Elite Sanitizer, Wipe Away Sanitizer)

Commercial/Institutional/Household Use

DISINFECTS • SANITIZES • DEODORIZES • VIRUCIDE • TUBERCULOCIDE • GERMICIDE

- KILLS HIV AND TB ON HARD NON-POROUS ENVIRONMENTAL SURFACES
- FOR FOOD SERVICE APPLICATIONS • RESTAURANT USE • HOSPITAL USE
- NEW HIGHER CONCENTRATION FORMULA • REMOVES STAINS
- DISINFECTS AND DEODORIZES • WHITENS • BRIGHTENS LAUNDRY
- FOOD PROCESSING PLANT USE • FARM - DAIRY USE
- KILLS GERMS
- BACTERICIDAL
- MILDEWCIDAL
- FUNGICIDAL
- GERMICIDAL
- CONCENTRATED

This product may be used on hard non-porous surfaces in commercial, institutional, hospital and household premises (including kitchens, bathrooms, nurseries, sick rooms, laundry rooms) eating establishments, pet kennels, veterinary premises, farms, dairies, and food processing plants.

ACTIVE INGREDIENT:	
Sodium Hypochlorite.....	6.0%
OTHER INGREDIENTS: 94.0%	
TOTAL:	100.0%
Yields 5.7% available chlorine.	

Contains no phosphorus

KEEP OUT OF REACH OF CHILDREN
DANGER

See Back Panel for Additional Cautions.

ACCEPTED

05/13/2020

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 1672-65

FIRST AID	
If in eyes	<ul style="list-style-type: none"> •Hold eye open and rinse slowly and gently with water for 15-20 minutes. •Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
If on skin or clothing	<ul style="list-style-type: none"> •Take off contaminated clothing. •Rinse skin immediately with plenty of water for 15-20 minutes.
If inhaled	<ul style="list-style-type: none"> •Move person to fresh air. •If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
If swallowed	<ul style="list-style-type: none"> •Have person sip a glass of water if able to swallow. •Do not induce vomiting unless told to do so by the poison control center or doctor. •Do not give anything by mouth to an unconscious person.
NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.	
<p>Call poison control center or doctor immediately for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the Poison Control Center at 1-800-222-1222. Hotline: 1-866-359-5662</p>	

Manufactured by:

James Austin Company – PO Box 827 Mars PA 16046

MSDS: 1-800-245-1942 – Hotline: 1-866-359-5662

Net Contents: 96 fl. oz. (3 Qt.) (2.84 L) [other sizes permitted]

EPA Reg. No. 1672-65

EPA Est. No. 1672-PA-1, FL-1, MA-1 or NC-1

[Alternate EPA Establishments: May be jet coded or designated in the Lot# or Batch#]

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

DANGER: Corrosive. Causes skin burns and substantial but temporary eye injury. Do not get in eyes, on skin, or on clothing. Wear safety glasses and rubber gloves when handling this product. Harmful if inhaled. Avoid breathing vapors. Harmful if swallowed. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using toilet.

Environmental Hazards

[Containers less than five gallons]

This product is toxic to fish and other aquatic organisms.

[Five gallon containers and larger]

This product is toxic to fish and aquatic organisms. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

Physical and Chemical Hazards

Strong oxidizer. Flush drains before and after use. Do not use or mix with other household chemicals, such as toilet bowl cleaners, rust removers, acid or ammonia containing products. To do so will release hazardous gasses. Prolonged contact with metal may cause pitting or discoloration.

DO NOT MIX WITH AMMONIA - DO NOT MIX WITH ACIDS

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal. Store this product in a cool, dry area, away from direct sunlight and heat to avoid deterioration. In case of spill, flood area with large quantities of water. Product or rinsates that cannot be used should be diluted with water before disposal in a sanitary sewer. Do not re-use empty container, offer for recycling if available, or place in trash collection.

Do not contaminate food or feed by storage, disposal or cleaning of equipment.

[For containers one gallon or less]

STORAGE: Store in a cool, dry area away from direct sunlight and heat to avoid deterioration.

DISPOSAL: Nonrefillable container. Do not reuse empty container. Offer for recycling, if available, or place in trash collection.

- Bottle is coded for recycling where HDPE containers are accepted.
- Recyclable Container
- Bleach does not contain any phosphorus or phosphates.
- Not tested on animals.

[Bracketed text is optional]

GENERAL / CLEANING / STAIN REMOVAL / AND DEODORIZING CLAIMS

- Removes Stains
- Gets whites even whiter than before
- Removes and Eliminates Odors
- Deodorizer
- Deodorizes
- Whites just got whiter
- Whitens bleachable fabrics
- Whitens whites
- For a cleaner, fresher laundry and household
- Anti-Allergen (non-living)
- Boosts Cold Water Cleaning Power
- Boosts Cold Water Washing Power
- Brightens Laundry
- Do Not Break Open – To Be Sold As A *[Two] [or Three] [or Four]* Pack
- Easy Pour Spout
- Do Not Drop
- For *[Use in]* Standard *[or Top Load]* & HE *[or Front Load]* *[Washing]* Machines
- For *[Use in]* HE *[or Front Load]* *[Washing]* Machines
- For *[Use in]* Standard *[or Top Load]* *[Washing]* Machines
- For Cold Washing
- Safe in HE *[or Front loading]* Washing Machines
- Smaller Bottle Is Easier To handle and Store Than Before
- Made In The USA
- Sparkling White
- Reduces Non-living Allergens

GENERAL SANITIZING / DISINFECTING / VIRUCIDAL CLAIMS

- Removes mold *[and mildew]*
- Sanitizer
- Sanitizes
- Fungicidal *[Fungicide]*
- Germicidal *[Germicide]*
- Virucide *[Virucidal]*
- Remove *[bacteria] [germs] [viruses]* -and/or- *[body soil]*, detergent leaves behind.
- Powerful germ killer
- Gets Rid of Germs *[And Odors]*
- *[Helps]* Prevent*[s]* The Spread Of The Cold And Flu Virus†† *[In Your Home] [and/or Office]* from treated surfaces
- Kills mold *[and mildew]*
- Kills most mold
- Kills Athlete's Foot Fungus †† *[In 5 Minutes]*
- Cleans *[and disinfects]*
- Disinfects
- Disinfects and Deodorizes by Killing *[Most]* Germs and Their Odors
- Disinfects hard, nonporous surfaces
- Disinfects day care centers
- Eliminates germs and bacteria
- Kills surface germs and bacteria *[including Staphylococcus aureus (Staph) [ATCC 6538], [Salmonella choleraesuis] -or- Salmonella enterica [ATCC 10708] and [the] Influenza A2 [virus] [ATCC VR-544]]*

- Also kills in 10 minutes
Adenovirus Type 2,
Avian Influenza [virus] Type A [ATCC VR-2072],
Hepatitis [type] A [ATCC VR-2093] [strain HM-175],
Cytomegalovirus [ATCC VR-578] [Strain AD-169],
Respiratory Syncytial Virus [ATCC VR-26] [Strain Long],
Varicella zoster Virus [ATCC VR-586], [Strain G],
Herpes Simplex Virus [Type] 2 (Herpes) [ATCC VR-734] [Strain G],
Rubella virus (German measles virus) [ATCC VR-315],
Mycobacterium bovis [ATCC 35743] (Tuberculosis),
Shigella dysenteriae [ATCC 13313]
- [Cleans away] [Kills] [Eliminates] [Destroys] [Removes] [Wipes Away] [99.9% of] the bacteria, germs and viruses commonly found in kitchens, bathrooms, [restrooms] [households] [homes] [offices] [work] [environments] [areas] [laundry].
- [Fights] [Cleans away] [Kills] [Eliminates] [Destroys] [Removes] [Wipes Away] [99.9% of] the bacteria, germs and viruses that [can] cause the common cold or flu in your [household] [home] [office] [work place] [environment] [area] [school] [classroom].*
- Help prevent the spread of [the] cold and flu viruses [in your home -or-office].*
- Kill[s] [99.9% of] [common] [household -or-laundry] germs.*
- Kill[s] bacteria on the surfaces your kids touch every day.*
- Kills [99.9% of] [household] mold [and mildew].
- Kills [household] viruses that cause colds and the flu: Rhinovirus and Influenza A2.
- Kills Virus That Causes The Flu‡
- Kills [Eliminates] Flu Virus ‡
- Kills [Eliminates] MRSA †††
- Kills H1N1 Influenza A virus
- Kills Pandemic 2009 H1N1 Influenza A virus [Formerly Called Swine Flu]
- Kills Clostridium difficile [C. diff] On Hard Non-Porous [Environmental] Surfaces

‡ Influenza A virus –and/or - H1N1 Influenza A virus
 ‡‡ Influenza A virus –and/or - H1N1 Influenza A virus & Rhinovirus type 37
 †† Trichophyton mentagrophytes (Athlete's Foot Fungus) [ATCC 9533],
 ††† Methicillin Resistant Staphylococcus aureus (MRSA)

- **for front of product label**

Kills [99.9%] of [common household] germs*

- **for back of product label**

*Kills [99.9%] of [common household] germs: Staphylococcus aureus (Staph) [ATCC 6538], Streptococcus pyogenes (Strep) [ATCC 9342], [Salmonella choleraesuis] -or- Salmonella enterica (Salmonella) [ATCC 10708], Pseudomonas aeruginosa (Pseudomonas) [ATCC 15442], Influenza A2 (Hong Kong) [VR-544], Rhinovirus Type 37 [ATCC VR-1147] (virus that causes cold and flu), Trichophyton mentagrophytes (Athlete's Foot Fungus) [ATCC 9533], Rotavirus [Strain WA] [ATCC VR-2018], Escherichia coli O157:H7. (E. coli) [ATCC 35150] and Aspergillus niger [ATCC 6275]

- **for front of product label**

Kills [99.9%] of [common household] bacteria*

- **for back of product label**

*Kills [99.9%] of [common household] bacteria: Staphylococcus aureus (Staph) [ATCC 6538], Streptococcus pyogenes (Strep) [ATCC 9342], [Salmonella choleraesuis] -or- Salmonella enterica (Salmonella) [ATCC 10708]. Pseudomonas aeruginosa (Pseudomonas) [ATCC 15442] and Escherichia coli O157:H7 (E coli) [ATCC 35150]

Emerging Viral Pathogens Claims

[This product qualifies for emerging viral pathogen claims per EPA's "Guidance to Registrants: Process for Making Claims Against Emerging Viral Pathogens not on EPA-Registered Disinfectant Labels" when used in accordance with the appropriate use directions indicated below.]

[This product meets the criteria to make claims against emerging viral pathogens from the following viral categories:]

For an emerging viral pathogen that is a/an:	...follow the directions for use for the following organism on the label:
Enveloped Virus	Hepatitis A
Large Non-Enveloped Virus	Hepatitis A
Small Non-Enveloped Virus	Hepatitis A Canine Parvovirus

[Statements must adhere to one or both of the following formats:]

[Product name] has demonstrated effectiveness against viruses similar to **[name of emerging virus]** on hard, non-porous surfaces. Therefore, **[product name]** can be used against **[name of emerging virus]** when used in accordance with the directions for use [against **[name of supporting virus(es) – see table above]**] on hard non-porous surfaces. Refer to the **[CDC or OIE]** website at **[pathogen-specific website address]** for additional information.

[Name of illness/outbreak] is caused by **[name of emerging virus]**. **[Product name]** kills similar viruses and therefore can be used against **[name of emerging virus]** when used in accordance with the directions for use [against **[name of supporting virus(es) – see table above]**] on hard non-porous surfaces. Refer to the **[CDC or OIE]** website at **[website address]** for additional information.

[These statements shall not appear on marketplace (final printed) labels]

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

One Tbsp. ($\frac{1}{2}$ oz.) of this bleach in 1 gallon of water is equivalent to approximately 200 parts per million (ppm) available chlorine. Use a chlorine test kit to determine exact available chlorine concentration and adjust as necessary to obtain specified ppm.

LAUNDRY

Use to bleach white and colorfast Acrylics, Cotton, Nylon, Polyester, Rayon (test to be sure). Do not use on Acetate, Leather, Silk, Spandex or Wool.

BLEACH TEST: Before using, mix one tablespoon of bleach with $\frac{1}{2}$ cup of water in a glass or plastic container and test a small piece of fabric in a place that doesn't show.

[or]

To test-dye colorfastness, add tablespoon of Bleach to $\frac{1}{4}$ cup of water and then place one drop of the solution on the hidden part of a seam. Blot dry after 1 minute. No color change means the piece may be bleached safely. Do not bleach wool, silk, mohair, leather, spandex and non-fast colors.

Standard Washer [Top Load] $\frac{3}{4}$ cup **Extra Large Washer [Top Load]** 1 $\frac{1}{4}$ cups

Standard Washer [Front Load] $\frac{1}{2}$ cup

For HE Washers, add using the bleach dispenser following the machine manufacturer's instructions.

LAUNDRY: Sort laundry by color. Before adding clothes, mix $\frac{3}{4}$ cup of bleach with water in standard top-loading [16 gallon] machines or mix a $\frac{1}{2}$ cup of bleach with water in front-loading [8 gallon] machines. Add Bleach to dispenser when available, otherwise, add bleach and detergent to wash water before the soiled laundry is put in. For HE Washers, fill dispenser to maximum level. *[If sanitization is desired, soak clothes for 5 minutes. Wash and rinse with usual cycles.]* If clothes are in machine the addition of bleach can cause damage.

HAND WASHING: Rinse article to remove any loose soils and then soak in a mixture of $\frac{1}{8}$ cup bleach and 2 gallons of cool water.

REMOVE STAINS: Mix $\frac{1}{4}$ cup of bleach with a gallon of water. Soak stained area for 5 minutes to remove grass, ink, coffee, tea, scorch, fruit, etc. Rinse thoroughly.

Disinfects, sanitizes, and deodorizes by killing most germs and their odors.

DISINFECTING KITCHEN, DISHES, SINKS: Use $\frac{1}{4}$ cup bleach mixed with a quart of water to soak cleaned dishes, teapot, cups, sinks, etc. for 10 minutes. Rinse with a solution of approximately 1 Tbsp. of bleach per gallon of water to prepare a 200 ppm solution. Do not use on silverware. Bleach solution can be used on porcelain, enamel, etc. surfaces after cleaning. Let air dry.

DISINFECTING WALLS, FLOORS, AND OTHER HARD INANIMATE SURFACES NOT IN DIRECT CONTACT WITH FOOD: Prewash surfaces and rinse. Mix $\frac{3}{4}$ cup bleach per gallon of water. Spray, rinse, or wipe surface with bleach solution and let stand for 10 minutes. Drain and air dry.

DISINFECTING BATHROOM: Prewash toilet and flush. Pour $\frac{1}{2}$ cup bleach into toilet bowl, scrub with a brush, making sure to get under rim, let stand 10 minutes, flush. Do not use with bowl cleaners or any other household chemicals.

DEODORIZING GARBAGE CANS: Wash and rinse. Use $\frac{3}{4}$ cup bleach for each gallon of water in can. Empty and let drain.

EGG SHELL SANITIZING: Thoroughly clean eggs. Mix approximately 1 Tbsp. ($\frac{1}{2}$ oz) of bleach per gallon of warm water to produce a 200 ppm available chlorine solution. The sanitizer temperature should not exceed 130 degree F. Spray the warm sanitizer so that the eggs are thoroughly wetted. Allow the eggs to thoroughly dry before casing or breaking. Do not apply a potable rinse. The solution should not be re-used to sanitize eggs.

One Tbsp. (½ ounce) of this bleach in 1 gallon of water is equivalent to approximately 200 parts per million (ppm) available chlorine. Use a chlorine test kit to determine exact available chlorine concentration.

FOR SANITIZING	Amount Bleach	Amount Water	INSTRUCTIONS
Work Surfaces	1 Tbsp.	1 Gallon	Pre-wash with detergent, rinse, cover surface with bleach solution for at least 5 minutes, drain, let air dry.
Dishes, Glassware, Utensils	1 Tbsp.	1 Gallon	After washing, soak for at least 5 minutes in bleach solution. Drain and let air dry.
Bathtubs, Showers	1 Tbsp.	1 Gallon	Wash, rinse, apply bleach solution for at least 5 minutes, drain, let air dry.
Refrigerators, Freezers	1 Tbsp.	1 Gallon	Wash, rinse, apply bleach solution for at least 5 minutes, drain, let air dry.
FOR DISINFECTING			
Hard Surface Floors, Glazed Tiles & Walls	¾ Cup (6 oz)	1 Gallon	Pre-wash surfaces and rinse. Spray, rinse or wipe surface with bleach solution, let stand for 10 minutes. Drain and air dry.
Mops, Brushes, Brooms & Rugs	¾ Cup (6 oz)	1 Gallon	Wash with detergent, apply bleach solution, soak at least 10 minutes. Rinse well.
Bathtubs, Showers & Kitchen Sinks	¾ Cup (6 oz)	1 Gallon	Pre-wash surface and wipe with bleach solution. Allow solution to stand at least 10 minutes. Rinse well. Air dry.
Toilet Bowls	1 Cup (8 oz)		Flush Toilet. Pour 1 cup of bleach into bowl. Brush entire bowl, including rim, with a scrub brush. Let stand 10 minutes before flushing again.
FOR DEODORIZING			
Garbage Cans	¾ Cup (6 oz)	1 Gallon	After washing and rinsing, brush inside with solution. Empty and let drain.
Drains	1 Cup (8 oz)		Pour into drain. Flush with hot water.
FOR BLEACHING/WHITENING			
Wooden Surfaces	½ Cup (4 oz)	1 Gallon	Apply for 5 minutes, rinse.
FOR MOLD, MILDEW, & STAIN REMOVAL			
All Surfaces	½ Cup (4 oz)	1 Gallon	Add bleach to detergent solution, apply, rinse.
FOR AGRICULTURAL USES			
Food Egg	1 Tbsp.	1 Gallon	Wash, rinse, spray with 130 degree F max bleach solution until completely wet; let air dry before casing or breaking. Once solution is used it must not be reused for this purpose.
Fruits and Vegetables	1 Tbsp.	20 Gallons	Wash, rinse, soak in circulating bleach solution (25 ppm) for 2 min. Spray vegetables with bleach solution before packing; use potable water to rinse fruit before packaging. Rinse is to occur only immediately prior to packaging.

DISINFECTING HARD, NONPOROUS SURFACES: Kills in 10 minutes

Staphylococcus aureus (Staph) [ATCC 6538],
Streptococcus pyogenes (Strep) [ATCC 9342],
[Salmonella choleraesuis] -or- Salmonella enterica (Salmonella) [ATCC 10708],
Pseudomonas aeruginosa (Pseudomonas) [ATCC 15442],
Influenza A2 (Hong Kong) [VR-544],
Adenovirus Type 2,
Rhinovirus Type 37 [ATCC VR-1147] (viruses that cause colds and flus),
Trichophyton mentagrophytes (Athlete's Foot Fungus) [ATCC 9533],
Rotavirus [Strain WA] [ATCC VR-2018],
Hepatitis [Type] A [ATCC VR-2093] [strain HM-175],
Cytomegalovirus [ATCC VR-578] [Strain AD-169],
Respiratory Syncytial Virus [ATCC VR-26] [Strain Long],
Varicella zoster Virus [ATCC VR-586], [Strain G]
Herpes Simplex Virus [Type] 2 (Herpes) [ATCC VR-734] [Strain G],
Rubella virus [ATCC VR-315], (German measles virus),
Escherichia coli O157:H7 [ATCC 35150] (E. coli),
Mycobacterium bovis [ATCC 35743] (Tuberculosis),
Shigella dysenteriae [ATCC 13313],
Aspergillus niger [ATCC 6275] (mildew) –or- Aspergillus brasiliensis (mildew) [ATCC 16404],
Avian Influenza [virus] Type A [ATCC VR-2072].
Methicillin Resistant Staphylococcus aureus (MRSA) [ATCC 33592]
Clostridium difficile spore (*C. diff*) [#] [ATCC 700792]
Canine parvovirus [ATCC 953]
Feline parvovirus [ATCC VR-648]
H1N1 Influenza A virus [ATCC VR-1469, Strain A/PR/8/34]
Human coronavirus [ATCC VR-740]

If Clostridium difficile claim is used, Clostridium difficile directions must be included on the printed label.

Laundering: To bleach and sanitize white and colorfast cotton, linen, nylon, Dacron, Orlon, polyester, Dynel and rayon in washing machine: $\frac{3}{4}$ cup of this product per load for conventional washing machine; $\frac{1}{2}$ cup for front load automatic. Add to pre-soak, wash water or first rinse. If clothes are in machine, dilute product in 1 quart water before adding.

To Whiten Nylon and Other Synthetics that has turned yellow or grey. 1 tablespoon of this product per gallon water. Soak clean fabric in solution for 15 to 20 minutes. Rinse well. Repeat if necessary.

To Remove Stains. Berry, wine, coffee, tea, ink, grass, dye, medicine stains, scorch and mildew. Make solutions of 2 tablespoons of this product to each quart water. Immerse fabric for 5 to 10 minutes. Rinse well in clear water. Repeat if necessary.

Today's permanent Press Fabrics are Bleachable, and need this product to get out stains and help prevent dirt build up. Wash with regular laundry as directed: Top-load automatics – $\frac{3}{4}$ cup per load. Wringer-type washers – $\frac{3}{4}$ cup per load. Front-load automatics – $\frac{1}{2}$ cup per load. Use this product with any good laundry soap or detergent. If your washer has an automatic bleach dispenser, follow washer directions. If not, add this product to wash water before laundry is put in. If laundry is put in before wash water then dilute this product in quart of water and add after machine has started agitating and fabrics are thoroughly wet.

Disinfecting and Deodorizing Bathrooms: To disinfect, deodorize, and eliminate mold and mildew from washable surfaces such as tubs, showers, countertops, sinks, glazed ceramic tile and vinyl flooring, spread a solution of $1\frac{1}{2}$ cups of this product per 2 gallons of water on clean surface. Let stand 10 minutes, then drain.

Toilet Bowls: To sanitize and deodorize pre-cleaned toilet bowls, use $\frac{1}{2}$ cup of this product. Flush, pour in bleach – swab with brush, making sure to get under the rim, and let stand for 10 minutes. Flush. DO NOT use with bowl cleaners or any other household chemicals.

Sickroom Equipment – Wash all surfaces thoroughly. Rinse, then spread a solution of $1\frac{1}{2}$ cups of this product per 2 gallons of water over all surfaces. Let stand 10 minutes, then drain.

Garbage cans – Wash thoroughly with warm soapy solution. Rinse then spread a solution of $1\frac{1}{2}$ cups of this product per 2 gallons of water over all surfaces. Let stand 10 minutes, then drain.

Avoid prolonged contact with metal since corrosion or discoloration may occur. Do not use this product on chipped enamel.

Sanitizing Nonporous Food Contact Surfaces –

Prepare a sanitizing solution by thoroughly mixing 2 Tbsp. (1 oz.) of this product with 2 gallons of water to provide approximately 200 ppm available chlorine by weight. Clean all surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 2 minutes. Do not rinse equipment with water after treatment and do not soak equipment overnight.

To Sanitize Milking Equipment: Prepare sanitizing solution as above immediately prior to use. All surfaces to be sanitized should be properly cleaned before application of chlorine solution. Milking utensils should be submerged in the solution for at least 2 minutes and allowed to drain. Do not rinse equipment with water after treatment.

If solution contains less than 50 ppm available chlorine, as determined by a suitable test kit, either discard the solution or add sufficient product to reestablish a 200 ppm residual. Sanitizers used in automated systems may be used for general cleaning but may not be reused for sanitizing purposes.

Sanitizing Porous Food Contact Surfaces: Prepare a solution of approximately 600 ppm by thoroughly mixing 6 Tbsp. (3 oz) of this product with 2 gallons of water. Clean surfaces in the normal manner. Rinse all surfaces thoroughly with the 600 ppm solution, maintaining contact with the sanitizer for at least 2 minutes. Prepare a 200 ppm sanitizing solution by thoroughly mixing 2 Tbsp. (1 oz.) of this product with 2 gallons of water. Prior to using equipment, rinse all surfaces with 200 ppm available chlorine solution. Do not rinse with water and do not soak equipment overnight.

PROPORTIONS FOR DILUTION OF THIS PRODUCT

200 ppm: 1 oz. (2 Tbsp) in 2 gallons of water

600 ppm: 3 oz. (6 Tbsp) in 2 gallons of water

(Use a chlorine test kit to determine exact available chlorine concentration and adjust dosage as necessary.)

RESTAURANTS, TAVERNS, SODA FOUNTAINS, DAIRIES, ETC.

Directions for Sanitizing Eating and Drinking Utensils:

Prepare sanitizing solution immediately prior to use.

1. Scrape and pre-wash utensils and glass whenever possible.
2. Wash with good detergent or compatible cleaner.
3. Rinse with clean water.
4. Sanitize in solution of 1 oz to 2 gallons of water (200 ppm).
5. Immerse utensils at least 2 minutes or for contact time specified by governing sanitary code.
6. Do not reuse sanitizing solution.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

NOTE: This product degrades with age. Use a chlorine test kit and increase dosage, as necessary, to obtain the required level of available chlorine.

TABLE OF PROPORTIONS AVAILABLE CHLORINE

600 ppm – 12 oz. in 10 gal. water
200 ppm – 4 oz. in 10 gal. water

SWIMMING POOL WATER DISINFECTION

For a new pool or spring start-up, super chlorinate with 107 to 213 oz. of product for each 10,000 gallons of water to yield 5 to 10 ppm available chlorine by weight. Check the level of available chlorine with a test kit. Adjust and maintain pool water pH to between 7.2 to 7.6. Adjust and maintain the alkalinity of the pool to between 50 to 100 ppm.

To maintain the pool, add manually or by a feeder device 21 oz. of this product for each 10,000 gallons of water to yield an available chlorine residual between 0.6 to 1.0 ppm by weight. Stabilized pools should maintain a residual of 1.0 to 1.5 ppm available chlorine. Test the pH, available chlorine residual and alkalinity of the water frequently with appropriate test kits. Frequency of water treatment will depend upon temperature and number of swimmers.

Every 7 days, or as necessary, super chlorinate the pool with 107 to 213 oz. of product for each 10,000 gallons of water to yield 5 to 10 ppm available chlorine by weight. Check the level of available chlorine with a test kit. Re-entry into treated pools is prohibited above 4 ppm due to risk of bodily injury.

At the end of the swimming pool season or when water is to be drained from the pool, chlorine must be allowed to dissipate from treated pool water before discharge. Do not chlorinate the pool within 24 hours prior to discharge.

WINTERIZING POOLS - While water is still clear & clean, apply 6.5 oz. of product per 1000 gallons, while filter is running, to obtain 3 ppm available chlorine residual, as determined by a suitable test kit. Cover pool, prepare heater, filter and heater components for winter by following manufacturers' instructions.

SPAS, HOT-TUBS, IMMERSION TANKS, ETC.

SPAS/HOT-TUBS - Apply 11 oz. of product per 1000 gallons of water to obtain a free available chlorine concentration of 5 ppm, as determined by a suitable chlorine test kit. Adjust and maintain pool water pH to between 7.2 and 7.8. Some oils, lotions, fragrances, cleaners, etc. may cause foaming or cloudy water as well as reduce the efficiency of the product.

To maintain the water, apply 11 oz. of product per 1000 gallons of water over the surface to maintain a chlorine concentration of 5 ppm. Re-entry into treated pools is prohibited above 4 ppm due to risk of bodily injury.

After each use, shock treat with 17 oz. of this product per 500 gallons of water to control odor and algae.

During extended periods of disuse, add 6.5 oz. of product daily per 1000 gallons of water to maintain a 3 ppm chlorine concentration.

HUBBARD AND IMMERSION TANKS - Add 11 oz. of this product per 200 gallons of water before patient use to obtain a chlorine residual of 25 ppm, as determined by a suitable test kit. Adjust and maintain the water pH to between 7.2 and 7.6. After each use drain the tank. Add 11 oz. to a bucket of water and circulate this solution through the agitator of the tank for 15 minutes and then rinse out the solution. Clean tank thoroughly and dry with clean cloths. *[Not for use in California]*

HYDROTHERAPY TANKS - Add 2 oz. of this product per 1000 gallons of water to obtain a chlorine residual of 1 ppm, as determined by a suitable chlorine test kit. Pool should not be entered until the chlorine residual is below 3 ppm. Adjust and maintain the water pH to between 7.2 and 7.6. Operate pool filter continuously. Drain pool weekly, and clean before refilling.

SANITIZATION OF NONPOROUS FOOD CONTACT SURFACES

RINSE METHOD - A solution of 100 ppm available chlorine may be used in the sanitizing solution if a chlorine test kit is available. Solutions containing an initial concentration of 100 ppm available chlorine must be tested and adjusted periodically to ensure that the available chlorine does not drop below 50 ppm. Prepare a 100 ppm sanitizing solution by thoroughly mixing 2 oz. of this product with 10 gallons of water. If no test kit is available, prepare a sanitizing solution by thoroughly mixing 4 oz. of this product with 10 gallons of water to provide approximately 200 ppm available chlorine by weight.

Clean equipment surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 2 minutes. If solution contains less than 50 ppm available chlorine, as determined by a suitable test kit, either discard the solution or add sufficient product to reestablish a 200 ppm residual. Do not rinse equipment with water after treatment and do not soak equipment overnight.

Sanitizers used in automated systems may be used for general cleaning but may not be re-used for sanitizing purposes.

IMMERSION METHOD - A solution of 100 ppm available chlorine may be used in the sanitizing solution if a chlorine test kit is available. Solutions containing an initial concentration of 100 ppm available chlorine must be tested and adjusted periodically to ensure that the available chlorine does not drop below 50 ppm. Prepare a 100 ppm sanitizing solution by thoroughly mixing 2 oz. of this product with 10 gallons of water. If no test kit is available, prepare a sanitizing solution by thoroughly mixing 4 oz. of this product with 10 gallons of water to provide approximately 200 ppm available chlorine by weight.

Clean equipment in the normal manner. Prior to use, immerse equipment in the sanitizing solution for at least 2 minutes and allow the sanitizer to drain. If solution contains less than 50 ppm available chlorine, as determined by a suitable test kit, either discard the solution or add sufficient product to reestablish a 200 ppm residual. Do not rinse equipment with water after treatment.

Sanitizers used in automated systems may be used for general cleaning but may not be re-used for sanitizing purposes.

FLOW/PRESSURE METHOD - Disassemble equipment and thoroughly clean after use. Assemble equipment in operating position prior to use. Prepare a volume of a 200 ppm available chlorine sanitizing solution equal to 110% of volume capacity of the equipment by mixing the product in a ratio of 4 oz. product with 10 gallons of water. Pump solution through the system until full flow is obtained at all extremities, the system is completely filled with the sanitizer and all air is removed from the system. Close drain valves and hold under pressure for at least 2 minutes to ensure contact with all internal surfaces. Remove some cleaning solution from drain valve and test with a chlorine test kit. Repeat entire cleaning/sanitizing process if effluent contains less than 50 ppm available chlorine.

CLEAN-IN-PLACE METHOD - Thoroughly clean equipment after use. Prepare a volume of a 200 ppm available chlorine sanitizing solution equal to 110% of volume capacity of the equipment by mixing the product in a ratio of 4 oz. product with 10 gallons of water. Pump solution through the system until full flow is obtained at all extremities, the system is completely filled with the sanitizer and all air is removed from the system. Close drain valves and hold under pressure for at least 10 minutes to ensure contact with all internal surfaces. Remove some cleaning solution from drain valve and test with a chlorine test kit. Repeat entire cleaning/sanitizing process if effluent contains less than 50 ppm available chlorine.

SPRAY METHOD - Preclean all surfaces after use. Use a 200 ppm available chlorine solution to control bacteria, mold or fungi and a 600 ppm solution to control bacteriophage. Prepare a 200 ppm sanitizing solution of sufficient size by thoroughly mixing the product in a ratio of 4 oz. product with 10 gallons of water. Prepare a 600 ppm solution by thoroughly mixing the product in a ratio of 13 oz. product with 10 gallons of water. Use spray equipment which can resist hypochlorite solutions. Always empty and rinse spray equipment with potable water after use. Thoroughly spray all surfaces until wet, allowing excess sanitizer to drain. Vacate area for at least 2 hours. Prior to using equipment, rinse all surfaces treated with 600 ppm solution with a 200 ppm solution.

SANITIZING OF POROUS FOOD CONTACT SURFACES

RINSE METHOD - Prepare a sanitizing solution by thoroughly mixing 13 oz. of this product with 10 gallons of water to provide 600 ppm available chlorine by weight. Clean surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 2 minutes and allow the sanitizer to drain. Prepare a 200 ppm sanitizing solution by thoroughly mixing 4 oz. of this product with 10 gallons of water and rinse all surfaces with this 200 ppm solution. Do not rinse with water and do not soak equipment overnight.

IMMERSION METHOD - Prepare a sanitizing solution by thoroughly mixing, in an immersion tank, 13 oz. of this product with 10 gallons of water to provide 600 ppm available chlorine by weight. Clean equipment in the normal manner. Prior to use, immerse equipment in the sanitizing solution, maintaining contact for at least 2 minutes and allow the sanitizer to drain. Following this, prepare a 200 ppm sanitizing solution by thoroughly mixing 4 oz. of this product with 10 gallons of water and rinse all surfaces with this 200 ppm solution. Do not rinse with water and do not soak equipment overnight.

SPRAY METHOD - Preclean all surfaces after use. Prepare a 600 ppm available chlorine sanitizing solution of sufficient size by thoroughly mixing the product in a ratio of 13 oz. product with 10 gallons of water. Use spray equipment which can resist hypochlorite solutions. Always empty and rinse spray equipment with potable water after use. Thoroughly spray all surfaces until wet, allowing excess sanitizer to drain. Vacate area for at least 2 hours. Prior to using equipment, rinse all surfaces with a 200 ppm available chlorine solution. Prepare a 200 ppm sanitizing solution by thoroughly mixing 4 oz. of this product with 10 gallons of water.

SANITIZATION OF NONPOROUS NON-FOOD CONTACT SURFACES

RINSE METHOD - Prepare a sanitizing solution by thoroughly mixing 4 oz. of this product with 10 gallons of water to provide approximately 200 ppm available chlorine by weight. Clean equipment surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 5 minutes. Do not rinse equipment with water after treatment and do not soak equipment overnight.

IMMERSION METHOD - Prepare a sanitizing solution by thoroughly mixing, in an immersion tank, 4 oz. of this product with 10 gallons of water to provide approximately 200 ppm available chlorine by weight. Clean equipment in the normal manner. Prior to use, immerse equipment in the sanitizing solution for at least 5 minutes and allow the sanitizer to drain. Do not rinse equipment with water after treatment.

SPRAY METHOD - Preclean all surfaces after use. Prepare 200 ppm available chlorine sanitizing solution of sufficient size by thoroughly mixing the product in a ratio of 4oz. product with 10 gallons of water. Use spray equipment which can resist hypochlorite solutions. Prior to using equipment, thoroughly spray all surfaces until wet, allowing excess sanitizer to drain. Vacate area for at least 2 hours.

SANITIZATION OF POROUS NON-FOOD CONTACT SURFACES

RINSE METHOD - Prepare a sanitizing solution by thoroughly mixing 13 oz. of this product with 10 gallons of water to provide approximately 600 ppm available chlorine by weight. Clean surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 2 minutes. Do not rinse equipment with water after treatment and do not soak equipment overnight.

IMMERSION METHOD - Prepare a sanitizing solution by thoroughly mixing, in an immersion tank, 13 oz. of this product with 10 gallons of water to provide approximately 600 ppm available chlorine by weight. Clean equipment in the normal manner. Prior to use, immerse equipment in the sanitizing solution for at least 2 minutes and allow the sanitizer to drain. Do not rinse equipment with water after treatment.

SPRAY METHOD - After cleaning, sanitize non-food contact surfaces with 600 ppm available chlorine by thoroughly mixing the product in a ratio of 13 oz. of this product with 10 gallons of water. Use spray equipment which can resist hypochlorite solutions. Always empty and rinse spray equipment with potable water after use. Prior to using equipment, thoroughly spray all surfaces until wet, allowing excess sanitizer to drain. Vacate area for at least 2 hours.

SEWAGE & WASTEWATER EFFLUENT TREATMENT

The disinfection of sewage effluent must be evaluated by determining the total number of coliform bacteria and/or fecal coliform bacteria, as determined by the Most Probable Number (MPN) procedure, to ensure that the chlorinated effluent has been reduced to or below the maximum permitted by the controlling regulatory jurisdiction.

On the average, satisfactory disinfection of secondary wastewater effluent can be obtained when the chlorine residual is 0.5 ppm after 15 minutes contact. Although the chlorine residual is the critical factor in disinfection, the importance of correlating chlorine residual with bacterial kill must be emphasized. The MPN of the effluent, which is directly related to the water quality standards requirements, should be the final and primary standard and the chlorine residual should be considered an operating standard valid only to the extent verified by the coliform quality of the effluent.

The following are critical factors affecting wastewater disinfection.

1. **Mixing:** It is imperative that the product and the wastewater be instantaneously and completely flash mixed to assure reaction with every chemically active soluble and particulate component of the wastewater.
2. **Contacting:** Upon flash mixing, the flow through the system must be maintained.
3. **Dosage/Residual Control:** Successful disinfection is extremely dependant on response to fluctuating chlorine demand to maintain a predetermined, desirable chlorine level. Secondary effluent should contain 0.2 to 1.0 ppm chlorine residual after a 15 to 30 minute contact time. A reasonable average of residual chlorine is 0.5 ppm after 15 minutes contact time.

SEWAGE AND WASTEWATER TREATMENT

EFFLUENT SLIME CONTROL - Apply a 100 to 1000 ppm available chlorine solution at a location which will allow complete mixing. Prepare this solution by mixing 21 to 213 oz. of this product with 100 gallons of water. Once control is evident, apply a 15 ppm available chlorine solution. Prepare this solution by mixing 3 oz. of this product with 100 gallons of water.

FILTER BEDS - SLIME CONTROL: Remove filter from service, drain to a depth of 1 ft. above filter sand, and add 170 oz. of product per 20 sq/ft evenly over the surface. Wait 30 minutes before draining water to a level that is even with the top of the filter. Wait for 4 to 6 hours before completely draining and back washing filter.

DISINFECTION OF DRINKING WATER (EMERGENCY/PUBLIC/INDIVIDUAL/SYSTEMS)

PUBLIC SYSTEMS - Mix a ratio of 2 oz. of this product to 100 gallons of water. Begin feeding this solution with a hypochlorinator until free available chlorine residual of at least 0.2 ppm and no more than 0.6 ppm is attained throughout the distribution system. Check water frequently with a chlorine test kit. Bacteriological sampling must be conducted at a frequency no less than that prescribed by the National Primary Drinking Water Regulations. Contact your local Health Department for further details.

INDIVIDUAL SYSTEMS: DUG WELLS - Upon completion of the casing (lining), wash the interior of the casing (lining) with a 100 ppm available chlorine solution using a stiff brush. This solution can be made by thoroughly mixing 2 oz. of this product into 10 gallons of water. After covering the well, pour the sanitizing solution into the well through both the pipe sleeve opening and the pipeline. Wash the exterior of the pump cylinder also with the sanitizing solution. Start pump and pump water until strong odor of chlorine in water is noted. Stop pump and wait at least 24 hours. After 24 hours flush well until all traces of chlorine have been removed from the water. Consult your local Health Department for further details.

INDIVIDUAL WATER SYSTEMS: DRILLED, DRIVEN & BORED WELLS - Run pump until water is as free from turbidity as possible. Pour a 100 ppm available chlorine sanitizing solution into the well. This solution can be made by thoroughly mixing 2 oz. of this product into 10 gallons of water. Add 5 to 10 gallons of clean, chlorinated water to the well in order to force the sanitizer into the rock formation. Wash the exterior of the pump cylinder with the sanitizer. Drop pipeline into the well, start pump and pump water until strong odor of chlorine in water is noted. Stop pump and wait at least 24 hours. After 24 hours, flush well until all traces of chlorine have been removed from the water. Deep wells with high water levels may necessitate the use of special methods for introduction of the sanitizer to the well. Consult your local Health Department for further details.

INDIVIDUAL WATER SYSTEMS: FLOWING ARTESIAN WELLS - Artesian Wells generally do not require disinfection. If analyses indicate persistent contamination, the well should be disinfected. Consult your local Health Department for further details.

EMERGENCY DISINFECTION - When boiling of water for 1 minute is not practical, water can be made potable by using this product. Prior to addition of the sanitizer, remove all suspended material by filtration or by allowing it to settle to the bottom. Decant the clarified, contaminated water to a clean container and add 3 drops of this product to 20 gallons of water. Allow the treated water to stand for 30 minutes. Properly treated water should have a slight chlorine odor, if not, repeat dosage and allow the water to stand an additional 15 minutes. The treated water can then be made palatable by pouring it between clean containers for several times. This process has not been demonstrated to inactivate Cryptosporidium cysts.

PUBLIC WATER SYSTEMS

RESERVOIRS: ALGAE CONTROL - Hypo chlorinate streams feeding the reservoir. Suitable feeding points should be selected on each stream at least 50 yards upstream from the points of entry into the reservoir.

MAINS - Thoroughly flush section to be sanitized by discharging from hydrants. Permit a water flow of at least 2.5 feet per minute to continue under pressure while injecting this product by means of a hypochlorinator. Stop water flow when a chlorine residual test of 50 ppm is obtained at the low pressure end of the new main section after a 24 hour retention time. When chlorination is completed, the system must be flushed free of all heavily chlorinated water.

NEW TANKS, BASINS, ETC. - Remove all physical soil from surfaces. Place 43 oz. of this product for each 5 cubic feet of working capacity (500 ppm available chlorine). Fill to working capacity and allow to stand for at least 4 hours. Drain and flush with potable water and return to surface.

NEW FILTER SAND - Apply 170 oz. of this product for each 150 to 200 cubic feet of sand. The action of the product dissolving as the water passes through the bed will aid in sanitizing the new sand.

NEW WELLS - Flush the casing with a 50 ppm available chlorine solution of water containing 11 oz. of this product for each 100 gallons of water. The solution should be pumped or fed by gravity into the well after thorough mixing with agitation. The well should stand for several hours or overnight under chlorination. It may then be pumped until a representative raw water sample is obtained. Bacterial examination of the water will indicate whether further treatment is necessary.

EXISTING EQUIPMENT - Remove equipment from service, thoroughly clean surfaces of all physical soil. Sanitize by placing 45 oz. of this product for each 5 cubic feet capacity (approximately 500 ppm available chlorine). Fill to working capacity and let stand at least 4 hours. Drain and place in service. If the previous treatment is not practical, surfaces may be sprayed with a solution containing 11 oz. of this product for each 5 gallons of water (approximately 1000 ppm available chlorine). After drying, flush with water and return to service.

EMERGENCY DISINFECTION AFTER FLOODS

WELLS - Thoroughly flush contaminated casing with a 500 ppm available chlorine solution. Prepare this solution by mixing 11 oz. of this product with 10 gallons of water. Backwash the well to increase yield and reduce turbidity, adding sufficient chlorinating solution to the backwash to produce a 10 ppm available chlorine residual, as determined by a chlorine test kit. After the turbidity has been reduced and the casing has been treated, add sufficient chlorinating solution to produce a 50 ppm available chlorine residual. Agitate the well water for several hours and take a representative water sample. Retreat well if water samples are biologically unacceptable.

RESERVOIRS - In case of contamination by overflowing streams, establish hypochlorinating stations upstream of the reservoir. Chlorinate the inlet water until the entire reservoir obtains a 0.2 ppm available chlorine residual, as determined by a suitable chlorine test kit. In case of contamination from surface drainage, apply sufficient product directly to the reservoir to obtain a 0.2 ppm available chlorine residual in all parts of the reservoir.

BASINS, TANKS, FLUMES, ETC. - Thoroughly clean all equipment, then apply 43 oz. of product per 5 cu. ft. of water to obtain 500 ppm available chlorine, as determined by a suitable test kit. After 24 hours, drain, flush, and return to service. If the previous method is not suitable, spray or flush the equipment with a solution containing 11 oz. of this product for each 5 gallons of water (1000 ppm available chlorine). Allow to stand for 2 to 4 hours, flush and return to service.

FILTERS - When the sand filter needs replacement, apply 170 oz. of this product for each 150 to 200 cubic feet of sand. When the filter is severely contaminated, additional product should be distributed over the surface at the rate of 100 oz. per 20 sq. ft. Water should stand at a depth of 1 foot above the surface of the filter bed for 4 to 24 hours. When filter beds can be backwashed of mud and silt, apply 170 oz. of this product per each 50 sq. ft., allowing the water to stand at a depth of 1 foot above the filter sand. After 30 minutes, drain water to the level of the filter. After 4 to 6 hours, drain, and proceed with normal backwashing.

DISTRIBUTION SYSTEM - Flush repaired or replaced section with water. Establish a hypochlorinating station and apply sufficient product until a consistent available chlorine residual of at least 10 ppm remains after 24 hour retention time. Use a chlorine test kit.

EMERGENCY DISINFECTION AFTER FIRES

CROSS CONNECTIONS OF EMERGENCY CONNECTIONS - Hypochlorination or gravity feed equipment should be set up near the intake of the untreated water supply. Apply sufficient product to give a chlorine residual of at least 0.1 to 0.2 ppm at the point where the untreated supply enters the regular distribution system. Use a chlorine test kit.

EMERGENCY DISINFECTION AFTER DROUGHTS

SUPPLEMENTARY WATER SUPPLIES - Gravity or mechanical hypochlorite feeders should be set up on a supplementary line to dose the water to a minimum chlorine residual of 0.2 ppm after a 20 minute contact time. Use a chlorine test kit.

WATER SHIPPED IN BY TANKS, TANK CARS, TRUCKS, ETC. - Thoroughly clean all containers and equipment. Spray a 500 ppm available chlorine solution and rinse with potable water after 5 minutes. This solution is made by mixing 11 oz. of this product for each 10 gallons of water. During the filling of the containers, dose with sufficient amounts of this product to provide at least a 0.2 ppm chlorine residual. Use a chlorine test kit.

EMERGENCY DISINFECTION AFTER MAIN BREAKS

MAINS - Before assembly of the repaired section, flush out mud and soil. Permit a water flow of at least 2.5 feet per minute to continue under pressure while injecting this product by means of a hypochlorinator. Stop water flow when a chlorine residual of test of 50 ppm is obtained at the low pressure end of the new main section after a 24 hour retention time. When chlorination is completed, the system must be flushed free of all heavily chlorinated water.

COOLING TOWER/EVAPORATIVE CONDENSER WATER

SLUG FEED METHOD - Initial Dose: When system is noticeably fouled, apply 107 to 213 oz. of this product per 10,000 gallons of water in the system to obtain from 5 to 10 ppm available chlorine. Repeat until control is achieved.

Subsequent Dose: When microbial control is evident, add 21 oz. of this product per 10,000 gallons of water in the system daily, or as needed to maintain control and keep the chlorine residual at 1 ppm. Badly fouled systems must be cleaned before treatment is begun.

INTERMITTENT FEED METHOD - Initial Dose: When system is noticeably fouled, apply 107 to 213 oz. of this product per 10,000 gallons of water in the system to obtain 5 to 10 ppm available chlorine. Apply half (or 1/3, 1/4, or 1/5) of this initial dose when half (or 1/3, 1/4, or 1/5) of the water in the system has been lost by blowdown.

Subsequent Dose: When microbial control is evident, add 21 oz. of this product per 10,000 gallons of water in the system to obtain a 1 ppm residual. Apply half (or 1/3, 1/4, or 1/5) of this initial dose when half (or 1/3, 1/4, or 1/5) of the water in the system has been lost by blowdown. Badly fouled systems must be cleaned before treatment is begun.

CONTINUOUS FEED METHOD - Initial Dose: When system is noticeably fouled, apply 107 to 213 oz. of this product per 10,000 gallons of water in the system to obtain 5 to 10 ppm available chlorine.

Subsequent Dose: Maintain this treatment level by starting a continuous feed of 2 oz. of this product per 1,000 gallons of water lost by blowdown to maintain a 1 ppm residual. Badly fouled systems must be cleaned before treatment is begun.

LAUNDRY SANITIZERS

Household Laundry Sanitizers

IN SOAKING SUDS - Thoroughly mix 4 oz. of this product to 10 gallons of wash water to provide 200 ppm available chlorine. Wait 5 minutes, then add soap or detergent. Immerse laundry for at least 11 minutes prior to starting the wash/rinse cycle.

IN WASHING SUDS - Thoroughly mix 4 oz. of this product to 10 gallons of wash water containing clothes to provide 200 ppm available chlorine. Wait 5 minutes, then add soap or detergent and start the wash/rinse cycle.

Commercial Laundry Sanitizers

Wet fabrics or clothes should be spun dry prior to sanitization. Thoroughly mix 4 oz. of this product with 10 gallons of water to yield 200 ppm available chlorine. Promptly after mixing the sanitizer, add the solution into the prewash prior to washing fabrics/clothes in the regular wash cycle with a good detergent. Test the level of available chlorine, if solution has been allowed to stand. Add more of this product if the available chlorine level has dropped below 200 ppm.

FARM PREMISES

Remove all animals, poultry, and feed from premises, vehicles, and enclosures. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities occupied or traversed by animals or poultry. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. To disinfect, saturate all surfaces with a solution of at least 1000 ppm available chlorine for a period of 10 minutes. A 1000 ppm solution can be made by thoroughly mixing 21 oz. of this product with 10 gallons of water. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals or poultry, as well as the cleaned forks, shovels, and scrapers used for removing litter and manure. Ventilate buildings, cars, boats, and other closed spaces. Do not house livestock or poultry or employ equipment until chlorine has been dissipated. All treated feed racks, mangers, troughs, automatic feeders, fountains, and waterers must be rinsed with potable water before reuse.

PULP AND PAPER MILL PROCESS WATER SYSTEMS

SLUG FEED METHOD - Initial Dose: When system is noticeably fouled, apply 107 to 213 oz. of this product per 10,000 gallons of water in the system to obtain from 5 to 10 ppm available chlorine. Repeat until control is achieved.

Subsequent Dose: When microbial control is evident, add 21 oz. of this product per 10,000 gallons of water in the system daily, or as needed to maintain control and keep the chlorine residual at 1 ppm. Badly fouled systems must be cleaned before treatment is begun.

INTERMITTENT FEED METHOD - Initial Dose: When system is noticeably fouled, apply 107 to 213 oz. of this product per 10,000 gallons of water in the system to obtain 5 to 10 ppm available chlorine. Apply half (or 1/3, 1/4, or 1/5) of this initial dose when half (or 1/3, 1/4, or 1/5) of the water in the system has been lost by blowdown.

Subsequent Dose: When microbial control is evident, add 21 oz. of this product per 10,000 gallons of water in the system to obtain a 1 ppm residual. Apply half (or 1/3, 1/4, or 1/5) of this initial dose when half (or 1/3, 1/4, or 1/5) of the water in the system has been lost by blowdown. Badly fouled systems must be cleaned before treatment is begun.

CONTINUOUS FEED METHOD - Initial Dose: When system is noticeably fouled, apply 107 to 213 oz. of this product per 10,000 gallons of water in the system to obtain 5 to 10 ppm available chlorine.

Subsequent Dose: Maintain this treatment level by starting a continuous feed of 2 oz. of this product per 1,000 gallons of water lost by blowdown to maintain a 1 ppm residual. Badly fouled systems must be cleaned before treatment is begun.

AGRICULTURAL USES

POST-HARVEST PROTECTION - Potatoes can be sanitized after cleaning and prior to storage by spraying with a sanitizing solution at a level of 1 gallon of sanitizing solution per ton of potatoes. Thoroughly mix 2 oz. of this product to 2 gallons of water to obtain 500 ppm available chlorine.

Disinfect leaf-cutting bee cells and bee boards by immersion in a solution containing 1 ppm available chlorine for 3 minutes. Allow cells to drain for 2 minutes and dry for 4 to 5 hours or until no chlorine odor can be detected. This solution is made by thoroughly mixing 2 tsp. of this product to 100 gallons of water. The bee domicile is disinfected by spraying with a 0.1 ppm solution until all surfaces are thoroughly wet. Allow the domicile to dry until all chlorine odor has dissipated. [Not for use in California]

FOOD EGG SANITIZATION - Thoroughly clean all eggs. Thoroughly mix 4 oz. of this product with 10 gallons of warm water to produce a 200 ppm available chlorine solution. The sanitizer temperature should not exceed 130 degrees F. Spray the warm sanitizer so that the eggs are thoroughly wetted. Allow the eggs to thoroughly dry before casing or breaking. Do not apply a potable water rinse. The solution should not be re-used to sanitize eggs.

FRUIT & VEGETABLE WASHING - Thoroughly clean all fruits and vegetables in a wash tank. Thoroughly mix 11 oz. of this product in 200 gallons of water to make a sanitizing solution of 25 ppm available chlorine. After draining the tank, submerge fruit or vegetables for 2 minutes in a second wash tank containing the recirculating sanitizing solution. Spray rinse vegetables with the sanitizing solution prior to packaging. Rinse fruit with potable water only prior to packaging.

AQUACULTURAL USES

FISH PONDS - Remove fish from ponds prior to treatment. Thoroughly mix 213 oz. of this product to 10,000 gallons of water to obtain 10 ppm available chlorine. Add more product to the water if the available chlorine level is below 1 ppm after 5 minutes. Return fish to pond after the available chlorine level reaches zero.

FISH POND EQUIPMENT - Thoroughly clean all equipment prior to treatment. Thoroughly mix 4 oz. of this product to 10 gallons of water to obtain 200 ppm available chlorine. Porous equipment should soak for one hour.

MAINE LOBSTER PONDS - Remove lobsters, seaweed, etc. from ponds prior to treatment. Drain the pond. Thoroughly mix 12,800 oz. of this product to 10,000 gallons of water to obtain at least 600 ppm available chlorine. Apply so that all barrows, gates, rocks and dams are treated with product. Permit high tide to fill the pond and then close the gates. Allow water to stand for 2 to 3 days until the available chlorine level reaches zero. Open and allow 2 tidal cycles to flush the pond before returning lobsters to the pond. [Not for use in California]

CONDITIONING LIVE OYSTERS - Thoroughly mix 11 oz. of this product to 10,000 gallons of water at 50 to 70 degrees F to obtain 0.5 ppm available chlorine. Expose oysters to this solution for at least 15 minutes, monitoring the available chlorine level so that it does not fall below 0.05 ppm. Repeat entire process if the available chlorine level drops below 0.05 ppm or the temperature falls below 50 degrees F. [Not for use in California]

CONTROL OF SCAVENGERS IN FISH HATCHERY PONDS - Prepare a solution containing 200 ppm of available chlorine by mixing 4 oz. of product with 10 gallons of water. Pour into drained pond potholes. Repeat if necessary. Do not put desirable fish back into refilled ponds until chlorine residual has dropped to 0 ppm, as determined by a test kit.

SANITIZATION OF DIALYSIS MACHINES

Flush equipment thoroughly with water prior to using this product. Thoroughly mix 13 oz. of this product to 10 gallons of water to obtain at least 600 ppm available chlorine. Immediately use this product in the hemodialysate system allowing for a minimum contact time of 15 minutes at 20 degrees C. Drain system of the sanitizing solution and thoroughly rinse with water. Discard and DO NOT reuse the spent sanitizer. Rinsate must be monitored with a suitable test kit to ensure that no available chlorine remains in the system.

This product is recommended for decontaminating single and multi patient hemodialysate systems. This product has been shown to be an effective disinfectant (virucide, fungicide, bactericide, pseudomonicide) when tested by AOAC and EPA test methods. This product may not totally eliminate all vegetative microorganisms in hemodialysate delivery systems due to their construction and/or assembly, but can be relied upon to reduce the number of microorganisms to acceptable levels when used as directed. This product should be used in a disinfectant program which includes bacteriological monitoring of the hemodialysate delivery system. This product is NOT recommended for use in hemodialysate or reverse osmosis (RO) membranes.

Consult the guidelines for hemodialysate systems which are available from the Hepatitis Laboratories, CDC, Phoenix, AZ 85021.

ASPHALT OR SEALED WOOD ROOFS AND SIDINGS

To control fungus and mildew, first remove all physical soil by brushing and hosing with clean water, and apply a 5000 ppm available chlorine solution. Mix 11 oz. of this product per gallon of water and brush or spray roof or siding. After 30 minutes, rinse by hosing with clean water. *[Not for use in California]*

BOAT BOTTOMS

To control slime on boat bottoms, sling a plastic tarp under boat, retaining enough water to cover the fouled bottom area, but not allowing water to enter enclosed area. This envelope should contain approximately 500 gallons of water for a 14 foot boat. Add 37 oz. of this product to this water to obtain a 35 ppm available chlorine concentration. Leave immersed for 8 to 12 hours. Repeat if necessary. Do not discharge the solution until the free chlorine level has dropped to 0 ppm, as determined by a swimming pool test kit. *[Not for use in California]*

ARTIFICIAL SAND BEACHES

To sanitize the sand, spray a 500 ppm available chlorine solution containing 11 oz. of this product per 10 gal. of water at frequent intervals. Small areas can be sprinkled with a watering can. *[Not for use in California]*

WAREWASHING

FOR SANITIZING TABLEWARE IN LOW TEMPERATURE DISHWASHING MACHINE -Dispense this product into final rinse water at 100 ppm available chlorine. Do not allow concentration to fall below 50 ppm. Air dry. Dispenser should be set to deliver 6.5 cc of sanitizing solution per gallon of water to give 100 ppm of available chlorine. Only a qualified service representative should set or adjust dispenser on the machine.

HOSPITAL USE

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 (AIDS VIRUS) ON SURFACES OR OBJECTS SOILED WITH BLOOD OR BODY FLUIDS

Kills HIV-1 on PRE-CLEANED ENVIRONMENTAL SURFACES/ OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS in health care settings (hospitals, nursing homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS).

Personal Protection: Wear disposable latex gloves, protective gown, face mask and eye covering, as appropriate, when handling HIV-1 infected blood or body fluids.

Cleaning Procedure: All blood and other body fluids must be thoroughly cleaned from surfaces and objects before applications of this disinfectant. Product can be used for this purpose.

Contact Time: Prepare a 2700 ppm available chlorine solution. Thoroughly wet surface with disinfectant spray. Allow to air dry. The efficacy of a 5 minute contact time has been shown to be adequate against HIV-1 (AIDS Virus).

Disposal of Infectious Materials: Blood and other body fluids should be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

SPECIAL LABEL INSTRUCTIONS FOR CLEANING PRIOR TO DISINFECTION AGAINST CLOSTRIDIUM DIFFICILE ENDOSPORES

Personal Protection: Wear appropriate barrier protection such as gloves, gowns, masks or eye covering.

Cleaning Procedure: Fecal matter and waste must be thoroughly cleaned from surfaces and objects before disinfection by application with clean cloth, mop and/or sponge saturated with product intended for disinfection. Cleaning should include vigorous wiping and/or scrubbing until visible soil is removed. Special attention is needed for high-touch surfaces. Surfaces in patient rooms should be cleaned in an appropriate manner, with restrooms and other 'dirty' areas cleaned last. Do not reuse soiled cloths.

Infectious Materials Disposal: Cleaning materials used that may contain feces/wastes should be disposed of immediately in accordance with local regulations for infectious materials disposal.

For Killing Clostridium difficile spores: Use 1 part bleach to 5 parts water to achieve a 1:5 dilution (10,000 ppm available chlorine) before use. Clean hard, non-porous surfaces by removing gross filth. Apply 1:5 solution and let stand for 10 minutes. Rinse and air dry. Do not use on non-stainless steel, aluminum, silver or chipped baked enamel.

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST MYCOBACTERIUM TUBERCULOSIS (TB) OF SURFACES/OBJECTS SOILED WITH BODY FLUIDS.

Kills Mycobacterium tuberculosis (TB) on hard non-porous surfaces soiled with body fluids in healthcare settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with body fluids, and in which the surfaces/objects likely to be soiled with body fluids can be associated with the potential for transmission of Mycobacterium tuberculosis.

Personal Protection: Disposable latex or vinyl gloves, gowns, masks, and/or eye coverings as appropriate must be worn during all cleaning and decontamination procedures of body fluids.

Cleaning Procedures: Remove gross filth or heavy soil from surfaces or objects prior to application of product.

Disinfectant Use and Contact Time: Effective against Mycobacterium tuberculosis on hard non-porous surfaces in the presence of a moderate amount of organic soil (e.g. sputum). Prepare disinfectant by mixing 22 oz. of this product per gallon of water to provide 10,000 ppm of available chlorine. Leave surfaces wet for 10 minutes. Drain and let air dry.

Disposal of Infectious Materials: Blood and other body fluids should be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

[Include the following only if critical or semi-critical devices included on label]

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

[Optional H1N1 Language]

Respiratory illnesses attributable to Pandemic 2009 H1N1 are caused by Influenza A virus. This product (Product Name) is a broad-spectrum hard surface disinfectant that has been shown to be effective against (Influenza A virus tested and listed on the label) and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 (formerly called swine flu).

This product has demonstrated effectiveness against Influenza A virus and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 Influenza A virus.

This product has demonstrated effectiveness against (Influenza A virus tested and listed on the label) and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 (formerly called swine flu).

Kills Pandemic 2009 H1N1 Influenza A virus (formerly called swine flu).

Kills Pandemic 2009 H1N1 Influenza A virus.

[Optional Logos]



Solid/Spot
PMS 361 C

Process Color
C=80% Y=100% M=0% K=0%

[Optional Table of Proportioning]

PPM	Quantity of Bleach	Quantity of Water
100	0.25 FL OZ	1 GAL
	0.50 FL OZ	2 GAL
	1.25 FL OZ	5 GAL
	2.00 FL OZ	10 GAL
200	0.50 FL OZ	1 GAL
	1.00 FL OZ	2 GAL
	2.50 FL OZ	5 GAL
	4.00 FL OZ	10 GAL
400	1.00 FL OZ	1 GAL
	2.00 FL OZ	2 GAL
	5.00 FL OZ	5 GAL
	8.00 FL OZ	10 GAL
500	1.25 FL OZ	1 GAL
	2.50 FL OZ	2 GAL
	6.25 FL OZ	5 GAL
	10.00 FL OZ	10 GAL
600	1.50 FL OZ	1 GAL
	3.00 FL OZ	2 GAL
	7.50 FL OZ	5 GAL
	12.00 FL OZ	10 GAL

PPM	Quantity of Bleach	Quantity of Water
1,000	2.50 FL OZ	1 GAL
	5.00 FL OZ	2 GAL
	12.50 FL OZ	5 GAL
	20.00 FL OZ	10 GAL
2,700	6.75 FL OZ	1 GAL
	13.50 FL OZ	2 GAL
	33.75 FL OZ	5 GAL
	54.00 FL OZ	10 GAL
5,000	12.50 FL OZ	1 GAL
	25.00 FL OZ	2 GAL
	62.50 FL OZ	5 GAL
	100.00 FL OZ	10 GAL
10,000	25.00 FL OZ	1 GAL
	50.00 FL OZ	2 GAL
	125.00 FL OZ	5 GAL
	200.00 FL OZ	10 GAL
1 teaspoon (tsp.) = 0.16 FL OZ 1 tablespoon (Tbsp.) = 0.50 FL OZ 8 tablespoons = 4 FL OZ (1/2 cup) PPM = parts per million		