

**XY-12  
LIQUID SANITIZER**

**ACTIVE INGREDIENT:**

Sodium hypochlorite ..... 8.4%

**INERT INGREDIENTS:** ..... 91.6%**KEEP OUT OF REACH OF CHILDREN****DANGER****PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**DANGER:** Corrosive. May cause severe skin and eye irritation or chemical burns to broken skin. Causes eye damage. Wear rubber gloves, chemical goggles and protective clothing. Wash after handling. Avoid breathing vapors. Vacate poorly ventilated areas as soon as possible. Do not return until strong odors have dissipated.

**STATEMENT OF PRACTICAL TREATMENT**

**IF IN EYES:** Flush immediately with cool water. Remove contact lenses. Continue flushing for 15 minutes, holding eyelids apart. Get prompt medical attention.

**IF ON SKIN:** Wash with plenty of soap and water.

**IF SWALLOWED:** Rinse mouth at once; then drink one or two glasses of water or milk. DO NOT give vinegar or other acids. DO NOT induce vomiting. Never give anything by mouth to an unconscious person.

**CALL A POISON CONTROL CENTER OR PHYSICIAN IMMEDIATELY**

**FOR EMERGENCY MEDICAL INFORMATION, CALL TOLL-FREE 1-800-328-0026**

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Do not discharge into lakes, streams, ponds or public waterways unless in accordance with a NPDES permit. For guidance, contact the regional office of the U.S. Environmental Protection Agency.

**PHYSICAL AND CHEMICAL HAZARDS:**

**STRONG OXIDIZING AGENT:** Mix only with water according to label directions. Mixing this product with chemicals (e.g. ammonia, acids, detergents, etc.) or organic matter (e.g. urine, feces, etc.) will release chlorine gas which is irritating to eyes, lungs and mucous membranes.

**FOR INDUSTRIAL USE ONLY****DO NOT MIX WITH ANYTHING BUT WATER**

Net Contents:	1 gallon (3.78 l)	30 gallons (113.6 l)
	5 gallons (18.9 l)	55 gallons (208.2 l)
	15 gallons (56.8 l)	

EPA Reg. No. 1677-52


EPA Est. 1677-IL-2 (J), 1677-NJ-1 (W), 1677-TX-1 (D), 1677-GA-1 (M), 1677-MN-1 (F),  
1677-CA-1 (S), 1677-PR-1 (B), 6574-CA-1 (L)

The superscript refers to the first letter of the date code.


Ecolab Inc.

Ecolab Center

St. Paul, MN 55102



INSTITUTIONS - SCHOOLS - RESTAURANTS - FOOD SERVICES -  
DAIRIES - FOOD PROCESSING AND BEVERAGE PLANTS



# **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in manner inconsistent with its labeling.

**NOTE:** This product degrades with age. Use a chlorine test kit and increase dosage, as necessary to obtain the required level of available chlorine.

## **SANITIZING NONPOROUS FOOD CONTACT EQUIPMENT - RINSE METHOD**

A solution of 100 ppm available chlorine may be used in the sanitizing solution if a chlorine test kit is available. Solutions containing an initial concentration of 100 ppm av. Cl must be tested and adjusted periodically to insure that the av. Cl does not drop below 50 ppm. Prepare a 100 ppm sanitizing solution by thoroughly mixing 2 oz of XY-12 with 13 gl of water. If no test kit is available, prepare a sanitizing solution by mixing 4 oz of XY-12 with 13 gallons of water to provide approximately 200 ppm av. Cl by weight.

Clean all surfaces with proper detergent and rinse with water. Just prior to use, rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 2 minutes. If solution contains less than 50 ppm av. Cl as determined by a suitable test kit, either discard the solution or add sufficient product to reestablish a 200 ppm residual. Allow equipment to drain thoroughly. Do not rinse and do not soak overnight.

## **SANITIZING POROUS FOOD CONTACT EQUIPMENT - RINSE METHOD**

Prepare a 600 ppm solution by thoroughly mixing 3 oz of this product in 3 gl water. Clean surfaces in the normal manner. Rinse all surfaces thoroughly with the 600 ppm solution, maintaining contact for at least 2 minutes. Prior to using equipment, rinse all surfaces with a 200 ppm av. Cl solution (4 oz/13 gl). Do not rinse and do not soak overnight.

## **SANITIZING EATING AND DRINKING UTENSILS - IMMERSION METHOD**

1. Scrape and preflush utensils to remove excess soil.
2. Wash with good detergent or compatible cleaner (see your Ecolab representative for a recommendation).
3. Rinse with clear water
4. Sanitize in a solution of 100 ppm av. Cl if a chlorine test kit is available. Test and adjust solution periodically to insure that the av. Cl does not drop below 50 ppm. If no test kit is available, prepare sanitizing solution to provide 200 ppm av. Cl. Immerse all utensils for at least one minute. Use 2 minutes exposure time if required by governing sanitary code.
5. Drain and air dry.

**NOTE:** FOR MECHANICAL OPERATIONS prepared use solutions may not be reused for sanitizing but may be reused for other purposes such as cleaning.

FOR MANUAL OPERATION fresh sanitizing solutions should be prepared as soon as they become diluted or soiled.

## **TABLEWARE SANITIZER AND DESTAINER FOR MECHANICAL SPRAY WAREWASHING MACHINES**

For sanitizing tableware in low-temperature warewashing machines, inject XY-12 into the final rinse water at a concentration of 100 ppm available chlorine. Do not exceed 200 ppm. Air dry.

To insure that available chlorine concentration does not fall below 50 ppm, periodically test the rinse solution with a suitable test kit and adjust the dispensing rate accordingly. Consult your local Ecolab Specialist for technical assistance and further information on sanitizing tableware in warewashing machines.

FORMULATED FOR USE WITH ECOLAB AUTOMATIC DISPENSING SYSTEMS.

DO NOT USE ON SILVER AND SILVER PLATE.

For brand name products sold for dispensing through computerized systems the following directions for use will appear on the label: For use only with the (Brand Name) Allocation System.

## **WATER CHLORINATION**

For farm, private or small municipal water chlorination, use 0.1 to 0.5 ppm, available chlorine. Use chlorine test kit for the determination of proper concentrations and amount of residual chlorine.

**BEST AVAILABLE COPY**

**FOR CONTINUOUS TREATMENT OF MEAT AND POULTRY OR FRUIT AND VEGETABLE CONVEYORS -**

wash, rinse and sanitize conveyance equipment. During processing, apply XY-12 at a 200 ppm available chlorine level to conveyors with MIKRO MASTER or other suitable feeding equipment. Controlled volumes of sanitizer are applied to return portion of conveyor through nozzles so located as to permit maximum drainage of sanitizer from equipment and to prevent puddles on top of belt. During interruptions in operation, spray equipment, peelers, collators, slicers and saws with MIKRO MASTER dispensed XY-12 solution of 200 ppm av. Cl. Conveyor equipment should be free of product when applying this spray.

**BACTERIOPHAGE CONTROL**

XY-12 will significantly reduce the incidence of Streptococcus cremoris and Streptococcus diacetilactis bacteriophage in cheese manufacturing establishments by fogging at a concentration of 600 ppm available chlorine. Fogging should be used as a supplement to acceptable manual cleaning and sanitizing of room surfaces as described above.

**DIRECTIONS FOR FOGGING**

Prior to fogging, food products and packaging material must be removed from the room or carefully protected. Fog desired areas using one quart per 1000 cu. ft. of room area with an XY-12 solution containing 600 ppm available chlorine. Vacate the area of all personnel for a minimum of 2 hours after fogging. All food contact surfaces must then be thoroughly rinsed with an XY-12 solution at 200 ppm available chlorine. All surfaces to drain thoroughly before operation are resumed.

**LAUNDRY SANITIZATION****HOUSEHOLD LAUNDRY SANITIZING**

In Soaking Suds - Thoroughly mix 4 oz. XY-12 to 13 gallons of wash water to provide 200 ppm available chlorine. Wait 5 minutes, then add soap or detergent. Immerse laundry for at least 11 minutes prior to starting the wash/rinse cycle.

In Washing Suds - thoroughly mix 4 oz. XY-12 to 13 gallons of wash water containing clothes to provide 200 ppm available chlorine. Wait 5 minutes, then add soap or detergent and start the wash/rinse cycle.

**COMMERCIAL LAUNDRY SANITIZING**

Wet fabrics or clothes should be spun dry prior to sanitization. Thoroughly mix 4 oz. XY-12 to 13 gallons of water to yield 200 ppm available chlorine. Promptly add the solution into the prewash prior to washing fabrics/clothes in the regular wash cycle with a good detergent. Test the level of available chlorine, if solution has been allowed to stand. Add more XY-12 if the available chlorine level has dropped below 200 ppm.



**FOOD EGG SANITIZATION** - Thoroughly clean all eggs before proceeding. To sanitize clean shell eggs intended for food or food products, spray with a solution of 2 oz of product in 6.5 gallons of water (providing 200 ppm available chlorine). The solution must be equal to or warmer than the eggs, by not to exceed 130 deg F. Wet eggs thoroughly and allow to drain. Eggs that have been sanitized with this chlorine compound may be broken for use in the manufacture of egg products without a prior potable water rinse. Eggs must be thoroughly dry before casing or breaking. The solution must not be reused for sanitizing eggs.

**LAUNDRY BLEACH** - Add 2 oz of product per 10 gallons of wash water (125 ppm av. Cl).

**FRUIT AND VEGETABLE WASHING** - Thoroughly clean all fruits and vegetables in a wash tank. Thoroughly mix 8 oz of product in 200 gallons of water to make a sanitizing solution of 25 ppm available chlorine. After draining the tank, submerge fruits or vegetables for 2 minutes in a second wash tank containing the recirculating sanitizing solution. Spray rinse vegetables with the sanitizing solution prior to packaging. Rinse fruit with potable water only prior to packaging.

**AVAILABLE CHLORINE TABLE OF PROPORTIONS**

0.5 ppm - 1 oz in 1300 gl water	
50 ppm - 1 oz in 13 gl water	
100 ppm - 2 oz in 13 gl water	1 oz in 6.5 gl water
200 ppm - 4 oz in 13 gl water	2 oz in 6.5 gl water
600 ppm - 12 oz in 13 gl water	



For other specialized cleaning and sanitizing operations, consult your local Ecolab representative.

For service or additional information, call 1-800-35-CLEAN (352-5326).

**STORAGE & DISPOSAL**

Store this product in a cool, dry area, away from direct sunlight and heat to avoid deterioration. In case of spill, flood the area with large quantities of water.

**PESTICIDE DISPOSAL:** Product or rinsates that cannot be used should be diluted with water before disposal in a sanitary sewer. Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

**CONTAINER DISPOSAL:** (1 gal) Do not reuse empty containers. Wrap container and put in trash. (5, 55 gal plastic) Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

