

**DAIRY AND FOOD
PROCESSING EQUIPMENT**

Rinse equipment with clean lukewarm water immediately after use. Flush thoroughly to remove all loose material.

Brush wash all surfaces with proper Klenzade detergent.

Rinse thoroughly with hot (145°F.) water.

Sanitize just before using equipment with K-SAN at 1 oz. to 2 gallons of water (100 ppm. of sodium dodecylbenzene sulfonate). All surfaces should be exposed to the sanitizer for a period of not less than two minutes.

CONTAINS NOT MORE THAN
7.0% PHOSPHORUS.
1.15 GMS. PER GAL. OF AVERAGE
RECOMMENDED USE CONCENTRATION.

Rinse empty container thoroughly with water and discard.

Manufactured by
ECONOMICS LABORATORY, INC.
KLENZADE DIVISION
Coborn Bldg. - St. Paul, Minnesota 55102

EPA Est. 1677-IL-2

50301-0375

PRINTED IN USA

**KLENZADE®
ACID
SANITIZER**

ACCEPTED

JAN. 21 1976

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 1677-56



formula S-15
EPA Reg. No. 1677-56AA

ACTIVE INGREDIENTS:	
Phosphoric Acid	22.00%
Sodium Dodecylbenzene sulfonate	2.75%
INERT INGREDIENTS:	
(Contains a non-ionic detergent)	75.25%
TOTAL	100.00%

DANGER

**KEEP OUT OF REACH OF CHILDREN.
SEE FIRST AID AND OTHER PRECAUTIONS ON SIDE PANEL**

NINE CONTENTS 5.31 U.S. GALLONS

42622

1677-56

INSTITUTIONAL APPLICATION:

**DISHES, GLASSES, UTENSILS
AND EQUIPMENT**

1. Scrape and prewash dishes, glasses, utensils and equipment whenever possible.
2. Wash with appropriate Klenzade detergent.
3. Rinse with clear water.
4. Sanitize in a solution of 1 oz. K-SAN to 2 gallons of water (100 ppm sodium dodecylbenzene sulfonate). Immerse all utensils for at least two minutes, or for contact time specified by governing sanitary code.
5. Place sanitized utensils on rack or drainboard to air dry.

DANGER

Keep out of reach of children. Corrosive. Cause eye damage and skin irritation. Do not get in eyes on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal when swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol.

CALL A PHYSICIAN IMMEDIATELY.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respirator depression and convulsion may be needed.

01/21/1976

1/1