

STER-BAC UDDER WASH COMPOUND

- Sanitizes udder before milking
- Sanitizes paper towel
- Helps prevent spread of udder infection.

DIRECTIONS FOR USE

1. Use one ounce Ster-Bac Udder Wash Compound to 4 gallons of warm water for a 200 ppm solution.
2. For each cow, place one clean towel in bucket containing 200 ppm of fresh, warm Ster-Bac Udder Wash Compound solution
3. Wash udder and discard towel in second bucket
4. Wipe off excess solution with clean, dry paper towel
5. Examine each quarter with strip cup.
6. After milking, dip each teat in dip cup containing 200 ppm Ster-Bac Udder Wash Compound solution.

8778/0301/0571 1-Ph in USA



Ster-Bac

for

FORMULA KQ-11

USDA Reg. No. 292-64

UDDER WASHING

A Quaternary Ammonium Compound

ACTIVE INGREDIENTS:	
n-Alkyl (50% C ₁₂ , 40% C ₁₄ , 10% C ₁₆) dimethyl benzyl ammonium chloride	10%
INERT INGREDIENTS	
TOTAL	100%

Contents: One U. S. Gallon

LOCAL & STATE REGULATIONS

Where local or state regulations are in effect concerning quaternary ammonium compounds for udder sanitizing, consult them for recommended dilutions and procedures.

Ster-Bac Udder Wash Compound fulfills the criteria of Grade 'A' Pasteurized Milk Ordinance 1965 Recommendations of U.S. Public Health Service in waters up to 550 ppm of hardness calculated as CaCO₃, when tested by the method outlined by Chambers.

CAUTION

Undiluted Ster-Bac may cause skin irritation or damage to the eyes. Harmful if swallowed. Avoid contact with skin and eyes. Do not take internally. For eyes, flush immediately with plenty of water and get immediate medical attention. Avoid contamination of food.

GUARANTEED PHENOL COEFFICIENT

AT 20° C. ORGANISM

Staphylococcus aureus	.75
Salmonella typhosa	.50

Manufactured Exclusively by
KLENZADE PRODUCTS

Division of Economics Laboratory, Inc.
Osborn Bldg. — St. Paul, Minnesota 55102

1671
Reg