

1677-158

1/8/2008

lof8

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

JAN 28 2008



Office of Pesticide Programs

Ecolab, Inc.
380 N. Wabasha Street
St. Paul, MN 55102

Attention: Joy A. Salverda
Regulatory Analyst

Subject: Vortexx
EPA Registration No. 1677-158
Your Amendment Dated August 17, 2007

The amendment, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, to revise the product labeling based on the two new toxicity studies (Acute Dermal, MRID # 472134-01 and Acute Inhalation, MRID #472134-02) , is acceptable, provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3©(5) and sec. 4 when the Agency requires all registrants of similar products to submit such data.
2. Please note that the Registrant must include all available ATCC organism reference numbers on their label as a point of reference.
3. Refer to the enclosed copy of the efficacy review under "FINDINGS" and "RECOMMENDATIONS" for further comments.

Submit two (2) copies of final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Page 2
EPA Registration No. 1677-158

A stamped copy of the "accepted" product labeling is enclosed for your records.

If you have any questions concerning this letter, please contact Martha Terry at (703) 308-6217.

Sincerely



Marshall Swindell
Product Manager 33
Regulatory Management Branch 1
Antimicrobials Division (7510P)

Enclosure

VORTEXX

ACID LIQUID SANITIZER FOR FOOD PROCESSING EQUIPMENT
in Dairies, Dairy Farms, Breweries, Wineries, Beverage and Food Processing Plants

DISINFECTANT
Veterinary Clinics, Animal Life Science Laboratories, Industrial Facilities, Office Buildings, Recreational Facilities, Retail and Wholesale Establishments, Animal Care Facilities, Veterinary Facilities, Farms, Livestock Quarters, Poultry Premises, and Poultry Hatcheries

STERILANT FOR ASEPTIC MANUFACTURING AND PACKAGING EQUIPMENT

DISINFECTANT FOR THE PHARMACEUTICAL AND COSMETIC INDUSTRY

Active Ingredients:	
Hydrogen Peroxide.....	6.9%
Peroxyacetic Acid.....	4.4%
Octanoic Acid	3.3%
Inert Ingredients:	85.4%
Total:	100.0%

ACCEPTED
with **COMMENTS**
EPA Letter Dated:

JAN - 8 2008

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide, registered under EPA Reg. No. 1677-158

KEEP OUT OF REACH OF CHILDREN
DANGER

PRECAUTIONARY STATEMENTS - HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: CORROSIVE: Causes irreversible eye damage and skin burns. Do not get in eyes or on clothing. Harmful if inhaled or swallowed. Avoid breathing vapor or spray mist. Wear protective eyewear (goggles, face shield, or safety glasses) and protective gloves. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco. Remove and wash contaminated clothing before reuse.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 -20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

FOR EMERGENCY MEDICAL INFORMATION CALL TOLL FREE: 1-800-328-0026

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

PHYSICAL AND CHEMICAL HAZARDS: Strong oxidizing agent. Corrosive. Do not use in concentrated form. Mix only with water according to label instructions. Never bring concentrate in contact with other sanitizers, cleaners or organic substances.

ENVIRONMENTAL HAZARDS: (containers 5 gallons or greater) Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

4 of 8

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SANITIZATION

VORTEXX is recommended for use on pre-cleaned surfaces such as equipment, pipelines, tanks, vats, fillers, evaporators, pasteurizers and aseptic equipment in dairies, dairy farms, breweries, wineries, beverage and food processing plants. This product is effective as a sanitizer when solution is prepared in water of up to 500 ppm hardness as CaCO₃.

NOTE: FOR MECHANICAL OPERATIONS, prepared use solutions may not be reused for sanitizing but may be reused for other purposes such as cleaning.
FOR MANUAL OPERATIONS, fresh sanitizing solutions should be prepared at least daily or more often if the solution becomes diluted or soiled.

SANITIZING FOOD CONTACT SURFACES

Prior to sanitizing, remove gross food particles, then wash with a detergent solution, followed by a potable water rinse. Sanitize with a concentration of 1 to 2 oz. **VORTEXX** per 6 gallons of water (0.13 - 0.26% v/v concentration). At this dilution **VORTEXX** is effective against *Staphylococcus aureus*, *Escherichia coli*, *Escherichia coli* 0157:H7, *Listeria monocytogenes*, *Salmonella typhimurium* (milk isolate), *Pseudomonas aeruginosa*, *Enterobacter sakazakii* and *Vibrio cholerae* as well as beverage spoilage organism *Lactobacillus malefermentans*. Use immersion, coarse spray or circulation techniques as appropriate to the equipment. All surfaces should be exposed to the sanitizing solution for a period of not less than one minute unless a longer time is specified by the governing sanitary code. Drain thoroughly. Do not rinse.

SANITIZING EATING, DRINKING, AND FOOD PREP UTENSILS

1. Remove gross food particles by a prescrape, preflush and, when necessary, a presoak treatment.
2. Wash with detergent.
3. Rinse with clean water.
4. Sanitize in a solution of 1 to 2 oz. **VORTEXX** per 6 gallons of water (0.13 - 0.26% v/v concentration). Immerse all utensils for at least 1 minute or contact time specified by governing sanitary code.
5. Drain and air dry.

ELEVATED TEMPERATURE SANITIZING

At a temperature of 120° F. **VORTEXX** can be used as a food contact sanitizer at 1 oz. per 14 gallons water (0.055% v/v concentration). At this concentration **VORTEXX** is effective against *Staphylococcus aureus* and *Escherichia coli*. All surfaces should be exposed to the sanitizing solution for a period of not less than one minute, unless otherwise specified by governing sanitary code. Drain thoroughly. Do not rinse.

SANITIZING TABLEWARE

For sanitizing tableware in low-temperature (minimum 120° F) warewashing machines, inject **VORTEXX** into the final rinse water at a concentration of 0.055% v/v (1 oz. per 14 gallons of water) at 120° F. Allow 1 minute contact. Air dry.

To insure that the **VORTEXX** concentration does not fall below 0.055%, periodically test the rinse solution with a suitable test kit and adjust the dispensing rate accordingly. Consult your local Ecolab Specialist for technical assistance and further information on sanitizing tableware in warewashing machines.

FINAL SANITIZING BOTTLE RINSE

VORTEXX may be used as a final sanitizing rinse for returnable and non-returnable bottles (e.g. glass or PET) at a 0.13% dilution (1 oz. to 6 gallons). At this dilution **VORTEXX** is effective against *Staphylococcus aureus*, *Escherichia coli*, *Escherichia coli* 0157:H7, *Listeria monocytogenes*, *Salmonella typhimurium*, *Pseudomonas aeruginosa*, *Enterobacter sakazakii*, *Vibrio cholerae*, and *Lactobacillus malefermentans*. All surfaces should be exposed to the sanitizing solution for a period of not less than 1 minute. Drain thoroughly. No rinse necessary.

ACCEPTED
with COMMENTS
EPA Letter Dated:

JAN - 8 2009

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide, registered under EPA Reg. No.

1177-158

CONTINUOUS TREATMENT OF CONVEYORS

Wash, rinse and sanitize conveyor equipment. During processing, apply **VORTEXX** at no more than 0.13% - 0.26% v/v (1 to 2 oz. per 6 gallons of water) concentration to conveyor with Mikro Master or other suitable feeding equipment. At this dilution **VORTEXX** is effective against *Staphylococcus aureus*, *Escherichia coli*, *Escherichia coli* 0157:H7, *Listeria monocytogenes*, *Salmonella typhimurium* (milk isolate), *Pseudomonas aeruginosa*, *Enterobacter sakazakii* and *Vibrio cholerae*. Controlled volumes of **VORTEXX** are applied to return portion of conveyor through nozzles so located as to permit maximum drainage of **VORTEXX** from equipment and to prevent puddles on top of belt. During interruptions in operations, coarse spray the processing equipment with **VORTEXX** solution at not more than 0.26% v/v concentration. Conveyor equipment must be free of product when applying coarse spray. Conveyor surface should be exposed to the sanitizing solution for a period of not less than 60 seconds.

SANITIZING NON-FOOD CONTACT SURFACES

Preclean surfaces as directed above. Sanitize non-food contact surfaces such as floors, walls, tables, chairs, benches, drains, troughs, drip pans with 1 oz. **VORTEXX** per 8 gallons water (0.1% v/v concentration). At this concentration the product is effective against *Staphylococcus aureus*, *Enterobacter aerogenes*, *Escherichia coli* 0157:H7, *Listeria monocytogenes*, *Salmonella typhimurium* (milk isolate), *Pseudomonas aeruginosa* in 500 ppm hard water. The product is also effective against beverage spoilage organisms: *Aspergillus niger*, *Saccharomyces cerevisiae*, *Pediococcus damnosus* and *Lactobacillus malefermentans*. All surfaces should be exposed to the sanitizing solution for a period of not less than 5 minutes. Drain thoroughly and allow to air dry. No rinse necessary.

SANITIZING HARD, NON-POROUS OUTSIDE SURFACES OF AIRTIGHT, SEALED PACKAGES CONTAINING FOOD OR NON-FOOD PRODUCTS

VORTEXX may be used as a final sanitizing rinse for hard, non-porous outside surfaces of airtight, sealed packages containing food or non-food products at a 0.13% - 0.26% v/v dilution (1 to 2 oz. per 6 gallons of water). All surfaces should be exposed to the sanitizing solution for a period of not less than 5 minutes. Drain thoroughly. No rinse necessary.

FOAM SANITIZING NON-FOOD CONTACT SURFACES

VORTEXX is an effective foam sanitizer of precleaned non-food contact surfaces, such as boots, floors, walls, and drains, and equipment surfaces. For this application, prepare a solution of 0.13% v/v (1 oz. per 6 gallons water) **VORTEXX** and 0.13% v/v (1 oz. per 6 gallons water) **Liquid K**. For example, in 6 gallons of water, add 1 ounce of **VORTEXX** and 1 ounce of **Liquid K**. Apply solution as a foam using recommended equipment. Wet surfaces thoroughly. At this concentration, the product is effective against *Staphylococcus aureus*, *Enterobacter aerogenes*, and *Listeria monocytogenes*. Surfaces should be exposed to the sanitizing foam for a period of not less than 5 minutes. No rinse is necessary. Contact your Ecolab representative for information on Liquid K or EcoCare 600 FA foaming agents and a recommended foam generation device.

DIRECTIONS FOR FOGGING

To sanitize hard surfaces as an adjunct to acceptable manual cleaning and disinfecting of room surfaces: Prior to fogging, food products and packaging materials must be removed from the room or carefully protected. Fog desired areas using one quart of a 0.2% to 1.3 % **VORTEXX** solution (3 oz. to 14 oz. per 8 gallons of water) per 1000 cu. ft. of room volume. Vacate the area of all personnel during fogging and until the hydrogen peroxide air concentration is below 0.5 ppm. Allow surfaces to drain thoroughly before operations are resumed. Solutions above 0.5% may be corrosive and are not to be used on all surfaces. Test solution on surfaces prior to use. All hard non-porous food contact surfaces treated with the disinfectant and fog must be rinsed thoroughly with a potable water rinse.

SANITIZING NON-FOOD CONTACT PACKAGING EQUIPMENT

Prior to use of this product, remove gross soil particles from surfaces. Wash with a recommended detergent solution, rinse thoroughly with potable water. For sanitization against *Staphylococcus aureus*, *Enterobacter aerogenes*, *Escherichia coli* 0157:H7, *Listeria monocytogenes*, *Salmonella typhimurium* (milk isolate), *Pseudomonas aeruginosa* as well as beverage spoilage organisms, *Aspergillus niger*, *Pediococcus damnosus*, *Lactobacillus malefermentans* and *Saccharomyces cerevisiae*, apply 0.1 - 0.5% (1 to 5 oz. per 8 gallons of water) of **VORTEXX** to surfaces at a temperature of 25° to 45°C and allow to remain wet for at least 5 minutes. Allow surfaces to drain thoroughly before operations are resumed. Drainage may be followed by a potable or sterile water rinse.

ACCEPTED
with COMMENTS

EPA Letter Dated:

JAN - 8 2008

under the Federal Insecticide,
Fungicide, and Rodenticide Act
amended, for the pesticide,
registered under EPA Reg. No.

1677-158

SANITIZE PRECLEANED OR NEW RETURNABLE OR NON-RETURNABLE BOTTLED WATER CONTAINERS

To sanitize precleaned or new returnable or non-returnable containers for bottled water processing, apply **VORTEXX** at a concentration of 0.75 to 2.2% (9 to 26 oz. per 10 gallons of water) at a temperature of 40° to 60° C for at least 7 seconds. At these conditions, **VORTEXX** is effective against *Staphylococcus aureus*, *Escherichia coli*, *Salmonella typhi*, *Pediococcus damnosus*, *Lactobacillus malefermentans*, *Saccharomyces cerevisiae* and *Aspergillus niger*. After thorough draining, rinse interior container surfaces with a disinfected water rinse free of pathogenic bacteria.

ANTIMICROBIAL TREATMENT OF WATER FILTERS

To reduce the number of beverage spoilage organisms, *Pediococcus damnosus*, *Lactobacillus malefermentans*, *Saccharomyces cerevisiae* and *Aspergillus niger*, apply **VORTEXX** as a 0.75 to 2.2% (9 to 26 oz. per 10 gallons of water) solution at 25° to 45° C for a minimum contact time of 5 minutes. After thorough draining, rinse filters with a disinfected water rinse free of pathogenic bacteria.

ANTIMICROBIAL RINSE OF PRECLEANED OR NEW RETURNABLE OR NON-RETURNABLE CONTAINERS

To reduce the number of beverage spoilage organisms, *Pediococcus damnosus*, *Lactobacillus malefermentans*, *Saccharomyces cerevisiae* and *Aspergillus niger*, apply **VORTEXX** at a concentration of 0.75 to 2.2% (9 to 26 oz. per 10 gallons of water) at a temperature of 40° to 60° C for at least 7 seconds. After thorough draining, rinse interior container surfaces with a disinfected water rinse free of pathogenic bacteria.

ANTIMICROBIAL RINSE OF PRECLEANED OR NEW RETURNABLE OR NON-RETURNABLE CONTAINERS WITH THE ADDITION OF SURFACTANT

To reduce the number of beverage spoilage organisms, *Pediococcus damnosus*, *Lactobacillus malefermentans*, *Saccharomyces cerevisiae*, and *Aspergillus niger*, apply **VORTEXX** as follows. Add 6.7 oz. of surfactant product to 10 gallons of prepared **VORTEXX** solution. Prepare **VORTEXX** at a concentration of 0.75 to 2.2% (9 to 26 oz. per 10 gallons of water) solution. Use one of the following surfactants, KX-7014, KX-7015, KX-7016, or KX-7017. Apply at a temperature of 40° to 60° C with a contact time of at least 7 seconds. After thorough draining, rinse interior container surfaces with a disinfected water rinse free of pathogenic bacteria.

BACTERIOPHAGE CONTROL

When applied to pre-cleaned surfaces, **VORTEXX** will reduce the incidence of *Lactococcus lactis* subsp. *Lactis*, *Lactococcus lactis* subsp. *Cremonis*, and *Streptococcus thermophilus* bacteriophage in cheese manufacturing establishments by spraying or immersion of equipment at concentrations of 1-2 oz. per 6 gallons (0.13 - 0.26% v/v concentration) of water. All surfaces should be exposed to the solution for a period of not less than one minute. Drain thoroughly. Do not rinse.

ACCEPTED
with COMMENTS
EPA Letter Dated:
JAN - 8 2008
under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No.
1677-158

DISINFECTION

VORTEXX disinfects as it cleans in one operation. **VORTEXX** can be used to disinfect floors, walls and other hard, nonporous surfaces such as tables, chairs, countertops, bathroom fixtures, sinks, shelves, racks, carts, refrigerators, coolers, tile, linoleum, vinyl, glazed porcelain, plastic (such as polypropylene and polyethylene), stainless steel, or glass.

Areas of Use: Use **VORTEXX** in veterinary clinics, animal life science laboratories, industrial facilities, office buildings, recreational facilities, retail and wholesale establishments.

Combination Disinfection and Cleaning

VORTEXX is effective against *Staphylococcus aureus*, *Salmonella choleraesuis*, and *Pseudomonas aeruginosa* at 0.2% - 1.0% (1 oz. /4 gallons to 5 oz. /4 gallons) in hard water (500 ppm as CaCO₃), 5% blood serum and dried soap film residue on hard nonporous surfaces. For heavily soiled areas a precleaning step is required. Apply solution with mop, cloth, sponge, brush, scrubber, or coarse spray device or by soaking so as to wet all surfaces thoroughly. Allow to remain wet for 10 minutes, then remove solution and entrapped soil with a clean wet mop, cloth, or wet vacuum pickup. Prepare a fresh solution daily or when it becomes soiled or diluted. Rinse food contact surfaces that come in contact with food with potable water prior to reuse.

Virucidal

At 0.2% (1 oz./4 gals.) VORTEXX is effective against Influenza B, Influenza A(H3N2), Influenza A (H1N1), Influenza A (H10N7), Avian Influenza A (H3N2) and Reovirus on hard inanimate surfaces when used at 20° C with a 10 minute contact time in the presence of 500 ppm hard water and organic soil. Apply as directed under disinfection.

Fungicidal

VORTEXX can be used on hard nonporous inanimate surfaces such as shower room floors, locker room benches, shower stalls and bath mats. At 0.2% (1oz./4 gallons) the product is effective against *Trichophyton mentagrophytes* (Athletes Foot Fungi) and *Candida albicans* (pathogenic yeast) in the presence of protein (5% blood serum) and soap scum in 500 ppm hard water with a 10 minute contact time. Apply as directed under disinfection.

STERILIZATION OF MANUFACTURING, FILLING, AND PACKAGING EQUIPMENT IN ASEPTIC PROCESSES

Prior to use of this product, remove gross soil particles from processing surfaces, then wash with a recommended detergent solution, followed by a thorough potable water rinse. Prepare a 5% sterilizing solution of VORTEXX. Circulate, coarse spray, or flood the sterilizing solution as appropriate to the equipment. All surfaces should be exposed to the sterilizing solution for 30 minutes. This product is effective against *Clostridium sporogenes* and *Bacillus subtilis* at 5% (25 oz./4 gallons) in 100 ppm hard water and 5% organic load at 20° C. Thoroughly rinse food contact surfaces with either a sterile water or potable water rinse. This product is not for use as a sterilant on medical devices.

DISINFECTING PHARMACEUTICAL AND COSMETIC SURFACES

VORTEXX is recommended for use on hard, non-porous, environmental surfaces such as floors, walls and processing equipment in pharmaceutical and cosmetic processing facilities.

VORTEXX is effective against *Staphylococcus aureus*, *Salmonella choleraesuis*, and *Pseudomonas aeruginosa* at 0.2% (1 oz./4 gallons) in hard water (500 ppm as CaCO₃), 5% blood serum and dried soap film residue. For heavily soiled areas a precleaning step is required. Rinse all surfaces thoroughly with the disinfecting solution and maintain a contact time of at least 10 minutes. Product contact surfaces must be rinsed with sterile water.

ACCEPTED
with COMMENTS
EPA Letter Dated:

POULTRY HOUSE DISINFECTION

VORTEXX is effective against the Avian Influenza A (H3N2) virus.

Special Instructions for Inactivating Avian Influenza A.

1. Remove all poultry and feeds from premises, trucks, coops, and crates.
2. Remove all litter and droppings from floors, walls, and surfaces of facilities occupied or traversed by poultry.
3. Empty all troughs, racks, and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with water.
5. Saturate surfaces with the recommended disinfecting solution for a period of 10 minutes.
6. Ventilate buildings, coops, and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set, or dried.
7. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains, and waterers with soap and detergent, and rinse with potable water before reuse.

JAN - 8 2008

added to EPA Pesticide, Fungicide, and Rodenticide Act as amended, for the pesticide, registered under EPA Reg. No. 1677-158

FARM PREMISE DISINFECTION

VORTEXX is effective against the Avian Influenza A (H3N2) virus.

Special Instructions for Inactivating Avian Influenza A.

1. Remove all animals and feed from premises, vehicles, and enclosures.
2. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes, and other facilities and fixtures occupied or traversed by animals.
3. Empty all troughs, racks, and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with water.
5. Saturate all surfaces with the recommended disinfecting solution for a period of 10 minutes.
6. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
7. Ventilate buildings, cars, boats, and other closed spaces. Do not house livestock or employ equipments until treatment has been absorbed, set, or dried.
8. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains, and waterers with soap and detergent and rinse with potable water before reuse.

8098

VORTEX label

08/17/07

NOTE: This product in its use solutions is compatible with stainless steel and aluminum surfaces. If product is intended to be used on any other surface, it is recommended that you apply product to a smaller test area to determine compatibility before proceeding with its use.

STORAGE & DISPOSAL

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL

PESTICIDE STORAGE: Product should be kept cool and in a vented container to avoid any explosion hazard.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

(Totes) Verify that the tote is empty. Do not rinse or clean. Seal tote and contact Ecolab for return.

FOR COMMERCIAL USE ONLY
STRONG OXIDIZING AGENT

EPA Reg. No. 1677-158
EPA Est. 1677-MN-1 (P), 60156-IL-1 (SE), 1677-CA-2(R),
1677-TX-1(D), 1677-OH-1(H), 1677-IL-1(J), 1677-GA-1(M),
1677-PR-1(B), 1677-CA-1(S), 1677-WV-1(V)
Superscript refers to first letter of date code

Ecolab Inc., Food & Beverage Division
370 Wabasha Street No.
St. Paul, MN 55102

U.S. Patent No. 5200189 and 5314687
Other Patents Pending

Net Contents:	1 U.S. Gallon (3.78 L)
	4 U.S. Gallons (15.1 L)
	2.5 U.S. Gallons (9.45 L)
	15 U.S. Gallons (56.8 L)
	30 U.S. Gallons (113.5 L)
	50 U.S. Gallons (189 L)
	300 U.S. Gallons (tote)

ACCEPTED
with COMMENTS
EPA Letter Dated:
JAN - 8 2008

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No.
1677-158