	ACCEPT		
ECOLAB.	FEB 0 6 2001		EPA Reg. No. 1677-164 EPA Est. 60156-IL-1
Oxy-15	Under the Federal Insecticida, Fung Rodenticide Act as a hendau, for th pesticide, registered under 2PA Reg. No. 1677		
Water Additive for Controlling the Growth of N		spoilage on Fresh Cut, Post Harvest, and Further Processed Fruits and Vegetables in transport, storage and processir	ng
ACTIVE INGREDIENTS: Peroxyacetic acid	AND DOMESTIC ANIMALS Imful or fatal if swallowed. Do not get in loves, and protective clothing if handling ng. Remove any contaminated clothing Get medical attention. act lenses. Hold eyelids open and flush k large quantities of water. Avoid alcohol. A POISON CONTROL CENTER OR REE: 1-800-328-0026 indicate the use of gastric lavage. form. Mix only with water according to the r sanitizers, cleaners, or organic fish, and aquatic invertebrates. Do not bonds, estuaries, oceans, or other waters ant Discharge Elimination System (NPDES) prior to discharge. Do not discharge effluent ifying the local sewage treatment plant	<ul> <li>U.S. Patent No. 5,409,713 Other Patents Pending</li> <li>Dxy-15 is intended for use in the control of microorganisms (or bactena) which cause decay (and/or spoilage) and that may be present in the process water and on the surface of frex-out and post-harvest fruits and vegetables, as well as processed fruits and vegetables. Examples of process water are flumes, chill tanks and wash water systems. This product is not intended for control of any public health organisms.</li> <li>DIRECTIONS FOR USE: It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. CORTECATMENT OF PROCESS WATER STREAMS:</li> <li>A. Batch systems with no makeup water added: <ol> <li>Fill vessel containing fruits or vegetables with known amount of water.</li> <li>Ensure that water is recirculating in vessel.</li> </ol> </li> <li>Add Cxy-15 to no more than 533 pom (wt/wt) lotal product (80 ppm residual peroxyacetic acid) in use solution. This can be accomplished by initially adding 53.3 grams (47.8 mis) Cxy-15 per 100 liters of water, or 1.0 fluid ounces Cxy-15 per 16.4 gallons of water.</li> <li>Continuous systems with constant addition of makeup water: Initial dose (10 ky-15).</li> <li>Ensure that system is recirculating in process water up to an initial property dosed level of Cxy-15).</li> <li>Ensure that system is recirculating with known amount of water in vessels and pping.</li> <li>Add initial dose of Cxy-15 to no more 533 ppm (wt/wt) lotal product (80 ppm residual peroxyacetic acid) in use solution. This can be accomplished by initially 53.3 grams (47.8 mis) Cxy-15 per 100 liters of water, or 1.0 fluid ounces Cxy-15 per 16.4 gallons of water.</li> <li>Contact time of 45 seconds minimum is recommended.</li> </ul> Empression water and yselem is set staady stale dosing of Oxy-15 is maintained). Meter no more than 533 ppm (wt/wt) Oxy-15 lotal product (80 ppm residual peroxyacetic acid) in use solution. This can be accomplished by adding 53.3 grams (47.8 mis) per minute of Cxy-15. Makeup	<ul> <li>Oxy-15 can be used on the following types of fresh, post harvest and further processed fruits and vegetables: <u>Vegetables</u></li> <li>Root and tuber vegetables: Carrot, potato, radish, rutabaga, sweet potato, yam, sugar beet</li> <li>Leaves of root and tuber vegetables: Turnip greens, and sugar beet</li> <li>Bulb vegetables: Onion (dry butb and green), leek, garlic, shallot</li> <li>Leaves of root and tuber vegetables: Turnip greens, and sugar beet</li> <li>Bulb vegetables: Lettuce (head and leaf), celery, fennel, endive, escarole, parsley, radicchio, rhubarb, spinach</li> <li>Brassica leafy vegetables: Broccoti, Brussel sprouts, cabbage, cauliflower, mustard greens, mustard spinach</li> <li>Legumes [succulent or dried], bean (green, kidney, lima, mung, navy, pinto, snap, wax), pea (chickpea, lentil, dwarf, garden, English; field, edible pea pod), alfalfa, and soybean</li> <li>Fruiting vegetables: Pepper (bell, pimento, hot, sweet), tomato, tomatillo, eggplant</li> <li>Cucurbits: Cucumber, melon (cantaloupe, crenshaw melon, honeydew, honey ball melon, mango melon, muskmelon, pineapple melon, watermelon), summer squash, pumpkins, winter squash <u>Fruits</u></li> <li>Otirus fruits: Sweet and sour orange, lemon, lime, tangelo, tangerine, mandarin, citrus citron, kumquats, grapefruit</li> <li>Pome fruits: Apples and pears</li> <li>Stone fruits: Sour or sweet cherry, peach, nectarine, plum, prune</li> <li>Small fruits and berries: Blackberries, blueberries, red and black raspberries</li> <li>Sprouts and seeds of vegetables and fruits that are listed on this label including, root &amp; tuber vegetables, legumes, fruiting vegetables, cucurbits, citrus fruits, pome fruits, stone fruits, small fruits and berries, mustard</li> <li><u>Tree nuts</u>: Almond, Brazil, fibert, cashew, pecan , walnut (black &amp; English), macadamia, chestrut</li> <li><u>Cereal grains</u>: Com, barley, oats, rice, wheat, triticale, wild rice, sweet com</li> <li><u>Herbs and Spices</u>: Basil, chives, coriander, dill,lemongrass marjoram, sage, savory, tarragon, thy</li></ul>
STANG OXIDIZING AGENT		in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. [300 gallon tote] Verify that the tote is empty. Do not rinse or clean. Seal tote and contact Ecolab for return.	Miscellaneous: Asparagus, avocado, artichoke, banana, cranberry, fig, grape, kiwifruit, mango, mushroom, okra, peanut, persimmon, pineapple, raisins, strawberry, water chestnut, watercress
	FOOD & BEVERAGE DIVISION Ecolab Inc. 370 Wabasha Street St. Paul, MN 55102	NET CONTENTS: 50 U.S. Gals. (189 liters) or 300 U.	