

1677-164

3/13/2014

1/7



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

MAR 13 2014

Mr. Ron Derbyshire
Senior Manger, NA Biocides
Ecolab, Inc.
370 N. Wabasha Street
Saint Paul, MN 55102

Subject: Tsunami 100
EPA Registration Number 1677-164
Your Amendment Dated October 21, 2013
EPA Received Date October 25, 2013

The following amendment, submitted in connection with registration under section 3(c)(7)(B) of the Federal Insecticide, Fungicide, and Rodenticide Act, FIFRA, as amended, to add labeling for use dilution container to secondary container labeling, is acceptable.

The Confidential Statement of Formula, dated March 10, 2014, is acceptable.

A stamped copy of the labeling is enclosed. Submit three (3) copies of your final printed labeling before distributing or selling the product bearing the revised labeling.

Submit and/or cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.

If the above conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

If you have questions concerning this letter, please contact Karen M. Leavy at (703)-308-6237.

Sincerely,

Marshall Swindell
Product Manager 33
Regulatory Management Branch I
Antimicrobials Division (7510P)

2/7


Tsunami 100

Water Additive for Pathogen* Reduction in
(Fruit and Vegetable) Processing Water (and) Controlling the Growth of Spoilage and
Decay Causing Non-Public Health Organisms (on Fruit and Vegetable Surfaces)

For Organic Production.

Tsunami 100 may be used as a water additive in (fruit and vegetable) processing water on products labeled as organic in food processing facilities (on both raw agricultural commodities and on fruits and vegetables that will be further processed).

Active Ingredients:	
Peroxyacetic acid.....	15.2%
Hydrogen peroxide.....	11.2%
Inert Ingredients:	73.6%
Total:	100.0%



**KEEP OUT OF REACH OF CHILDREN
DANGER**

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CORROSIVE: Causes severe eye damage and skin burns. Harmful or fatal if swallowed. Do not get in eyes, on skin, or on clothing. Wear chemical goggles, rubber gloves, and protective clothing if handling concentrate. Wash thoroughly with soap and water after handling and before eating, drinking, and chewing gum, or using tobacco. Remove any contaminated clothing and wash before re-use.

FIRST AID

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 –20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

FOR EMERGENCY MEDICAL INFORMATION CALL TOLL-FREE: 1-800-328-0026

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

See side/back panel for First Aid

PHYSICAL AND CHEMICAL HAZARDS:

Strong oxidizing agent. Corrosive. Do not use in concentrated form. Mix only with water according to label instructions. Never bring concentrate in contact with other sanitizers, cleaners or organic substances.

ENVIRONMENTAL HAZARDS: (containers 5 gallons or greater) This product is toxic to birds, fish, and aquatic invertebrates. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

Used as directed, *Tsunami 100* reduces 99.9% of the pathogens *Escherichia coli* O157:H7 (ATCC 43895, 35150, 43890)*, *Listeria monocytogenes* (ATCC 49594, 19114, 19116)* and *Salmonella enterica* (ATCC 10721, 6962, 13311)* in fruit and vegetable processing waters. *Tsunami 100* also provides control of spoilage and decay causing non-public health organisms present in processing waters and on the surface of post-harvest, fresh-cut and processed fruits and vegetables.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FOR PATHOGEN* REDUCTION AND CONTROL IN FRUIT AND VEGETABLE PROCESSING WATERS:

- A. Batch systems with no makeup water added:
1. Ensure that water is mixing in the processing vessel.
 2. Add *Tsunami 100* at a rate from 2.5-6.7 fluid ounces per 100 gallons of process water. This will produce about 215-575 ppm total product and about 30-80 ppm peroxyacetic acid. At this use dilution, *Tsunami 100* will provide a 99.9% reduction against the pathogens *Escherichia coli* O157:H7*, *Listeria monocytogenes** and *Salmonella enterica**.
 3. Measure the residual peroxyacetic acid concentration in the water using a Test Kit (consult Ecolab Representative) and adjust dose as needed. Allow a 1.5 minute mixing time.
- B. Continuous systems with makeup water continuously added:
- Initial dose:**
1. Ensure that water is mixing in the processing vessel and/or piping.
 2. Add *Tsunami 100* at a rate from 2.5-6.7 fluid ounces per 100 gallons of process water. This will produce about 215-575 ppm total product and about 30-80 ppm peroxyacetic acid. At this use dilution, *Tsunami 100* will provide a 99.9% reduction against the pathogens *Escherichia coli* O157:H7*, *Listeria monocytogenes** and *Salmonella enterica**.
 3. Measure the residual peroxyacetic acid concentration in the water using a Test Kit (consult Ecolab Representative) and adjust dose as needed. Allow a 1.5 minute mixing time.
- Continuous Dosing:**
Meter *Tsunami 100* at a rate from 2.5-6.7 fluid ounces per 100 gallons of fresh makeup water added to the system. This will produce about 215-575 ppm total product and about 30-80 ppm peroxyacetic acid. Measure the residual peroxyacetic acid concentration in the water using a Test Kit (consult Ecolab Representative) and adjust dose as needed. Allow a 1.5 minute mixing time.

FOR TREATMENT OF FRUIT AND VEGETABLE SURFACES AND PROCESS WATERS:

This product is not intended for control of any public health organisms on fruit and vegetable surfaces. Mix *Tsunami 100* with water either batch-wise or continuously to produce about 36 - 575 ppm total product and about 5 - 80 ppm peroxyacetic acid in use solution. This can be accomplished by initially adding *Tsunami 100* at a rate from 0.42 - 6.7 fluid ounces per 100 gallons of process water. The fruits and vegetables can be sprayed or submerged in the resulting solution for a minimum contact time of 90 seconds, followed by adequate draining. At this use dilution, *Tsunami 100* will control the growth of spoilage and decay causing non-public health organisms, including odor causing organisms, in process waters and on the surface of fruits and vegetables.

Tsunami 100 can be used on the following types of fresh, post harvest and further processed fruits and vegetables:

Vegetables

- ◆ Root and tuber vegetables: Carrot, potato, radish, rutabaga, sweet potato, yam, sugar beet
- ◆ Leaves of root and tuber vegetables: Turnip greens and sugar beet
- ◆ Bulb vegetables: Onion (dry bulb and green), leek, garlic, shallot
- ◆ Leafy vegetables: Lettuce (head and leaf), celery, fennel, endive, escarole, parsley, radicchio, rhubarb, spinach
- ◆ Brassica leafy vegetables: Broccoli, Brussel sprouts, cabbage, cauliflower, mustard greens, mustard spinach

- ◆ Legumes [succulent or dried], bean (green, kidney, lima, mung, navy, pinto, snap, wax), pea (chickpea, lentil, dwarf, garden, English, field, edible pea pod), alfalfa, and soybean
- ◆ Fruiting vegetables: Pepper (bell, pimento, hot, sweet), tomato, tomatillo, eggplant
- ◆ Cucurbits: Cucumber, melon (cantaloupe, crenshaw melon, honeydew, honey ball melon, mango melon, muskmelon, pineapple melon, watermelon), summer squash, pumpkins, winter squash

Fruits

- ◆ Citrus fruits: Sweet and sour orange, lemon, lime, tangelo, tangerine, mandarin, citrus citron, kumquats, grapefruit
- ◆ Pome fruits: Apples and pears
- ◆ Stone fruits: Sour and sweet cherry, peach, nectarine, plum, and prune
- ◆ Small Fruits and berries: Blackberries, blueberries, red and black raspberries

Sprouts and seeds of: vegetables and fruits that are listed on this label including, root & tuber vegetables, bulb vegetables, leafy vegetables, *Brassica* leafy vegetables, legumes, fruiting vegetables, cucurbits, citrus fruits, pome fruits, stone fruits, small fruits and berries, mustard

Tree nuts: Almond, Brazil, filbert, cashew, pecan, walnut (black & English), macadamia, chestnut

Cereal grains: Corn, barley, oats, rice, wheat, triticale, wild rice, sweet corn

Herbs and Spices: Basil, chives, coriander, dill, lemongrass, marjoram, sage, savory, tarragon, thyme

Miscellaneous: Asparagus, avocado, artichoke, banana, cranberry, fig, grape, kiwifruit, mango, mushroom, okra, peanut, persimmon, pineapple, raisins, strawberry, water chestnut, watercress, coffee berry, coffee bean, seaweed

FOR TREATMENT OF SEEDS NOT INTENDED FOR HUMAN OR ANIMAL CONSUMPTION:

Apply to seeds as directed to control seedborne microorganisms that cause plant disease or spoilage and decay of developing seedlings. Only treat seeds of the crops listed on this label. Mix *Tsunami 100* with clean water either batchwise or continuously to no more than 11,500 ppm total product (1750 ppm residual peroxyacetic acid) in use solution. This can be accomplished by adding 20 fluid ounces *Tsunami 100* per 16.4 gallons of water. The volume of treatment solution should be at least two times greater than the volume of seeds to be treated. The seeds should be submerged in the treatment solution and agitated for 30 minutes. Following treatment, remove seeds from treatment solution and dry.

FOR PATHOGEN* REDUCTION AND CONTROL IN (INDUSTRIAL) PROCESSING WATERS:

A. Batch systems with no makeup water added:

1. Ensure that water is mixing in the processing vessel.
2. Add *Tsunami 100* at a rate from 2.5-6.7 fluid ounces per 100 gallons of process water. This will produce about 215-575 ppm total product and about 30-80 ppm peroxyacetic acid. At this use dilution, *Tsunami 100* will provide a 99.9% reduction against the pathogens *Escherichia coli* O157:H7*, *Listeria monocytogenes** and *Salmonella enterica**.
3. Measure the residual peroxyacetic acid concentration in the water using a Test Kit (consult Ecolab Representative) and adjust dose as needed. Allow a 1.5 minute mixing time.

B. Continuous systems with makeup water continuously added:

Initial dose:

1. Ensure that water is mixing in the processing vessel and/or piping.
2. Add *Tsunami 100* at a rate from 2.5-6.7 fluid ounces per 100 gallons of process water. This will produce about 215-575 ppm total product and about 30-80 ppm peroxyacetic acid. At this use dilution, *Tsunami 100* will provide a 99.9% reduction against the pathogens *Escherichia coli* O157:H7*, *Listeria monocytogenes** and *Salmonella enterica**.
3. Measure the residual peroxyacetic acid concentration in the water using a Test Kit (consult Ecolab Representative) and adjust dose as needed. Allow a 1.5 minute mixing time.

Continuous Dosing:

Meter *Tsunami 100* at a rate from 2.5-6.7 fluid ounces per 100 gallons of fresh makeup water added to the system. This will produce about 215-575 ppm total product and about 30-80 ppm peroxyacetic acid. Measure the residual peroxyacetic acid concentration in the water using a Test Kit (consult Ecolab Representative) and adjust dose as needed. Allow a 1.5 minute mixing time.

FOR TREATMENT OF (INDUSTRIAL) PROCESS WATERS:

This product is not intended for control of any public health organisms. Mix *Tsunami 100* with water either batch-wise or continuously to produce about 36 - 575 ppm total product and about 5 - 80 ppm peroxyacetic acid in use solution. This can be accomplished by initially adding *Tsunami 100* at a rate from 0.42--6.7 fluid ounces per 100 gallons of process water. At this use dilution, *Tsunami 100* will control the growth of spoilage and decay causing non-public health organisms, including odor causing organisms, in process waters.

SANITIZATION

Tsunami 100 acid sanitizer is recommended for use on pre-cleaned surfaces such as equipment, pipelines, tanks, vats, fillers, evaporators, pasteurizers and aseptic equipment dairies, breweries, wineries, beverage and food processing plants. This product is effective as a sanitizer when solution is prepared in water of up to 500 ppm hardness.

SANITIZING FOOD CONTACT SURFACES

Prior to use of this product, remove gross soil particles from surfaces, thoroughly clean surfaces, and follow with a potable water rinse. Sanitize clean surfaces with a concentration of 0.10 – 0.18% v/v (1000 to 1800 ppm v/v or 1 to 1.8 ounces per 8 gallons of water) at room temperature. Use immersion, coarse spray or circulation techniques as appropriate to sanitize the surfaces. All surfaces must be exposed to the sanitizing solution for a period of not less than one minute unless otherwise specified by governing sanitary code. Allow surfaces to drain thoroughly and air dry. Do not rinse. This product when used per label directions is effective against *Staphylococcus aureus* (ATCC 6538) and *Escherichia coli* (ATCC 11229).

CLEANING HARD SURFACE**CLEANING HARD SURFACE FOOD PROCESSING EQUIPMENT– NO RINSE**

For hard surface cleaning applications, remove gross soil particles from surfaces, then thoroughly clean surfaces with a concentration of 0.10 – 0.18% v/v (1000 to 1800 ppm v/v or 1 to 1.8 ounces per 8 gallons of water). Use immersion, coarse spray or circulation techniques as appropriate to clean surfaces. Allow surfaces to drain thoroughly. No rinse necessary.

FINAL BOTTLE AND CLOSURE CLEANING RINSE

Tsunami 100 may be used as a final cleaning rinse for returnable and non-returnable bottles (e.g. glass or PET) and closures not requiring a final food contact surface sanitizing rinse when used at a concentration of 0.10 – 0.18% v/v (1000 to 1800 ppm v/v or 1 to 1.8 ounces per 8 gallons of water). Drain thoroughly. No rinse necessary.

CLEANING HARD SURFACE PROCESSING EQUIPMENT– RINSE FOR FOOD CONTACT SURFACES

For hard surface cleaning applications, remove gross soil particles from surfaces, then thoroughly clean surfaces with 2 to 3 ounces product per 3 gallons of water (0.5% to 0.8% v/v). All treated hard non-porous food contact surfaces must be rinsed thoroughly with a potable water rinse.

FLOODING AND PRODUCED WATER

For Water Flooding operations, add *Tsunami 100* initially at 3.75 fluid ounces per 1000 gallons of water (5ppm peroxyacetic acid by weight) to 75.5 fluid ounces per 1000 gallons of water (100 ppm peroxyacetic acid by weight) and repeat until control is achieved. Subsequent treatment may be continued on a weekly basis or as required.

NOTE: This product in its use solutions is compatible with stainless steel and aluminum surfaces. If product is intended to be used on any other surface, it is recommended that you apply product to a smaller test area to determine compatibility before proceeding with its use.

STORAGE & DISPOSAL:**DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL**

PESTICIDE STORAGE: Product should be kept cool and in a vented container to avoid any explosion hazard.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING AND DISPOSAL:

(For containers 5 gallons or less.) Non-refillable container. Do not reuse or refill this container. Triple rinse as follows: Fill container ¼ full with water and recap. Shake for 10 seconds. Drain for 10 seconds after the flow begins to drip. Follow Pesticide Disposal instructions for rinsate disposal. Repeat procedure two more times. Then offer for recycling or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.

(For containers >5-55 gallons.) Non-refillable container. Do not reuse or refill this container. Triple rinse as follows: Empty remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat procedure two more times. Then offer for recycling or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.

(300 gallon tote) Verify that the tote is empty. Do not rinse or clean. Seal tote and contact Ecolab for return.

FOR INSTITUTIONAL USE

STRONG OXIDIZING AGENT

EPA Reg. No. 1677-164

EPA Est.: 1677-MN-1 (P), 60156-IL-1 (SI), 1677-CA-2 (R), 1677-TX-1 (D), 1677-OH-1 (H), 1677-IL-2 (J), 1677-PR-1 (B), 1677-CA-1 (S), 1677-GA-1 (M), 1677-WV-1 (V)

Superscript refers to first letter of date code

<p>Net Contents: 4 U.S. Gallons (15.1 liters) 50 U.S. Gallons (189 liters) 300 U.S. Gals. (1134 liters)</p>
--

Manufactured by: Ecolab Inc., Food & Beverage Division
370 N. Wabasha Street, St. Paul, MN 55102

7/7

SECONDARY CONTAINER LABEL

(Note to reviewer: This secondary container label will be used only when the product is diluted with no more than 20 oz. product per 16.4 gallon of water, 0.95% v/v.)

Tsunami 100

Concentrate Ingredient Statement

Active Ingredients:

Peroxyacetic acid	15.2%
Hydrogen peroxide	11.2%

Inert Ingredients:	<u>73.6%</u>
Total:	100.0%

The product in this container must be diluted at the use rate with no more than 20 oz. product per 16.4 gallon of water, or 0.95% v/v, as directed on the concentrate product label.

Dilution: _____ oz product per _____ gallons of water

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

**USE DILUTION PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Causes moderate eye irritation. Harmful if swallowed. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

If Swallowed: Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

Follow the directions for use listed on the concentrate label when applying this product.
EPA Reg. No. 1677-164