

1677-209

6/1/2015

1/9



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

June 1, 2015

Nicole Listner
Associate Regulatory Specialist 1
Ecolab Inc.
370 N. Wabasha Street
St. Paul, MN 55102

Subject: Notification per PRN 98-10 – Minor Label Revisions-
Product Name: OCTAVE FS
EPA Registration Number: 1677-209
Application Date: May 15, 2015
Decision Number: 505162

Dear Nicole Listner:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the above referenced product. The Antimicrobials Division (AD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10.

The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, you may contact Tuere Williams at 703-347-0316 or via email at williams.tuere@epa.gov.

Sincerely,

John Hebert
Branch Chief for,
Regulatory Management Branch I
Antimicrobials Division (7510P)
Office of Pesticide Programs

OCTAVE FS

SANITIZER AND DISINFECTANT

Acid Liquid Sanitizer for Food Processing Equipment in Dairies, Dairy Farms, Breweries, Wineries, Beverage and Food Processing Plants

Disinfectant for Farms, Livestock Quarters, Poultry Premises, Poultry Hatcheries, Veterinary Clinics, Animal Life Science Laboratories, Animal Care Facilities and Industrial Facilities

Active Ingredients:

Hydrogen Peroxide.....	7.52%
Peroxyoctanoic Acid.....	1.00%
Octanoic Acid.....	2.85%

Other Ingredients:	88.63%
Total:	100.00%

NOTIFICATION

1677-209

The applicant has certified that no changes, other than those reported to the Agency have been made to the labeling. The Agency acknowledges this notification by letter dated:

06/01/2015

**KEEP OUT OF REACH OF CHILDREN
DANGER**

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: CORROSIVE. Causes irreversible eye damage and skin burns. Harmful if swallowed. Do not get in eyes, on skin or on clothing. Wear protective eyewear (goggles, face shield or safety glasses), coveralls worn over long-sleeved shirt and long pants, socks, chemical resistant gloves, and chemical resistant footwear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove contaminated clothing and wash clothing before reuse. Prolonged or frequently repeated skin contact may cause an allergic reaction in some individuals.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 -20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or going for treatment.

FOR EMERGENCY MEDICAL INFORMATION CALL TOLL FREE: 1-800-328-0026

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

PHYSICAL AND CHEMICAL HAZARDS: Strong oxidizing agent. Corrosive. Do not use in concentrated form. Mix only with water according to label instructions. Never bring concentrate in contact with other sanitizers, cleaners or organic substances. This product is not to be used around electrical equipment.

ENVIRONMENTAL HAZARDS (containers 5 gallons or greater): This pesticide is toxic to birds, fish, and aquatic invertebrates. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

(Note to reviewer: Text appearing in parentheses / brackets is optional text)

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

SANITIZATION

Octave FS acid sanitizer is recommended for use on pre-cleaned surfaces such as equipment, pipelines, tanks, vats, fillers, evaporators, pasteurizers and aseptic equipment in dairies, dairy farms, breweries, wineries, beverage and food processing plants. This product is effective as a sanitizer when solution is prepared in water of up to 500 ppm hardness as CaCO_3 .

NOTE: FOR MECHANICAL OPERATIONS, prepared use solution may not be reused for sanitizing but may be reused for other purposes such as cleaning.

FOR MANUAL OPERATIONS, fresh sanitizing solution must be prepared at least daily or more often if the solution becomes diluted or soiled.

SANITIZING FOOD CONTACT SURFACES

Prior to sanitizing, remove gross food particles, and then wash with a detergent solution, followed by a potable water rinse. Sanitize with a concentration of 1 oz. **Octave FS** concentrate per 6 to 8 gals. of water (0.1% - 0.13% v/v concentration). At this dilution **Octave FS** is an effective food contact surface sanitizer against the following organisms: *Staphylococcus aureus* (ATCC 6538), *Escherichia coli* (ATCC 11229), *Escherichia coli* O157:H7 (ATCC 43895), *Pseudomonas aeruginosa* (ATCC 15442), *Listeria monocytogenes* (ATCC 7644), *Salmonella typhimurium* (ATCC 13311), *Enterobacter sakazakii* (ATCC 12868), *Campylobacter jejuni* (ATCC 33291) and *Candida albicans* (ATCC 18804). At the same dilution, **Octave FS** is also effective for sanitization against the common spoilage organisms *Pediococcus damnosus* (ATCC 25248), *Lactobacillus malefermentans* (ATCC 11305) and *Geotrichum candidum* (ATCC 34614). Use immersion, coarse spray or circulation techniques as appropriate to the equipment. All surfaces must be exposed to the sanitizing solution for a period of not less than one minute unless a longer time is specified by the governing sanitary code. Drain thoroughly and allow to air dry. Do not rinse.

FOAM SANITIZING FOOD CONTACT SURFACES

Octave FS is an effective foam sanitizer of pre-cleaned food contact surfaces. Foam sanitize with a concentration of 1 oz. **Octave FS** concentrate per 6 to 8 gals. of water (0.1% - 0.13% v/v concentration). At this dilution **Octave FS** is an effective foaming food contact surface sanitizer against the following organisms: *Staphylococcus aureus* (ATCC 6538) and *Escherichia coli* (ATCC 11229), *Escherichia coli* O157:H7 (ATCC 43895), *Listeria monocytogenes* (ATCC 7644), *Salmonella typhimurium* (ATCC 13311), *Enterobacter sakazakii* (ATCC 12868). At the same dilution, **Octave FS** is also effective for foam sanitization against the common spoilage organisms *Pediococcus damnosus* (ATCC 25248) and *Lactobacillus malefermentans* (ATCC 11305). Use an appropriate device to generate foam intended to cover the food contact surfaces. Contact your Ecolab representative for information on a recommended foamer. All surfaces must be exposed to the sanitizing solution for a period of not less than one minute unless a longer time is specified by the governing sanitary code. Drain thoroughly and allow to air dry. Do not rinse.

CONTINUOUS TREATMENT OF CONVEYORS

Wash, rinse and sanitize conveyor equipment. During processing, apply solution containing 1 oz. **Octave FS** concentrate per 6 to 8 gals. of water (0.1% - 0.13% v/v concentration) to a conveyor with Mikro Master or other suitable dispensing equipment. At this dilution **Octave FS** is effective against *Staphylococcus aureus* (ATCC 6538), *Escherichia coli* (ATCC 11229), *Escherichia coli* O157:H7 (ATCC 43895), *Pseudomonas aeruginosa* (ATCC 15442), *Listeria monocytogenes* (ATCC 7644), *Salmonella typhimurium* (ATCC 13311), *Enterobacter sakazakii* (ATCC 12868), *Campylobacter jejuni* (ATCC 33291) and *Candida albicans* (ATCC 18804). At the same dilution **Octave FS** is effective at reducing the common spoilage microorganisms *Pediococcus damnosus* (ATCC 25248), *Lactobacillus malefermentans* (ATCC 11305) and *Geotrichum candidum* (ATCC 34614). Controlled volumes of **Octave FS** are applied to return portion of conveyor through nozzles so located as to permit maximum drainage of **Octave FS** from equipment and to prevent puddles on top of belt. During interruptions in operations, coarse spray the processing

equipment with **Octave FS** solution at not more than 0.13% v/v concentration. Conveyor equipment must be free of product when applying coarse spray. Conveyor surface should be exposed to the sanitizing solution for a period of not less than one minute.

SANITIZING NON-FOOD CONTACT SURFACES

Pre-clean surfaces as directed above. Sanitize non-food contact surfaces such as floors, walls, tables, chairs, benches, troughs, and drip pans with 1 to 5 oz. **Octave FS** concentrate per 8 gals. of water (0.1% -0.5% v/v concentration). At this concentration the product is effective as a non-food contact surface sanitizer against the following organisms: *Staphylococcus aureus* (ATCC 6538), *Enterobacter aerogenes* (ATCC 13048), *Listeria monocytogenes* (ATCC7644), *Salmonella typhimurium* (ATCC 13311), *Pseudomonas aeruginosa* (ATCC 15442), *Enterobacter sakazakii* (ATCC 12868) and *Geotrichum candidum* (ATCC 34614). All surfaces must be exposed to the sanitizing solution for a period of not less than 5 minutes. Drain thoroughly and allow to air dry. No rinse necessary.

FOAM SANITIZING NON-FOOD CONTACT SURFACES (This use not approved in the state of California)

Octave FS is an effective foam sanitizer of pre-cleaned non-food contact surfaces, such as boots, floors, walls and associated equipment. To foam sanitize, prepare a solution of 1 to 5 oz. **Octave FS** concentrate per 8 gals. of water (0.1% - 0.5% v/v concentration) and 1.4 oz. **Liquid K** per 8 gals. (0.13% v/v concentration). **Liquid K** is the only approved foam generator. At this concentration the product is effective as a non-food contact surface sanitizer against *Staphylococcus aureus* (ATCC 6538), *Enterobacter aerogenes* (ATCC 13048), *Listeria monocytogenes* (ATCC7644), *Salmonella typhimurium* (ATCC 13311), *Enterobacter sakazakii* (ATCC 12868) and *Campylobacter jejuni* (ATCC 33291). Use an appropriate device to generate foam intended to cover non-food contact surfaces. Wet surfaces thoroughly. Surfaces should be exposed to the sanitizing foam for a period of not less than 5 minutes. No rinse is necessary. Contact your Ecolab representative for information on a recommended foamer.

FOAM SANITIZING NON-FOOD CONTACT SURFACES

Octave FS is an effective foam sanitizer of pre-cleaned non-food contact surfaces, such as boots, floors, walls and associated equipment. Sanitize with a concentration of 1 to 5 oz. **Octave FS** concentrate per 8 gallons of water (0.1% - 0.5% v/v concentration). At this concentration the product is effective as a non-food contact surface sanitizer against *Staphylococcus aureus* (ATCC 6538), *Enterobacter aerogenes* (ATCC 13048), *Listeria monocytogenes* (ATCC7644), *Salmonella typhimurium* (ATCC 13311), *Enterobacter sakazakii* (ATCC 12868) and *Campylobacter jejuni* (ATCC 33291). Use an appropriate device to generate foam intended to cover non-food contact surfaces. Wet surfaces thoroughly. Surfaces must be exposed to the sanitizing foam for a period of not less than 5 minutes. No rinse is necessary. Contact your Ecolab representative for information on a recommended foamer.

ENTRYWAY FOAM SANITIZING SYSTEMS (This use not approved in the state of California)

To prevent cross contamination from area to area, set the system to deliver sanitizing solution at 1 oz. **Octave FS** per 1 to 8 gals. of water (0.1% - 0.78% v/v).

Alternately, **Octave FS** can be used along with **Liquid K** to apply the foam by setting the system to deliver 1 oz. **Octave FS** per 1 to 8 gals. of water (0.1% - 0.78% v/v) and 1.4 oz. **Liquid K** per 8 gals. of water (0.13% v/v concentration).

The foam (or spray) must cover the entire path of the doorway. Set the system so that a continuous wet blanket of sanitizer solution is delivered to the floor.

SHOE BATH SANITIZER DIRECTIONS:

To prevent tracking (into animal areas)(and the packaging and storage areas of food plants), shoe baths containing one inch of freshly made solution must be placed at all entrances to buildings and hatcheries. Scrape waterproof work boots (shoes) and place in solution of 1 oz. **Octave FS** per 6 to 8 gals. of water (0.1% - 0.13% v/v) for one minute prior to entering area. (If there is a heavy soil load or excessive traffic place work (boots) (shoes) in a use-solution of 1 oz. per 1 to 8 gals. of water (0.1 - 0.78% v/v) (or equivalent dilution) for one minute prior to entering area). Change the solution in the bath daily or more often if solution appears diluted or soiled.

SHOE FOAM SANITIZER DIRECTIONS

Octave FS can be used to prevent tracking into processing, packaging and storage areas of food and beverage plants. Apply a foam layer approximately 0.5 to 2 inches thick made from a solution of 1 oz. per 1 to 8 gals. of water (0.1% - 0.78% v/v) at all entrances to buildings, hatcheries, production and packaging rooms by using a foam generating machine or aerator to apply foam layer. Follow the foaming directions as specified by the manufacturer of the foam generator/aerator. Scrape waterproof shoes.

Stand and/or walk through foamed area for one minute prior to entering area. Foam area must be washed and replaced daily or when it appears dirty.

DISINFECTION

Octave FS disinfects as it cleans in one operation. **Octave FS** can be used to disinfect floors, walls and other hard nonporous surfaces such as tables, chairs, countertops, bathroom fixtures, sinks, shelves, racks, carts, exterior surfaces of refrigerators and coolers, tile, linoleum, vinyl, glazed porcelain, plastic (such as polypropylene and polyethylene), stainless steel, or glass. Areas of use include dairies, dairy farms, breweries, wineries, beverage and food processing plants, poultry farms, veterinary facilities, animal care facilities and industrial facilities.

Before use in federally inspected meat and poultry food processing plants and dairies, food products and packaging material must be removed from the room or carefully protected.

COMBINATION DISINFECTION AND CLEANING

Octave FS is effective as a disinfectant at a concentration of 1 – 2 oz. **Octave FS** concentrate per 2 gals. (0.39% - 0.78% v/v concentration) of hard water (500 ppm as CaCO₃), in the presence of 5% blood serum. At this dilution, **Octave FS** is effective against *Staphylococcus aureus* (ATCC 6538), *Salmonella enterica* (ATCC 10708), *Pseudomonas aeruginosa* (ATCC 15442) and *Candida albicans* (ATCC 18804). For heavily soiled areas a pre-cleaning step is required. Prepare a disinfecting and cleaning solution by diluting 1 – 2 oz. **Octave FS** concentrate per 2 gals. of water (0.39% - 0.78% v/v concentration). Apply use solution with mop, cloth, sponge, brush, scrubber, or coarse spray device or by soaking or foaming so as to wet all surfaces thoroughly. Allow to remain wet for 10 minutes, and then remove solution and entrapped soil with a clean wet mop, cloth, or wet vacuum pickup. Prepare a fresh solution daily or when it becomes soiled or diluted.

FOAM DISINFECTION

Octave FS is an effective foam disinfectant and cleaner on hard, non-porous surfaces, such as boots, floors, walls, drains, and associated equipment. For this application, prepare a solution of 1 – 2 oz. **Octave FS** concentrate per 2 gals. of water (0.39% - 0.78% v/v concentration). Apply solution as a foam using recommended equipment. Wet surfaces thoroughly. At this concentration, the product is effective against the following: *Staphylococcus aureus* (ATCC 6538), *Salmonella enterica* (ATCC 10708), *Pseudomonas aeruginosa* (ATCC 15442). Surfaces must be exposed to the disinfecting foam for a period of not less than 10 minutes. No rinse is necessary. Contact your Ecolab representative for information on a recommended foamer.

VIRUCIDAL

Octave FS is an effective virucide at a concentration of 1 - 2 oz. per 2 gals. (0.39%-0.78% v/v concentration) of hard water (500 ppm as CaCO₃), in the presence of 5% blood serum. At this dilution, **Octave FS** is effective against Influenza A (H1N1), Influenza A (H3N2), Influenza B, Reovirus, Influenza A, Norovirus, Feline calicivirus and Avian Influenza when used at 20°C with a 10 minute contact time. Apply as directed under Disinfection.

DISINFECTION OF POULTRY PREMISES, TRUCKS, COOPS AND CRATES

1. Remove all poultry and feed from premises, trucks, coops and crates.
2. Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by poultry.
3. Empty all troughs, racks and other feeding and watering appliances.
4. Thoroughly clean all surfaces with a detergent and rinse with water.

5. Saturate surfaces with a 1 - 2 oz. per 2 gals. (0.39%-0.78% v/v concentration) solution of **Octave FS** for a period of 10 minutes.
 6. Ventilate buildings, coops and other closed spaces. Do not house poultry or employ equipment until treatment has been absorbed, set or dried.
 7. Thoroughly scrub treated feed racks, troughs, automatic feeders, fountains and waterers with a detergent and rinse with potable water before reuse.
- See your Ecolab Representative for specific recommendations for all cleaning and rinsing requirements.

POULTRY HATCHERY DISINFECTION

Clean out any remaining eggs and chicks. Remove gross soils, such as litter, down, shell fragments or other hatching related debris. Empty all racks and other equipment. Thoroughly wash all surfaces, including floors, walls, conveyors, trays and water systems with a recommended detergent. Rinse thoroughly with water. Apply a 1 - 2 oz. per 2 gals. (0.39%-0.78% v/v concentration) solution of **Octave FS** with a mop, cloth, brush, foam or coarse spray. Wet all surfaces and allow to remain wet for 10 minutes. Ventilate buildings and other closed spaces. Allow to dry before reintroducing eggs.

DISINFECTION AND DEODORIZING OF ANIMAL HOUSING FACILITIES (BARNs, KENNELS, HUTCHES)

Remove animals and feed from facilities. Remove litter, waste matter and gross soils from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all troughs, rack and other feeding and watering equipment. Wash surfaces with a recommended alkaline detergent, by manual, foam, or spray application. Rinse with water. Apply a 1 - 2 oz. per 2 gals. (0.39%-0.78% v/v concentration) solution of **Octave FS** with a mop, cloth, brush, foam or coarse spray. Wet all surfaces and allow to remain wet for 10 minutes. Ventilate buildings and other closed spaces. Allow to air dry before reintroducing animals.

(VETERINARY PRACTICE / ANIMAL CARE / ANIMAL LABORATORY / ZOOS / PET SHOP / KENNELS DISINFECTION DIRECTIONS:)

For disinfection and deodorizing of hard nonporous surfaces of equipment used in animal housing facilities including food or water containers, utensils, instruments, cages, crates kennels, stables and catteries.

OR

(FARM PREMISE DISINFECTION DIRECTIONS:)

1. Remove all animals and feed from premises, vehicles, and enclosures.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals.
3. [For "Veterinary Practice..." :Empty all feeding and watering appliances.] [For "Farm Premise Use" Empty all troughs, racks and other feeding and watering appliances.]
4. Thoroughly clean surfaces with soap or detergent by manual, foam, or spray application and rinse with water.
5. Saturate surfaces with a use-solution of 1 – 2 oz. **Octave FS** per 2 gals. of water (0.39% - 0.78% v/v concentration or 3900 – 7800 ppm product). (or equivalent dilution) and allow to remain wet for a period of 10 minutes.
6. Immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels and scrapers used for removing litter and manure. Wipe or allow to air dry.
7. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried.
8. [For "Veterinary Practice..." Thoroughly scrub all treated feeding and watering appliances with soap or detergent, and rinse with potable water before reuse. [For "Farm Premise Use"; Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse.]

CLEANING

OCTAVE FS DIRECTIONS FOR USE AS A HARD SURFACE CLEANER

For hard surface cleaning applications, use **Octave FS** at a concentration of 1.0 – 2.7 oz. per gal. of detergent use solution (0.8-2.1% v/v or 8000 ppm – 21,000 ppm product) in water in conjunction with an alkaline pH modifier, such as KOH or NaOH to adjust cleaning solution to a pH >8.8. All hard food contact

surfaces treated with this cleaning system must be rinsed thoroughly with potable water followed by sanitizing with an approved food contact surface sanitizer.

For hard surface acid cleaning applications, remove gross food particles with a water rinse, then wash using **Octave FS** at a rate of 1.0 oz. **Octave FS** concentrate per 4.5-6.0 gallons of water (0.130 – 0.174% v/v or 1300 ppm – 1740 ppm product). All hard food contact surfaces treated with this cleaning system must be drained thoroughly, no rinse required.

NOTE: This product in its use solutions is compatible with stainless steel and aluminum surfaces. If product is intended to be used on any other surface, it is recommended that you apply product to a smaller test area to determine compatibility before proceeding with its use.

8/9

STORAGE & DISPOSAL

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL

PESTICIDE STORAGE: Product must be kept cool and in a vented container to avoid any explosion hazard.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING AND DISPOSAL:

(For containers 5 gallons or less.) Non-refillable container. Do not reuse or refill this container. Triple rinse as follows: Fill container ¼ full with water and recap. Shake for 10 seconds. Drain for 10 seconds after the flow begins to drip. Follow Pesticide Disposal instructions for rinsate disposal. Repeat procedure two more times. Then offer for recycling or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.

(For containers >5-55 gallons.) Non-refillable container. Do not reuse or refill this container. Triple rinse as follows: Empty remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat procedure two more times. Then offer for recycling or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by incineration.

(2.5 gl bladder in box) Do not reuse or refill bladder. Triple rinse as follows: Fill container ¼ full with water and recap. Shake for 10 seconds. Drain for 10 seconds after the flow begins to drip. Follow Pesticide Disposal instructions for rinsate disposal. Repeat procedure two more times. Offer box and bladder for recycling, if available, or discard in trash.

(Totes) Verify that the tote is empty. Do not rinse or clean. Seal tote and contact Ecolab for return.

**FOR COMMERCIAL USE
STRONG OXIDIZING AGENT**

EPA Reg. No. 1677-209
EPA Est. 1677-MN-1 (P), 60156-IL-1 (SI), 1677-CA-2(R),
1677-TX-1(D), 1677-IL-1(J), 1677-GA-1(M),
1677-WV-1(V)
Superscript refers to first letter of date code


Net Contents:	1 U.S. Gal. (3.78 L)
	2.5 U.S. Gals. (9.45 L)
	5 U.S. Gals. (18.9 L)
	15 U.S. Gals. (56.8 L)
	50 U.S. Gals. (189 L)
	300 U.S. Gals. (tote)

Ecolab Inc.
370 N. Wabasha Street
St. Paul, MN 55102

[Made in USA] [Made in United States of America]

Optional Marketing Language:

- Respiratory illnesses attributable to Pandemic 2009 H1N1 are caused by influenza A virus.
- This product is a broad-spectrum hard surface disinfectant that has been shown to be effective against (influenza A virus tested and listed on the label) and is expected to inactivate all influenza A viruses including Pandemic 2009 H1N1 (formerly called swine flu).
- This product has demonstrated effectiveness against influenza A virus and is expected to inactivate all influenza A viruses including Pandemic 2009 H1N1 influenza A virus.
- This product has demonstrated effectiveness against (influenza A virus tested and listed on the label) and is expected to inactivate all influenza A viruses including Pandemic 2009 H1N1 (formerly called swine flu).
- Kills Pandemic 2009 H1N1 influenza A virus (formerly called swine flu).
- Kills Pandemic 2009 H1N1 influenza A virus.
- See side/back panel for First Aid...



Antimicrobial