

FOR INDUSTRIAL AND INSTITUTIONAL USE ONLY



FORMULA 236 *Terg-o-cide*

*Cleaner-Disinfectant-Deodorizer-Fungicide-Virucide
For Hospitals and Institutional Use*

AOAC Phenol Coefficients

Staph. aureus (ATCC No. 6538) 100
Salmonella typhosa (ATCC No. 6539) 50

Active Ingredients

Didecyl dimethyl ammonium chloride 4.5%
Tetrasodium ethylenediamine tetraacetate 2.0%
Sodium carbonate 1.0%
Sodium metasilicate 0.5%

Inert Ingredients

92.0%
100.0%

EPA Registration No 1685-52 EPA Est No 1685-OH-1

TERG-O-CIDE is formulated for controlling contamination on environment surfaces in hospitals, schools, institutions and industry. It both cleans and disinfects effectively and is virucidal when used as directed. Its hard surface disinfecting action will reduce the hazard of cross-infection.

Two ounces of TERG-O-CIDE per gallon of water will deodorize bathrooms, garbage storage areas, and other areas where microorganism growth can cause mal odors

WARNING

KEEP OUT OF REACH OF CHILDREN:

Causes eye and skin irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Avoid contamination of food. Rinse empty container with water before discarding.

FIRST AID In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, do not induce vomiting. Drink large quantities of fluid and call a physician immediately.

SS-92L

USE DIRECTIONS:

Apply TERG-O-CIDE to walls, floors, and other similar hard surfaces with a cloth or mop. For heavily soiled areas, a preliminary cleaning is required.

FOR HOSPITALS AND NURSING HOMES: Add two ounces per gallon of water.

At two ounces per gallon use-level, TERG-O-CIDE is effective against Pseudomonas aeruginosa, Staphylococcus aureus, Salmonella choleraesuis, Escherichia coli, and pathogenic fungi. Germicidal performance against the first four organisms has been confirmed by the AOAC Use-Dilution Test. Fungicidal performance was determined against T. interdigitale by the AOAC Fungicidal Test.

At two ounces per gallon use-level, TERG-O-CIDE is virucidal against Herpes Simplex (a member of the virus family that causes infectious mononucleosis), Vaccinia (representative of the pox viruses), and influenza A₁ (the Hong Kong and the London flu virus) carried on inanimate environmental surfaces

FOR SCHOOLS, INSTITUTIONAL AND INDUSTRIAL USES: Add one ounce per gallon of water.

At one ounce per gallon use-level, TERG-O-CIDE delivers excellent cleaning and is germicidal. It is effective against Staphylococcus aureus, Salmonella choleraesuis, Escherichia coli, and Trichophyton interdigitale, the athlete's foot fungus. The same AOAC tests used above to confirm suitable performance for hospitals were used.

At one-half ounce per gallon of water, TERG-O-CIDE can be used for sanitizing and cleaning of utensils, equipment and other hard surfaces. When sanitizing food processing equipment and utensils and food contact surfaces, rinse thoroughly with potable water.

The **State** CHEMICAL MANUFACTURING CO.
MAIN OFFICE AND PLANT ■ CLEVELAND, OHIO 44114

NET CONTENTS

ACCEPTED