

# FORMULA NO. 235

## Heavy Duty Terg-O-Cide

### Detergent-Deodorizer-Sanitizer-Disinfectant

#### Active Ingredients

Didcyl dimethyl ammonium chloride	2.3%
Tetrasodium ethylenediamine tetraacetate	2.0%
Sodium metasilicate, anhydrous	1.5%
Trisodium Phosphate	1.5%

#### Inert Ingredients

92.7%  
100.0%

#### AOAC Phenol Coefficients

Staph. aureus (ATCC No. 6538)	50
Salmonella typhosa (ATCC No. 6539)	20

EPA Reg. No. 1685-6D

Formula 235 Heavy Duty Terg-O-Cide is a specially formulated germicidal cleaner that provides a dual action of cleaning and disinfecting.

Formula 235 is designed for use in schools, industry and institutions where heavy duty cleaning and sanitizing is required for routine maintenance. Not effective against PSEUDOMONAS AERUGINOSA when used as directed.

### DANGER

### KEEP OUT OF REACH OF CHILDREN

SEE RIGHT PANEL FOR ADDITIONAL  
PRECAUTIONARY STATEMENTS

NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

The **State** CHEMICAL MANUFACTURING CO.  
MAIN OFFICE AND PLANT ■ CLEVELAND OHIO

### USE DIRECTIONS:

Cleaning, disinfecting, sanitizing and deodorizing: Use 2 ozs. per gallon of water. For best results apply diluted Formula 235 with a cloth or mop to walls, floors and other similar hard surfaces.

At 2 ounces per gallon use-level, will provide 360 ppm of active quaternary. Effective against STAPHYLOCOCCUS AUREUS, SALMONELLA CHOLERAESUIS and ESCHERICHIA coli. Germicidal performance against above three organisms has been confirmed by the AOAC use-dilution test.

Rinse empty container thoroughly with water and discard it after use.

### DANGER

### KEEP OUT OF REACH OF CHILDREN

Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

### FIRST AID:

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink PROMPTLY a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

### NOTE TO PHYSICIAN:

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

NET CONTENTS